

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
6 May 2005 (06.05.2005)

PCT

(10) International Publication Number  
**WO 2005/040377 A2**

(51) International Patent Classification<sup>7</sup>: C12N 15/10

(74) Agent: IPRIS GMBH; Missionsstrasse 24, CH-4055 Basel (CH).

(21) International Application Number:  
PCT/EP2004/011974

(22) International Filing Date: 22 October 2004 (22.10.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/513,574 24 October 2003 (24.10.2003) US  
04002722.9 6 February 2004 (06.02.2004) EP

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(71) Applicant (*for all designated States except US*): SELEXIS S.A. [CH/CH]; 18, Chemin des Aulx, CH-1228 Plan-les-Ouates (CH).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): MERMOD, Nicolas [CH/CH]; Route des Deux-Communes 19, CH-1164 Buchillon (CH). GIROD, Pierre Alain [CH/CH]; Ch. du Muguet 15, CH-1007 Lausanne (CH). BUCHER, Philipp [CH/CH]; Avenue de Valmont 18, CH-1010 Lausanne (CH). NGUYEN, Duc-Quang [CH/CH]; Ch. des Etourneaux 24, CH-1162 Saint Prex (CH). CALABRESE, David [CH/CH]; Pl. du Tunnel 4, CH-1005 Lausanne (CH). SAUGY, Damien [CH/CH]; Avenue Mon-Loisir 13, CH-1006 Lausanne (CH). PUTTINI, Stefania [IT/CH]; Avenue Marc-Dufour 50, CH-1007 Lausanne (CH).

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Published:**

— without international search report and to be republished upon receipt of that report

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: HIGH EFFICIENCY GENE TRANSFER AND EXPRESSION IN MAMMALIAN CELLS BY A MULTIPLE TRANSFECTION PROCEDURE OF MAR SEQUENCES

(57) Abstract: The present invention relates to purified and isolated DNA sequences having protein production increasing activity and more specifically to the use of matrix attachment regions (MARs) for increasing protein production activity in a eukaryotic cell. Also disclosed is a method for the identification of said active regions, in particular MAR nucleotide sequences, and the use of these characterized active MAR sequences in a new multiple transfection method.



WO 2005/040377 A2

# HIGH EFFICIENCY GENE TRANSFER AND EXPRESSION IN MAMMALIAN CELLS BY A MULTIPLE TRANSFECTION PROCEDURE OF MAR SEQUENCES

## FIELD OF THE INVENTION

5 The present invention relates to purified and isolated DNA sequences having protein production increasing activity and more specifically to the use of matrix attachment regions (MARs) for increasing protein production activity in a eukaryotic cell. Also disclosed is a method for the identification of said active regions, in particular MAR  
10 nucleotide sequences, and the use of these characterized active MAR sequences in a new multiple transfection method.

## BACKGROUND OF THE INVENTION

15 Nowadays, the model of loop domain organization of eukaryotic chromosomes is well accepted (Boulikas T, "Nature of DNA sequences at the attachment regions of genes to the nuclear matrix", *J. Cell Biochem.*, 52:14-22, 1993). According to this model chromatin is organized in loops that span 50-100 kb attached to the nuclear matrix, a proteinaceous network made up of RNPs and other nonhistone proteins (Bode J,  
20 Stengert-Iber M, Kay V, Schalke T and Dietz-Pfeilstetter A, *Crit. Rev. Euk. Gene Exp.*, 6:115-138, 1996).

The DNA regions attached to the nuclear matrix are termed SAR or MAR for respectively scaffold (during metaphase) or matrix (interphase) attachment regions  
25 (Hart C and Laemmli U (1998), "Facilitation of chromatin dynamics by SARs" *Curr Opin Genet Dev* 8, 519-525.)

As such, these regions may define boundaries of independent chromatin domains, such that only the encompassing cis-regulatory elements control the expression of the  
30 genes within the domain.

However, their ability to fully shield a chromosomal locus from nearby chromatin elements, and thus confer position-independent gene expression, has not been seen in stably transfected cells (Poljak L, Seum C, Mattioni T and Laemmli U. (1994) "SARs stimulate but do not confer position independent gene expression", *Nucleic Acids Res*  
35 22, 4386-4394). On the other hand, MAR (or S/MAR) sequences have been shown to interact with enhancers to increase local chromatin accessibility (Jenuwein T, Forrester W, Fernandez-Herrero L, Laible G, Dull M, and Grosschedl R. (1997) "Extension of chromatin accessibility by nuclear matrix attachment regions" *Nature* 385, 269-272).  
40 Specifically, MAR elements can enhance expression of heterologous genes in cell culture lines (Kalos M and Fournier R (1995) "Position-independent transgene expression mediated by boundary elements from the apolipoprotein B chromatin domain" *Mol Cell Biol* 15,198-207), transgenic mice (Castilla J, Pintado B, Sola, I, Sanchez-Morgado J, and Enjuanes L (1998) "Engineering passive immunity in  
45 transgenic mice secreting virus-neutralizing antibodies in milk" *Nat Biotechnol* 16, 349-354) and plants (Allen G, Hall GJ, Michalowski S, Newman W, Spiker S, Weissinger A, and Thompson W (1996), "High-level transgene expression in plant cells: effects of a strong scaffold attachment region from tobacco" *Plant Cell* 8, 899-913). The utility of MAR sequences for developing improved vectors for gene therapy is also recognized  
50 (Agarwal M, Austin T, Morel F, Chen J, Bohnlein E, and Plavec I (1998), "Scaffold attachment region-mediated enhancement of retroviral vector expression in primary T

cells" *J Virol* 72, 3720-3728).

Recently, it has been shown that chromatin-structure modifying sequences including MARs, as exemplified by the chicken lysozyme 5' MAR is able to significantly enhance reporter expression in pools of stable Chinese Hamster Ovary (CHO) cells (Zahn-Zabal M, et al., "Development of stable cell lines for production or regulated expression using matrix attachment regions" *J Biotechnol*, 2001, 87(1): p. 29-42). This property was used to increase the proportion of high-producing clones, thus reducing the number of clones that need to be screened. These benefits have been observed both for constructs with MARs flanking the transgene expression cassette, as well as when constructs are co-transfected with the MAR on a separate plasmid. However, expression levels upon co-transfection with MARs were not as high as those observed for a construct in which two MARs delimit the transgene expression unit. A third and preferable process was shown to be the transfection of transgenes with MARs both linked to the transgene and on a separate plasmid (Girod et al., submitted for publication). However, one persisting limitation of this technique is the quantity of DNA that can be transfected per cell. Many multiples transfection protocols have been developed in order to achieve a high transfection efficiency to characterize the function of genes of interest. The protocol applied by Yamamoto et al, 1999 ("High efficiency gene transfer by multiple transfection protocol", *Histochem. J.* 31(4), 241-243) leads to a transfection efficiency of about 80 % after 5 transfections events, whereas the conventional transfection protocol only achieved a rate of <40%. While this technique may be useful when one wishes to increase the proportion of expressing cells, it does not lead to cells with a higher intrinsic productivity. Therefore, it cannot be used to generate high producer monoclonal cell lines. Hence, the previously described technique has two major drawbacks:

- i) this technique does not generate a homogenous population of transfected cells, since it cannot favour the integration of further gene copy, nor does it direct the transgenes to favorable chromosomal loci,
- ii) the use of the same selectable marker in multiple transfection events does not permit the selection of doubly or triply transfected cells.

In patent application WO02/074969, the utility of MARs for the development of stable eukaryotic cell lines has also been demonstrated. However, this application does not disclose neither any conserved homology for MAR DNA element nor any technique for predicting the ability for a DNA sequence to be a MAR sequence.

In fact no clear-cut MAR consensus sequence has been found (Boulikas T, "Nature of DNA sequences at the attachment regions of genes to the nuclear matrix", *J. Cell Biochem.*, 52:14-22, 1993) but evolutionarily, the structure of these sequences seem to be functionally conserved in eukaryotic genomes, since animal MARs can bind to plant nuclear scaffolds and vice versa (Mielke C, Kohwi Y, Kohwi-Shigematsu T and Bode J, "Hierarchical binding of DNA fragments derived from scaffold-attached regions: correlation of properties in vitro and function in vivo", *Biochemistry*, 29:7475-7485, 1990).

The identification of MARs by biochemical studies is a long and unpredictable process; various results can be obtained depending on the assay (Razin SV, "Functional architecture of chromosomal DNA domains", *Crit Rev Eukaryot Gene Expr.*, 6:247-269, 1996). Considering the huge number of expected MARs in a eukaryotic genome and the amount of sequences issued from genome projects, a tool able to filter potential MARs in order to perform targeted experiments would be greatly useful.

Currently two different predictive tools for MARs are available via the Internet.

The first one, MAR-Finder (<http://futuresoft.org/MarFinder>; Singh GB, Kramer JA and Krawetz SA, "Mathematical model to predict regions of chromatin attachment to the nuclear matrix", *Nucleic Acid Research*, 25:1419-1425, 1997) is based on set of patterns identified within several MARs and a statistical analysis of the co-occurrence of these patterns. MAR-Finder predictions are dependent of the sequence context, meaning that predicted MARs depend on the context of the submitted sequence. The other predictive software, SMARTest (<http://www.genomatix.de>; Frisch M, Frech K, Klingenhoff A, Cartharius K, Liebich I and Werner T, "In silico prediction of scaffold/matrix attachment regions in large genomic sequences", *Genome Research*, 12:349-354, 2001), use weight-matrices derived from experimentally identified MARs. SMARTest is said to be suitable to perform large-scale analyses. But actually aside its relative poor specificity, the amount of hypothetical MARs rapidly gets huge when doing large scale analyses with it, and in having no way to increase its specificity to restrain the number of hypothetical MARs, SMARTest becomes almost useless to screen for potent MARs from large DNA sequences.

Some other softwares, not available via the Internet, also exists; they are based as well on the frequency of MAR motifs (MRS criterion; Van Drunen CM et al., "A bipartite sequence element associated with matrix/scaffold attachment regions", *Nucleic Acids Res*, 27:2924-2930, 1999), (ChrClass; Glazko GV et al., "Comparative study and prediction of DNA fragments associated with various elements of the nuclear matrix", *Biochim. Biophys. Acta*, 1517:351-356, 2001) or based on the identification of sites of stress-induced DNA duplex (SID; Benham C and al., "Stress-induced duplex DNA destabilization in scaffold/matrix attachment regions", *J. Mol. Biol.*, 274:181-196, 1997). However, their suitability to analyze complete genome sequences remains unknown, and whether these tools may allow the identification of protein production-increasing sequences has not been reported.

Furthermore, due to the relatively poor specificity of these softwares (Frisch M, Frech K, Klingenhoff A, Cartharius K, Liebich I and Werner T, "In silico prediction of scaffold/matrix attachment regions in large genomic sequences", *Genome Research*, 12:349-354, 2001), the amount of hypothetical MARs identified in genomes rapidly gets unmanageable when doing large scale analyses, especially if most of these have no or poor activity in practice. Thus, having no way to increase prediction specificity to restrain the number of hypothetical MARs, many of the available programs become almost useless to identify potent genetic elements in view of efficiently increasing recombinant protein production.

Since all the above available predictive methods have some drawbacks that prevent large-scale analyses of genomes to identify reliably novel and potent MARs, the object of this invention is to 1) understand the functional features of MARs that allow improved recombinant protein expression; 2) get a new Bioinformatic tool compiling MAR structural features as a prediction of function, in order to 3) perform large scale analyses of genomes to identify novel and more potent MARs, and, finally 4) to demonstrate improved efficiency to increase the production of recombinant proteins from eukaryotic cells or organisms when using the newly identified MAR sequences.

## SUMMARY OF THE INVENTION

This object has been achieved by providing an improved and reliable method for the identification of DNA sequences having protein production increasing activity, in

particular MAR nucleotide sequences, and the use of these characterized active MAR sequences in a new multiple transfection method to increase the production of recombinant proteins in eukaryotic cells.

5

## BRIEF DESCRIPTION OF THE FIGURES

Fig. 1 shows the distribution plots of MARs and non-MARs sequences. Histograms are density plots (relative frequency divided by the bin width) relative to the score of the observed parameter. The density histogram for human MARs in the SMART DB database is shown in black, while the density histogram for the human chromosome 22 are in grey.

Fig. 2 shows Scatterplots of the four different criteria used by SMAR Scan® and the AT-content with human MARs from SMART DB.

Fig. 3 shows the distribution plots of MAR sequences by organism. MAR sequences from SMART DB of other organisms were retrieved and analyzed. The MAR sequences density distributions for the mouse, the chicken, the sorghum bicolor and the human are plotted jointly.

Fig. 4 shows SMAR Scan® predictions on human chromosome 22 and on shuffled chromosome 22. Top plot : Average number of hits obtained by SMAR Scan® with five: rubbed, scrambled, shuffled within nonoverlapping windows of 10 bp, order 1 Markov chains model and with the native chromosome 22. Bottom plot: Average number of MARs predicted by SMAR Scan® in five: rubbed, scrambled, shuffled within non-overlapping windows of 10 bp, order 1 Markov chains model and with the native chromosome 22.

Fig. 5 shows the dissection of the ability of the chicken lysozyme gene 5'-MAR to stimulate transgene expression in CHO-DG44 cells. Fragments B, K and F show the highest ability to stimulate transgene expression. The indicated relative strength of the elements was based on the number of high-expressor cells.

Fig. 6 shows the effect of serial-deletions of the 5'-end (upper part) and the 3'-end (lower part) of the 5'-MAR on the loss of ability to stimulate transgene expression. The transition from increased to decreased activity coincide with B-, K- and F-fragments.

Fig. 7 shows that portions of the F fragment significantly stimulate transgene expression. The F fragment regions indicated by the light grey arrow were multimerized, inserted in pGEGFP Control and transfected in CHO cells. The element that displays the highest activity is located in the central part of the element and corresponds to fragment FIII (black bar labelled minimal MAR). In addition, an enhancer activity is located in the 3'-flanking part of the FIII fragment (dark grey bar labelled MAR enhancer).

Fig. 8 shows a map of locations for various DNA sequence motifs within the *cLysMAR*. Fig. 8 (B) represents a Map of locations for various DNA sequence motifs within the *cLysMAR*. Vertical lines represent the position of the computer-predicted sites or sequence motifs along the 3034 base pairs of the *cLysMAR* and its active regions, as presented in Fig. 5. The putative transcription factor sites, (MEF2 05, Oct-1, USF-02, GATA, NFAT) for activators and (CDP, SATB1, CTCF, ARBP/MeCP2) for repressors of transcription, were identified using MatInspector (Genomatix), and CpG islands were identified with CPGPLOT. Motifs previously associated with MAR elements are labelled

in black and include CpG dinucleotides and CpG islands, unwinding motifs (AATATATT and AATATT), poly As and Ts, poly Gs and Cs, *Drosophila* topoisomerase II binding sites (GTNWAYATTNATTNATNNR) which had identity to the 6 bp core and High mobility group I (HMG-I/Y) protein binding sites. Other structural motifs include nucleosome-binding and nucleosome disfavoured sites and a motif thought to relieve the superhelical strain of DNA. Fig. 8(A) represents the comparison of the ability of portions of the cLysMAR to activate transcription with MAR prediction score profiles with MarFinder. The top diagram shows the MAR fragment activity as in Fig. 5, while the middle and bottom curves show MARFinder-predicted potential for MAR activity and for bent DNA structures respectively.

Fig. 9 shows the correlation of DNA physico-chemical properties with MAR activity. Fig. 9(A), represents the DNA melting temperature, double helix bending, major groove depth and minor groove width profiles of the 5'-MAR and were determined using the algorithms of Levitsky et al (Levitsky VG, Ponomarenko MP, Ponomarenko JV, Frolov AS, Kolchanov NA "Nucleosomal DNA property database", *Bioinformatics*, 15; 582592, 1999). The most active B, K and F fragments depicted at the top are as shown as in Figure 1. Fig. 9(B), represents the enlargement of the data presented in panel A to display the F fragment map aligned with the tracings corresponding to the melting temperature (top curve) and DNA bending (bottom curve). The position of the most active FIB fragment and protein binding site for specific transcription factors are as indicated.

Fig. 10 shows the distribution of putative transcription factor binding sites within the 5'-cLysMAR. Large arrows indicate the position of the CUE elements as identified with SMAR Scan®.

Fig. 11 shows the scheme of assembly of various portions of the MAR. The indicated portions of the cLysMAR were amplified by PCR, introducing BglII-BamHI linker elements at each extremity, and assembled to generate the depicted composite elements. For instance, the top construct consists of the assembly of all CUE and flanking sequences at their original location except that BglII-BamHI linker sequences separate each element.

Fig. 12 represents the plasmid maps.

Fig. 13 shows the effect of re-transfecting primary transfectants on GFP expression. Cells (CHO-DG44) were co-transfected with pSV40EGFP (left tube) or pMAR-SV40EGFP (central tube) and pSVneo as resistance plasmid. Cells transfected with pMAR-SV40EGFP were re-transfected 24 hours later with the same plasmid and a different selection plasmid, pSVpuro (right tube). After two weeks selection, the phenotype of the stably transfected cell population was analysed by FACS.

Fig. 14 shows the effect of multiple load of MAR-containing plasmid. The pMAR-SV40EGFP/ pMAR-SV40EGFP secondary transfectants were used in a third cycle of transfection at the end of the selection process. The tertiary transfection was accomplished with pMAR or pMAR-SV40EGFP to give tertiary transfectants. After 24 hours, cells were transfected again with either plasmid, resulting in the quaternary transfectants (see Table 4).

Fig. 15 shows comparative performance of SMAR prediction algorithms exemplified by region WP18A10A7. (A) SMAR Scan® analysis was performed with default settings. (B) SIDD analysis (top curve and left-hand side scale), and the attachment of several

DNA fragments to the nuclear matrix in vitro (bar-graph, right-hand side scale) was taken from Goetze et al ( Goetze S, Gluch A, Benham C, Bode J, "Computational and in vitro analysis of destabilized DNA regions in the interferon gene cluster: potential of predicting functional gene domains." *Biochemistry*, 42:154-166, 2003).

Fig. 16 represents the results of a a gene therapy-like protocol using MARs. The group of mice injected by MAR-network, induced from the beginning of the experiment, display a better induction of the hematocrit in comparison of mice injected by original network without MAR. After 2 months, hematocrits in "MAR-containing group" is still at values higher (65%) than normal hematocrit levels (45-55%).

Fig. 17 represents the scatterplot for the 1757 S/MAR sequences of the AT (top) and TA (bottom) dinucleotide percentages versus the predicted DNA bending as computed by SMAR Scan®.

Fig. 18 represents the dinucleotide percentage distribution plots over the 1757 non-S/MARs sequences.

Fig.19 shows the effect of various S/MAR elements on the production of recombinant green fluorescent protein (GFP). Populations of CHO cells transfected with a GFP expression vector containing or a MAR element, as indicated, were analyzed by a fluorescence-activated cell sorter (FACS®), and typical profiles are shown. The profiles display the cell number counts as a function of the GFP fluorescence levels.

Fig. 20 depicts the effect of the induction of hematocrit in mice injected by MAR-network.

## DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to a purified and isolated DNA sequence having protein production increasing activity characterized in that said DNA sequence comprises at least one bent DNA element, and at least one binding site for a DNA binding protein.

Certain sequences of DNA are known to form a relatively "static curve", where the DNA follows a particular 3-dimensional path. Thus, instead of just being in the normal B-DNA conformation ("straight"), the piece of DNA can form a flat, planar curve also defined as bent DNA (Marini, *et al.*, 1982 "Bent helical structure in kinetoplast DNA", *Proc. Natl. Acad. Sci. USA*, 79: 7664-7664).

Surprisingly, Applicants have shown that the bent DNA element of a purified and isolated DNA sequence having protein production increasing activity of the present invention usually contains at least 10% of dinucleotide TA, and/or at least 12% of dinucleotide AT on a stretch of 100 contiguous base pairs. Preferably, the bent DNA element contains at least 33% of dinucleotide TA, and/or at least 33% of dinucleotide AT on a stretch of 100 contiguous base pairs. These data have been obtained by the method described further.

According to the present invention, the purified and isolated DNA sequence usually comprises a MAR nucleotide sequence selected from the group comprising the sequences SEQ ID Nos 1 to 27 or a cLysMAR element or a fragment thereof. Preferably, the purified and isolated DNA sequence is a MAR nucleotide sequence

selected from the group comprising the sequences SEQ ID Nos 1 to 27, more preferably the sequences SEQ ID Nos 24 to 27.

5 Encompassed by the present invention are as well complementary sequences of the above-mentioned sequences SEQ ID Nos 1 to 27 and the cLysMAR element or fragment, which can be produced by using PCR or other means.

10 An "element" is a conserved nucleotide sequences that bears common functional properties (i.e. binding sites for transcription factors) or structural (i.e. bent DNA sequence) features.

15 A part of sequences SEQ ID Nos 1 to 27 and the cLysMAR element or fragment refers to sequences sharing at least 70% nucleotides in length with the respective sequence of the SEQ ID Nos 1 to 27. These sequences can be used as long as they exhibit the same properties as the native sequence from which they derive. Preferably these sequences share more than 80%, in particular more than 90% nucleotides in length with the respective sequence of the SEQ ID Nos 1 to 27.

20 The present invention also includes variants of the aforementioned sequences SEQ ID Nos 1 to 27 and the cLysMAR element or fragment, that is nucleotide sequences that vary from the reference sequence by conservative nucleotide substitutions, whereby one or more nucleotides are substituted by another with same characteristics.

25 The sequences SEQ ID Nos 1 to 23 have been identified by scanning human chromosome 1 and 2 using SMAR Scan®, showing that the identification of novel MAR sequences is feasible using the tools reported thereafter whereas SEQ ID No 24 to 27 have been identified by scanning the complete human genome using the combined SMAR Scan® method.

30 In a first step, the complete chromosome 1 and 2 were screened to identify bent DNA element as region corresponding to the highest bent, major groove depth, minor groove width and lowest melting temperature as shown in figure 3. In a second step, this collection of sequence was scanned for binding sites of regulatory proteins such as SATB1, GATA, etc. as shown in the figure 8B) yielding sequences SEQ ID 1-23.  
35 Furthermore, sequences 21-23 were further shown to be located next to known gene from the Human Genome Data Base.

40 With regard to SEQ ID No 24 to 27 these sequences have been yielded by scanning the human genome according to the combined method and were selected as examples among 1757 MAR elements so detected.

45 Molecular chimera of MAR sequences are also considered in the present invention. By molecular chimera is intended a nucleotide sequence that may include a functional portion of a MAR element and that will be obtained by molecular biology methods known by those skilled in the art.

50 Particular combinations of MAR elements or fragments or sub-portions thereof are also considered in the present invention. These fragments can be prepared by a variety of methods known in the art. These methods include, but are not limited to, digestion with restriction enzymes and recovery of the fragments, chemical synthesis or polymerase chain reactions (PCR).

Therefore, particular combinations of elements or fragments of the sequences SEQ ID

Nos 1 to 27 and cLysMAR elements or fragments are also envisioned in the present invention, depending on the functional results to be obtained. Elements of the cLysMAR are e.g. the B, K and F regions as described in WO 02/074969, the disclosure of which is hereby incorporated herein by reference, in its entirety. The preferred elements of the cLysMAR used in the present invention are the B, K and F regions. Only one element might be used or multiple copies of the same or distinct elements (multimerized elements) might be used (see Fig. 8 A)).

By fragment is intended a portion of the respective nucleotide sequence. Fragments of a MAR nucleotide sequence may retain biological activity and hence bind to purified nuclear matrices and/or alter the expression patterns of coding sequences operably linked to a promoter. Fragments of a MAR nucleotide sequence may range from at least about 100 to 1000 bp, preferably from about 200 to 700 bp, more preferably from about 300 to 500 bp nucleotides. Also envisioned are any combinations of fragments, which have the same number of nucleotides present in a synthetic MAR sequence consisting of natural MAR element and/or fragments. The fragments are preferably assembled by linker sequences. Preferred linkers are BglII-BamHI linker.

"Protein production increasing activity" refers to an activity of the purified and isolated DNA sequence defined as follows: after having been introduced under suitable conditions into a eukaryotic host cell, the sequence is capable of increasing protein production levels in cell culture as compared to a culture of cell transfected without said DNA sequence. Usually the increase is 1.5 to 10 fold, preferably 4 to 10 fold. This corresponds to a production rate or a specific cellular productivity of at least 10 pg per cell per day (see Example 11 and Fig.13).

As used herein, the following definitions are supplied in order to facilitate the understanding of this invention.

"Chromatin" is the protein and nucleic acid material constituting the chromosomes of a eukaryotic cell, and refers to DNA, RNA and associated proteins.

A "chromatin element" means a nucleic acid sequence on a chromosome having the property to modify the chromatin structure when integrated into that chromosome.

"Cis" refers to the placement of two or more elements (such as chromatin elements) on the same nucleic acid molecule (such as the same vector, plasmid or chromosome).

"Trans" refers to the placement of two or more elements (such as chromatin elements) on two or more different nucleic acid molecules (such as on two vectors or two chromosomes).

Chromatin modifying elements that are potentially capable of overcoming position effects, and hence are of interest for the development of stable cell lines, include boundary elements (BEs), matrix attachment regions (MARs), locus control regions (LCRs), and universal chromatin opening elements (UCOE).

Boundary elements ("BEs"), or insulator elements, define boundaries in chromatin in many cases (Bell A and Felsenfeld G. 1999; "Stopped at the border: boundaries and insulators, *Curr Opin Genet Dev* 9, 191-198) and may play a role in defining a transcriptional domain in vivo. BEs lack intrinsic promoter/enhancer activity, but rather are thought to protect genes from the transcriptional influence of regulatory elements in the surrounding chromatin. The enhancer-block assay is commonly used to identify

insulator elements. In this assay, the chromatin element is placed between an enhancer and a promoter, and enhancer-activated transcription is measured. Boundary elements have been shown to be able to protect stably transfected reporter genes against position effects in *Drosophila*, yeast and in mammalian cells. They have also been shown to increase the proportion of transgenic mice with inducible transgene expression.

Locus control regions ("LCRs") are cis-regulatory elements required for the initial chromatin activation of a locus and subsequent gene transcription in their native locations (Grosveld, F. 1999, "Activation by locus control regions?" *Curr Opin Genet Dev* 9, 152-157). The activating function of LCRs also allows the expression of a coupled transgene in the appropriate tissue in transgenic mice, irrespective of the site of integration in the host genome. While LCRs generally confer tissue-specific levels of expression on linked genes, efficient expression in nearly all tissues in transgenic mice has been reported for a truncated human T-cell receptor LCR and a rat LAP LCR. The most extensively characterized LCR is that of the globin locus. Its use in vectors for the gene therapy of sickle cell disease and  $\beta$ -thalassemias is currently being evaluated.

"MARs", according to a well-accepted model, may mediate the anchorage of specific DNA sequence to the nuclear matrix, generating chromatin loop domains that extend outwards from the heterochromatin cores. While MARs do not contain any obvious consensus or recognizable sequence, their most consistent feature appears to be an overall high A/T content, and C bases predominating on one strand (Bode J, Schlake T, RiosRamirez M, Mielke C, Stengart M, Kay V and KlehrWirth D, "Scaffold/matrix-attached regions: structural properties creating transcriptionally active loci", *Structural and Functional Organization of the Nuclear Matrix: International Review of Cytology*, 162A:389453, 1995). These regions have a propensity to form bent secondary structures that may be prone to strand separation. They are often referred to as base-unpairing regions (BURs), and they contain a core-unwinding element (CUE) that might represent the nucleation point of strand separation (Benham C and al., Stress induced duplex DNA destabilization in scaffold/matrix attachment regions, *J. Mol. Biol.*, 274:181-196, 1997). Several simple AT-rich sequence motifs have often been found within MAR sequences, but for the most part, their functional importance and potential mode of action remain unclear. These include the A-box (AATAAAYAAA), the T-box (TTWTTWTTWTT), DNA unwinding motifs (AATATATT, AATATT), SATB1 binding sites (H-box, A/T/C25) and consensus Topoisomerase II sites for vertebrates (RNYNNCNGYNGKTNYNY) or *Drosophila* (GTNWAYATTNATNNR).

Ubiquitous chromatin opening elements ("UCOE", also known as "ubiquitously-acting chromatin opening elements") have been reported in WO 00/05393.

An "enhancer" is a nucleotide sequence that acts to potentiate the transcription of genes independent of the identity of the gene, the position of the sequence in relation to the gene, or the orientation of the sequence. The vectors of the present invention optionally include enhancers.

A "gene" is a deoxyribonucleotide (DNA) sequence coding for a given mature protein. As used herein, the term "gene" shall not include untranslated flanking regions such as RNA transcription initiation signals, polyadenylation addition sites, promoters or enhancers.

A "product gene" is a gene that encodes a protein product having desirable characteristics such as diagnostic or therapeutic utility. A product gene includes, e. g.,

structural genes and regulatory genes.

5 A "structural gene" refers to a gene that encodes a structural protein. Examples of structural genes include but are not limited to, cytoskeletal proteins, extracellular matrix proteins, enzymes, nuclear pore proteins and nuclear scaffold proteins, ion channels and transporters, contractile proteins, and chaperones. Preferred structural genes encode for antibodies or antibody fragments.

10 A "regulatory gene" refers to a gene that encodes a regulatory protein. Examples of regulatory proteins include, but are not limited to, transcription factors, hormones, growth factors, cytokines, signal transduction molecules, oncogenes, proto-oncogenes, transmembrane receptors, and protein kinases.

15 "Orientation" refers to the order of nucleotides in a given DNA sequence. For example, an inverted orientation of a DNA sequence is one in which the 5' to 3' order of the sequence in relation to another sequence is reversed when compared to a point of reference in the DNA from which the sequence was obtained. Such reference points can include the direction of transcription of other specified DNA sequences in the source DNA and/or the origin of replication of replicable vectors containing the  
20 sequence.

"Eukaryotic cell" refers to any mammalian or non-mammalian cell from a eukaryotic organism. By way of non-limiting example, any eukaryotic cell that is capable of being maintained under cell culture conditions and subsequently transfected would be  
25 included in this invention. Especially preferable cell types include, e. g., stem cells, embryonic stem cells, Chinese hamster ovary cells (CHO), COS, BHK21, NIH3T3, HeLa, C2C12, cancer cells, and primary differentiated or undifferentiated cells. Other suitable host cells are known to those skilled in the art.

30 The terms "host cell" and "recombinant host cell" are used interchangeably herein to indicate a eukaryotic cell into which one or more vectors of the invention have been introduced. It is understood that such terms refer not only to the particular subject cell but also to the progeny or potential progeny of such a cell. Because certain modifications may occur in succeeding generations due to either mutation or  
35 environmental influences, such progeny may not, in fact, be identical to the parent cell, but are still included within the scope of the term as used herein.

40 The terms "introducing a purified DNA into a eukaryotic host cell" or "transfection" denote any process wherein an extracellular DNA, with or without accompanying material, enters a host cell. The term "cell transfected" or "transfected cell" means the cell into which the extracellular DNA has been introduced and thus harbours the extracellular DNA. The DNA might be introduced into the cell so that the nucleic acid is replicable either as a chromosomal integrant or as an extra chromosomal element.

45 "Promoter" as used herein refers to a nucleic acid sequence that regulates expression of a gene.

50 "Co-transfection" means the process of transfecting a eukaryotic cell with more than one exogenous gene, or vector, or plasmid, foreign to the cell, one of which may confer a selectable phenotype on the cell.

The purified and isolated DNA sequence having protein production increasing activity also comprises, besides one or more bent DNA element, at least one binding site for a DNA binding protein.

- 5 Usually the DNA binding protein is a transcription factor. Examples of transcription factors are the group comprising the polyQpolyP domain proteins.  
Another example of a transcription factor is a transcription factor selected from the group comprising SATB1, NMP4, MEF2, S8, DLX1, FREAC7, BRN2, GATA 1/3, TATA, Bright, MSX, AP1, C/EBP, CREBP1, FOX, Freac7, HFH1, HNF3alpha, Nkx25,  
10 POU3F2, Pit1, TTF1, XFD1, AR, C/EBPgamma, Cdc5, FOXD3, HFH3, HNF3 beta, MRF2, Oct1, POU6F1, SRF, V\$MTATA\_B, XFD2, Bach2, CDP CR3, Cdx2, FOXJ2, HFL, HP1, Myc, PBX, Pax3, TEF, VBP, XFD3, Brn2, COMP1, Evl, FOXP3, GATA4, HFN1, Lhx3, NKX3A, POU1F1, Pax6, TFIIA or a combination of two or more of these transcription factors are preferred. Most preferred are SATB1, NMP4, MEF2 and  
15 polyQpolyP domain proteins.

SATB1, NMP4 and MEF2, for example, are known to regulate the development and/or tissue-specific gene expression in mammals. These transcription factors have the capacity to alter DNA geometry, and reciprocally, binding to DNA as an allosteric ligand  
20 modifies their structure. Recently, SATB1 was found to form a cage-like structure circumscribing heterochromatin (Cai S, Han HJ, and Kohwi-Shigematsu T, "Tissue-specific nuclear architecture and gene expression regulated by SATB1" *Nat Genet*, 2003. 34(1): p. 42-51).

- 25 Yet another object of the present invention is to provide a purified and isolated cLysMAR element and/or fragment, a sequence complementary thereof, a part thereof sharing at least 70% nucleotides in length, a molecular chimera thereof, a combination thereof and variants.

- 30 More preferably, the cLysMAR element and/or fragment are consisting of at least one nucleotide sequence selected from the B, K and F regions.

A further object of the present invention is to provide a synthetic MAR sequence comprising natural MAR element and/or fragments assembled between linker  
35 sequences.

- Preferably, the synthetic MAR sequence comprises a cLysMAR element and/or fragment a sequence complementary thereof, a part thereof sharing at least 70% nucleotides in length, a molecular chimera thereof, a combination thereof and variants.  
40 Also preferably, linker sequences are BgIII-BamHI linker.

An other aspect of the invention is to provide a method for identifying a MAR sequence using a Bioinformatic tool comprising the computing of values of one or more DNA sequence features corresponding to DNA bending, major groove depth and minor  
45 groove width potentials and melting temperature. Preferably, the identification of one or more DNA sequence features further comprises a further DNA sequence feature corresponding to binding sites for DNA binding proteins, which is also computed with this method.

- 50 Preferably, profiles or weight-matrices of said bioinformatic tool are based on dinucleotide recognition.

The bioinformatic tool used for the present method is preferably, SMAR Scan®, which contains algorithms developed by Gene Express ([http://srs6.bionet.nsc.ru/srs6bin/cgi-bin/wgetz?-e+\[FEATURES-SiteID:'nR'\]](http://srs6.bionet.nsc.ru/srs6bin/cgi-bin/wgetz?-e+[FEATURES-SiteID:'nR'])) and based on Levitsky *et al.*, 1999. These algorithms recognise profiles, based on dinucleotides weight-matrices, to compute the theoretical values for conformational and physicochemical properties of DNA.

Preferably, SMAR Scan® uses the four theoretical criteria also designated as DNA sequence features corresponding to DNA bending, major groove depth and minor groove width potentials, melting temperature in all possible combination, using scanning windows of variable size (see Fig. 3). For each function used, a cut-off value has to be set. The program returns a hit every time the computed score of a given region is above the set cut-off value for all of the chosen criteria. Two data output modes are available to handle the hits, the first (called "profile-like") simply returns all hit positions on the query sequence and their corresponding values for the different criteria chosen. The second mode (called "contiguous hits") returns only the positions of several contiguous hits and their corresponding sequence. For this mode, the minimum number of contiguous hits is another cut-off value that can be set, again with a tunable window size. This second mode is the default mode of SMAR Scan®. Indeed, from a semantic point of view, a hit is considered as a core-unwinding element (CUE), and a cluster of CUEs accompanied by clusters of binding sites for relevant proteins is considered as a MAR. Thus, SMAR Scan® considers only several contiguous hits as a potential MAR.

To tune the default cut-off values for the four theoretical structural criteria, experimentally validated MARs from SMARt DB (<http://transfac.gbf.de/-> SMARt DB) were used. All the human MAR sequences from the database were retrieved and analyzed with SMAR Scan® using the "profile-like" mode with the four criteria and with no set cut-off value. This allowed the setting of each function for every position of the sequences. The distribution for each criterion was then computed according to these data (see Fig. 1 and 3).

The default cut-off values of SMAR Scan® for the bend, the major groove depth and the minor groove width were set at the average of the 75th quantile and the median. For the melting temperature, the default cut-off value should be set at the 75th quantile. The minimum length for the "contiguous-hits" mode should be set to 300 because it is assumed to be the minimum length of a MAR (see Fig. 8 and 9). However, one skilled in the art would be able to determine the cut-off values for the above-mentioned criteria for a given organism with minimal experimentation.

Preferably, DNA bending values are comprised between 3 to 5 ° (radial degree). Most preferably they are situated between 3.8 to 4.4 °, corresponding to the smallest peak of Fig. 1.

Preferably the major groove depth values are comprised between 8.9 to 9.3 Å (Angström) and minor groove width values between 5.2 to 5.8 Å. Most preferably the major groove depth values are comprised between 9.0 to 9.2 Å and minor groove width values between 5.4 to 5.7 Å.

Preferably the melting temperature is comprised between 55 to 75 ° C (Celsius degree). Most preferably, the melting temperature is comprised between 55 to 62 ° C.

The DNA binding protein of which values can be computed by the method is usually a transcription factor preferably a polyQpolyP domain or a transcription factor selected

from the group comprising SATB1, NMP4, MEF2, S8, DLX1, FREAC7, BRN2, GATA 1/3, TATA, Bright, MSX, AP1, C/EBP, CREBP1, FOX, Freac7, HFH1, HNF3alpha, Nkx25, POU3F2, Pit1, TTF1, XFD1, AR, C/EBPgamma, Cdc5, FOXD3, HFH3, HNF3 beta, MRF2, Oct1, POU6F1, SRF, V\$MTATA\_B, XFD2, Bach2, CDP CR3, Cdx2, FOXJ2, HFL, HP1, Myc, PBX, Pax3, TEF, VBP, XFD3, Brn2, COMP1, Evil, FOXP3, GATA4, HFN1, Lhx3, NKX3A, POU1F1, Pax6, TFIIA or a combination of two or more of these transcription factors.

However, one skilled in the art would be able to determine other kinds of transcription factors in order to carry out the method according to the present invention.

In case SMAR Scan® is envisaged to perform, for example, large scale analysis, then, preferably, the above-mentioned method further comprises at least one filter predicting DNA binding sites for DNA transcription factors in order to reduce the computation.

The principle of this method combines SMAR Scan® to compute the structural features as described above and a filter, such as for example, the pfsearch, (from the pftools package as described in Bucher P, Karplus K, Moeri N, and Hofmann K, "A flexible search technique based on generalized profiles", *Computers and Chemistry*, 20:324, 1996) to predict the binding of some transcription factors.

Examples of filters comprise, but are not limited to, pfsearch, MatInspector, RMatch Professional and TRANSFAC Professional

This combined method uses the structural features of SMAR Scan® and the predicted binding of specific transcription factors of the filter that can be applied sequentially in any order to select MARs, therefore, depending on the filter is applied at the beginning or at the end of the method.

The first level selects sequences out of the primary input sequence and the second level, consisting in the filter, may be used to restrain among the selected sequences those which satisfy the criteria used by the filter.

In this combined method the filter detects clusters of DNA binding sites using profiles or weightmatrices from, for example, MatInspector (Quandt K, Frech K, Karas H, Wingender E, Werner T, "MatInd and MatInspector New fast and versatile tools for detection of consensus matches in nucleotide sequence data", *Nucleic Acids Research*, 23, 48784884, 1995.). The filter can also detect densities of clusters of DNA binding sites.

The combined method is actually a "wrapper" written in Perl for SMAR Scan® and, in case the pfsearch is used as a filter, from the pftools. The combined method performs a twolevel processing using at each level one of these tools (SMAR Scan® or filter) as a potential "filter", each filter being optional and possible to be used to compute the predicted features without doing any filtering.

If SMAR Scan® is used in the first level to filter subsequences, it has to be used with the "all the contiguous hits" mode in order to return sequences. If the pfsearch is used in the first level as first filter, it has to be used with only one profile and a distance in nucleotide needs to be provided. This distance is used to group together pfsearch hits that are located at a distance inferior to the distance provided in order to return sequences; The combined method launches pfsearch, parses its output and returns

sequences corresponding to pfsearch hits that are grouped together according to the distance provided. Then whatever the tool used in the first level, the length of the sub-sequences thus selected can be systematically extended at both ends according to a parameter called "hits extension".

5 The second and optional level can be used to filter out sequences (already filtered sequences or unfiltered input sequences) or to get the results of SMAR Scan® and/or pfsearch without doing any filtering on these sequences. If the second level of combined method is used to filter, for each criteria considered cutoff values (hit per  
10 nucleotide) need to be provided to filter out those sequences (see Fig. 20).

Another concern of the present invention is also to provide a method for identifying a MAR sequence comprising at least one filter detecting clusters of DNA binding sites using profiles or weightmatrices. Preferably, this method comprises two levels of filters  
15 and in this case, SMAR Scan® is totally absent from said method. Usually, the two levels consist in pfsearch.

Also embraced by the present invention is a purified and isolated MAR DNA sequence identifiable according to the method for identifying a MAR sequence using the  
20 described bioinformatic tool, the combined method or the method comprising at least one filter.

Analysis by the combined method of the whole human genome yielded a total of 1757 putative MARs representing a total of 1 065 305 base paires. In order to reduce the number of results, a dinucleotide analysis was performed on these 1757 MARs,  
25 computing each of the 16 possible dinucleotide percentage for each sequence considering both strands in the 5' to 3' direction.

Surprisingly, Applicants have shown that all of the "super" MARs detected with the combined method contain at least 10% of dinucleotide TA on a stretch of 100  
30 contiguous base pairs. Preferably, these sequences contain at least 33% of dinucleotide TA on a stretch of 100 contiguous base pairs.

Applicants have also shown that these same sequences further contain at least 12% of dinucleotide AT on a stretch of 100 contiguous base pairs. Preferably, they contain at  
35 least 33% of dinucleotide AT on a stretch of 100 contiguous base pairs.

An other aspect of the invention is to provide a purified and isolated MAR DNA sequence of any of the preceding described MARs, comprising a sequence selected from the sequences SEQ ID Nos 1 to 27, a sequence complementary thereof, a part  
40 thereof sharing at least 70% nucleotides in length, a molecular chimera thereof, a combination thereof and variants.

Preferably, said purified and isolated MAR DNA sequence comprises a sequence selected from the sequences SEQ ID Nos 24 to 27, a sequence complementary thereof, a part thereof sharing at least 70% nucleotides in length, a molecular chimera  
45 thereof, a combination thereof and variants. These sequences 24 to 27 correspond to those detected by the combined method and show a higher protein production increasing activity over sequences 1 to 23.

The present invention also encompasses the use of a purified and isolated DNA  
50 sequence comprising a first isolated matrix attachment region (MAR) nucleotide sequence which is a MAR nucleotide sequence selected from the group comprising

- a purified and isolated DNA sequence having protein production increasing activity,
- a purified and isolated MAR DNA sequence identifiable according to the method for identifying a MAR sequence using the described bioinformatic tool, the combined method or the method comprising at least one filter,
- the sequences SEQ ID Nos 1 to 27,
- a purified and isolated cLysMAR element and/or fragment,
- a synthetic MAR sequence comprising natural MAR element and/or fragments assembled between linker sequences,

a sequence complementary thereof, a part thereof sharing at least 70% nucleotides in length, a molecular chimera thereof, a combination thereof and variants or a MAR nucleotide sequence of a cLysMAR element and/or fragment, a sequence complementary thereof, a part thereof sharing at least 70% nucleotides in length, a molecular chimera thereof, a combination thereof and variants for increasing protein production activity in a eukaryotic host cell.

Said purified and isolated DNA sequence usually further comprises one or more regulatory sequences, as known in the art e.g. a promoter and/or an enhancer, polyadenylation sites and splice junctions usually employed for the expression of the protein or may optionally encode a selectable marker. Preferably said purified and isolated DNA sequence comprises a promoter which is operably linked to a gene of interest.

The DNA sequences of this invention can be isolated according to standard PCR protocols and methods well known in the art.

Promoters which can be used provided that such promoters are compatible with the host cell are, for example, promoters obtained from the genomes of viruses such as polyoma virus, adenovirus (such as Adenovirus 2), papilloma virus (such as bovine papilloma virus), avian sarcoma virus, cytomegalovirus (such as murine or human cytomegalovirus immediate early promoter), a retrovirus, hepatitis-B virus, and Simian Virus 40 (such as SV 40 early and late promoters) or promoters obtained from heterologous mammalian promoters, such as the actin promoter or an immunoglobulin promoter or heat shock promoters. Such regulatory sequences direct constitutive expression.

Furthermore, the purified and isolated DNA sequence might further comprise regulatory sequences which are capable of directing expression of the nucleic acid preferentially in a particular cell type (e. g., tissue-specific regulatory elements are used to express the nucleic acid). Tissue-specific regulatory elements are known in the art. Non-limiting examples of suitable tissue-specific promoters include the albumin promoter (liver-specific; Pinkert, et al., 1987. *Genes Dev.* 1: 268-277), lymphoid-specific promoters (Calame and Eaton, 1988. *Adv. Immunol.* 43: 235-275), in particular promoters of T cell receptors (Winoto and Baltimore, 1989. *EMBOJ.* 8: 729-733) and immunoglobulins (Banerji, et al., 1983. *Cell* 33: 729-740; Queen and Baltimore, 1983. *Cell* 33:741-748), neuron-specific promoters (e. g., the neurofilament promoter; Byrne and Ruddle, 1989. *Proc. Natl. Acad. Sci. USA* 86: 5473-5477), pancreas-specific promoters (Edlund, et al., 1985. *Science* 230: 912-916), and mammary gland-specific promoters (e. g., milk whey promoter; U. S. Pat. No. 4,873,316 and European Application No. 264,166).

Developmentally-regulated promoters are also encompassed. Examples of such promoters include, e.g., the murine hox promoters (Kessel and Gruss, 1990. *Science* 249: 374-379) and thea-fetoprotein promoter (Campes and Tilghman, 1989. *Genes*

Dev. 3: 537-546).

Regulatable gene expression promoters are well known in the art, and include, by way of non-limiting example, any promoter that modulates expression of a gene encoding a desired protein by binding an exogenous molecule, such as the CRE/LOX system, the TET system, the doxycycline system, the NFkappaB/UV light system, the Leu3p/isopropylmalate system, and the GLVPc/GAL4 system (See e. g., Sauer, 1998, Methods 14 (4): 381-92; Lewandoski, 2001, Nat. Rev. Genet 2 (10): 743-55; Legrand-Poels et al., 1998, J. Photochem. Photobiol. B. 45: 18; Guo et al., 1996, FEBS Lett. 390 (2): 191-5; Wang et al., PNAS USA, 1999, 96 (15): 84838). However, one skilled in the art would be able to determine other kinds of promoters that are suitable in carrying out the present invention.

Enhancers can be optionally included in the purified DNA sequence of the invention then belonging to the regulatory sequence, e.g. the promoter.

The "gene of interest" or "transgene" preferably encodes a protein (structural or regulatory protein). As used herein "protein" refers generally to peptides and polypeptides having more than about ten amino acids. The proteins may be "homologous" to the host (i.e., endogenous to the host cell being utilized), or "heterologous," (i.e., foreign to the host cell being utilized), such as a human protein produced by yeast. The protein may be produced as an insoluble aggregate or as a soluble protein in the periplasmic space or cytoplasm of the cell, or in the extracellular medium. Examples of proteins include hormones such as growth hormone or erythropoietin (EPO), growth factors such as epidermal growth factor, analgesic substances like enkephalin, enzymes like chymotrypsin, receptors to hormones or growth factors, antibodies and include as well proteins usually used as a visualizing marker e.g. green fluorescent protein.

Preferably the purified DNA sequence further comprises at least a second isolated matrix attachment region (MAR) nucleotide sequence selected from the group comprising

- a purified and isolated DNA sequence having protein production increasing activity,
- a purified and isolated MAR DNA sequence identifiable according to the method for identifying a MAR sequence using the described bioinformatic tool, the combined method or the method comprising at least one filter,
- the sequences SEQ ID Nos 1 to 27,
- a purified and isolated cLysMAR element and/or fragment,
- a synthetic MAR sequence comprising natural MAR element and/or fragments assembled between linker sequences,

a sequence complementary thereof, a part thereof sharing at least 70% nucleotides in length, a molecular chimera thereof, a combination thereof and variants. The isolated matrix attachment region (MAR) nucleotide sequence might be identical or different.

Alternatively, a first and a second identical MAR nucleotide sequence are used.

Preferably, the MAR nucleotide sequences are located at both the 5' and the 3' ends of the sequence containing the promoter and the gene of interest. But the invention also envisions the fact that said first and or at least second MAR nucleotide sequences are located on a sequence distinct from the one containing the promoter and the gene of interest.

Embraced by the scope of the present invention is also the purified and isolated DNA sequence comprising a first isolated matrix attachment region (MAR) nucleotide sequence which is a MAR nucleotide sequence selected from the group comprising

- a purified and isolated DNA sequence having protein production increasing activity,
- a purified and isolated MAR DNA sequence identifiable according to the method for identifying a MAR sequence using the described bioinformatic tool, the combined method or the method comprising at least one filter,
- the sequences SEQ ID Nos 1 to 27,
- a purified and isolated cLysMAR element and/or fragment,
- a synthetic MAR sequence comprising natural MAR element and/or fragments assembled between linker sequences,

a sequence complementary thereof, a part thereof sharing at least 70% nucleotides in length, a molecular chimera thereof, a combination thereof and variants that can be used for increasing protein production activity in a eukaryotic host cell by introducing the purified and isolated DNA sequence into a eukaryotic host cell according to well known protocols. Usually applied methods for introducing DNA into eukaryotic host cells applied are e.g. direct introduction of cloned DNA by microinjection or microparticle bombardment; electrotransfer; use of viral vectors; encapsulation within a carrier system; and use of transfecting reagents such as calcium phosphate, diethylaminoethyl (DEAE) –dextran or commercial transfection systems like the Lipofect-AMINE 2000 (Invitrogen). Preferably, the transfection method used to introduce the purified DNA sequence into a eukaryotic host cell is the method for transfecting a eukaryotic cell as described below.

The purified and isolated DNA sequence can be used in the form of a circular vector. Preferably, the purified and isolated DNA sequence is used in the form of a linear DNA sequence as vector.

As used herein, "plasmid" and "vector" are used interchangeably, as the plasmid is the most commonly used vector form. However, the invention is intended to include such other forms of expression vectors, including, but not limited to, viral vectors (e. g., replication defective retroviruses, adenoviruses and adeno-associated viruses), which serve equivalent functions.

The present invention further encompasses a method for transfecting a eukaryotic host cell, said method comprising

- a) introducing into said eukaryotic host cell at least one purified DNA sequence comprising at least one DNA sequence of interest and/or at least one purified and isolated DNA sequence comprising a MAR nucleotide sequence or other chromatin modifying elements,
- b) subjecting within a defined time said transfected eukaryotic host cell to at least one additional transfection step with at least one purified DNA sequence comprising at least one DNA sequence of interest and/or with at least one purified and isolated DNA sequence comprising a MAR nucleotide sequence or other chromatin modifying elements
- c) selecting said transfected eukaryotic host cell.

Preferably at least two up to four transfecting steps are applied in step b).

In order to select the successful transfected cells, a gene that encodes a selectable marker (e. g., resistance to antibiotics) is generally introduced into the host cells along with the gene of interest. The gene that encodes a selectable marker might be located

on the purified DNA sequence comprising at least one DNA sequence of interest and/or at least one purified and isolated DNA sequence consisting of a MAR nucleotide sequence or other chromatin modifying elements or might optionally be co-introduced in separate form e.g. on a plasmid. Various selectable markers include those that confer resistance to drugs, such as G418, hygromycin and methotrexate. The amount of the drug can be adapted as desired in order to increase productivity

Usually, one or more selectable markers are used. Preferably, the selectable markers used in each distinct transfection steps are different. This allows selecting the transformed cells that are "multi-transformed" by using for example two different antibiotic selections.

Any eukaryotic host cell capable of protein production and lacking a cell wall can be used in the methods of the invention. Examples of useful mammalian host cell lines include human cells such as human embryonic kidney line (293 or 293 cells subcloned for growth in suspension culture, Graham et al., J. Gen Virol 36, 59 (1977)), human cervical carcinoma cells (HELA, ATCC CCL 2), human lung cells (W138, ATCC CCL 75), human liver cells (Hep G2, HB 8065); rodent cells such as baby hamster kidney cells (BHK, ATCC CCL 10), Chinese hamster ovary cells/-DHFR (CHO, Urlaub and Chasin, *Proc. Natl. Acad. Sci. USA*, 77, 4216 (1980)), mouse sertoli cells (TM4, Mather, *Biol. Reprod* 23, 243-251 (1980)), mouse mammary tumor (MMT 060562, ATCC CCL51); and cells from other mammals such as monkey kidney CV1 line transformed by SV40 (COS-7, ATCC CRL 1651); monkey kidney cells (CV1 ATCC CCL 70); African green monkey kidney cells (VERO-76, ATCC CRL-1587); canine kidney cells (MDCK, ATCC CCL 34); buffalo rat liver cells (BRL 3A, ATCC CRL 1442); myeloma (e.g. NS0) /hybridoma cells.

Preferably, the selected transfected eukaryotic host cells are high protein producer cells with a production rate of at least 10 pg per cell per day.

Most preferred for uses herein are mammalian cells, more preferred are CHO cells.

The DNA sequence of interest of the purified and isolated DNA sequence is usually a gene of interest preferably encoding a protein operably linked to a promoter as described above. The purified and isolated DNA sequence comprising at least one DNA sequence of interest might comprise additionally to the DNA sequence of interest MAR nucleotide sequence or other chromatin modifying elements.

Purified and isolated DNA sequence comprising a MAR nucleotide sequence are for example selected from the group comprising the sequences SEQ ID Nos 1 to 27 and/or particular elements of the cLysMAR e.g. the B, K and F regions as well as fragment and elements and combinations thereof as described above. Other chromatin modifying elements are for example boundary elements (BEs), locus control regions (LCRs), and universal chromatin opening elements (UCOEs) (see Zahn-Zabal et al. already cited). An example of multiple transfections of host cells is shown in Example 12 (Table 3).

The first transfecting step (primary transfection) is carried out with the gene of interest (SV40EGFP) alone, with a MAR nucleotide sequence (MAR) alone or with the gene of interest and a MAR nucleotide sequence (MAR-SV40EGFP). The second transfecting step (secondary transfection) is carried out with the gene of interest (SV40EGFP) alone, with a MAR nucleotide sequence (MAR) alone or with the gene of interest and a MAR nucleotide sequence (MAR-SV40EGFP), in all possible combinations resulting from the first transfecting step.

Preferably the eukaryotic host cell is transfected by:

- a) introducing a purified DNA sequence comprising one DNA sequence of interest and additionally a MAR nucleotide sequence,  
 b) subjecting within a defined time said transfected eukaryotic host cell to at least one additional transfection step with the same purified DNA sequence comprising one DNA sequence of interest and additionally a MAR nucleotide sequence of step a).

Also preferably, the MAR nucleotide sequence of the of the purified and isolated DNA sequence is selected from the group comprising

- a purified and isolated DNA sequence having protein production increasing activity,
- a purified and isolated MAR DNA sequence identifiable according to the method for identifying a MAR sequence using the described bioinformatic tool, the combined method or the method comprising at least one filter,
- the sequences SEQ ID Nos 1 to 27,
- a purified and isolated cLysMAR element and/or fragment,
- a synthetic MAR sequence comprising natural MAR element and/or fragments assembled between linker sequences,

a sequence complementary thereof, a part thereof sharing at least 70% nucleotides in length, a molecular chimera thereof, a combination thereof and variants.

Surprisingly, a synergy between the first and second transfection has been observed. A particular synergy has been observed when MAR elements are present at one or both of the transfection steps. Multiple transfections of the cells with pMAR alone or in combination with various expression plasmids, using the method described above have been carried out. For example, Table 3 shows that transfecting the cells twice with the pMAR-SV40EGFP plasmid gave the highest expression of GFP and the highest degree of enhancement of all conditions (4.3 fold). In contrast, transfecting twice the vector without MAR gave little or no enhancement, 2.8-fold, instead of the expected two-fold increase. This proves that the presence of MAR elements at each transfection step is of particular interest to achieve the maximal protein synthesis.

As a particular example of the transfection method, said purified DNA sequence comprising at least one DNA sequence of interest can be introduced in form of multiple unlinked plasmids, comprising a gene of interest operably linked to a promoter, a selectable marker gene, and/or protein production increasing elements such as MAR sequences.

The ratio of the first and subsequent DNA sequences may be adapted as required for the use of specific cell types, and is routine experimentation to one ordinary skilled in the art.

The defined time for additional transformations of the primary transformed cells is tightly dependent on the cell cycle and on its duration. Usually the defined time corresponds to intervals related to the cell division cycle.

Therefore this precise timing may be adapted as required for the use of specific cell types, and is routine experimentation to one ordinary skilled in the art.

Preferably the defined time is the moment the host cell just has entered into the same phase of a second or a further cell division cycle, preferably the second cycle.

This time is usually situated between 6h and 48 h, preferably between 20h and 24h after the previous transfecting event.

Also encompassed by the present invention is a method for transfecting a eukaryotic host cell, said method comprising co-transfecting into said eukaryotic host cell at least one first purified and isolated DNA sequence comprising at least one DNA sequence of

interest, and a second purified DNA comprising at least one MAR nucleotide selected from the group comprising:

- a purified and isolated DNA sequence having protein production increasing activity,
- a purified and isolated MAR DNA sequence identifiable according to the method for identifying a MAR sequence using the described bioinformatic tool, the combined method or the method comprising at least one filter,
- the sequences SEQ ID Nos 1 to 27,
- a purified and isolated cLysMAR element and/or fragment,
- a synthetic MAR sequence comprising natural MAR element and/or fragments assembled between linker sequences,

a sequence complementary thereof, a part thereof sharing at least 70% nucleotides in length, a molecular chimera thereof, a combination thereof and variants.

Said first purified and isolated DNA sequence can also comprise at least one MAR nucleotide as described above.

Also envisioned is a process for the production of a protein wherein a eukaryotic host cell is transfected according to the transfection methods as defined in the present invention and is cultured in a culture medium under conditions suitable for expression of the protein. Said protein is finally recovered according to any recovering process known to the skilled in the art.

Given as an example, the following process for protein production might be used.

The eukaryotic host cell transfected with the transfection method of the present invention is used in a process for the production of a protein by culturing said cell under conditions suitable for expression of said protein and recovering said protein. Suitable culture conditions are those conventionally used for in vitro cultivation of eukaryotic cells as described e.g. in WO 96/39488. The protein can be isolated from the cell culture by conventional separation techniques such as e.g. fractionation on immunoaffinity or ion-exchange columns; precipitation; reverse phase HPLC; chromatography; chromatofocusing; SDS-PAGE; gel filtration. One skilled in the art will appreciate that purification methods suitable for the polypeptide of interest may require modification to account for changes in the character of the polypeptide upon expression in recombinant cell culture.

The proteins that are produced according to this invention can be tested for functionality by a variety of methods. For example, the presence of antigenic epitopes and ability of the proteins to bind ligands can be determined by Western blot assays, fluorescence cell sorting assays, immunoprecipitation, immunochemical assays and/or competitive binding assays, as well as any other assay which measures specific binding activity.

The proteins of this invention can be used in a number of practical applications including, but not limited to:

1. Immunization with recombinant host protein antigen as a viral/pathogen antagonist.
2. Production of membrane proteins for diagnostic or screening assays.
3. Production of membrane proteins for biochemical studies.
4. Production of membrane protein for structural studies.
5. Antigen production for generation of antibodies for immuno-histochemical mapping, including mapping of orphan receptors and ion channels.

Also provided by the present invention is a eukaryotic host cell transfected according to any of the preceding transfection methods. Preferably, the eukaryotic host cell is a mammalian host cell line.

As already described, example of useful mammalian host cell lines include human cells such as human embryonic kidney line (293 or 293 cells subcloned for growth in suspension culture, Graham et al., *J. Gen Virol* 36, 59 (1977)), human cervical carcinoma cells (HELA, ATCC CCL 2), human lung cells (W138, ATCC CCL 75), human liver cells (Hep G2, HB 8065); rodent cells such as baby hamster kidney cells (BHK, ATCC CCL 10), Chinese hamster ovary cells/-DHFR (CHO, Urlaub and Chasin, *Proc. Natl. Acad. Sci. USA*, 77, 4216 (1980)), mouse sertoli cells (TM4, Mather, *Biol. Reprod* 23, 243-251 (1980)), mouse mammary tumor (MMT 060562, ATCC CCL51); and cells from other mammals such as monkey kidney CV1 line transformed by SV40 (COS-7, ATCC CRL 1651); monkey kidney cells (CV1 ATCC CCL 70); African green monkey kidney cells (VERO-76, ATCC CRL-1587); canine kidney cells (MDCK, ATCC CCL 34); buffalo rat liver cells (BRL 3A, ATCC CRL 1442); myeloma (e.g. NS0) /hybridoma cells.

Most preferred for uses herein are CHO cells.

The present invention also provides for a cell transfection mixture or Kit comprising at least one purified and isolated DNA sequence according to the invention.

The invention further comprises a transgenic organism wherein at least some of its cells have stably incorporated at least one DNA sequence of

- a purified and isolated DNA sequence having protein production increasing activity,
- a purified and isolated MAR DNA sequence identifiable according to the method for identifying a MAR sequence using the described bioinformatic tool, the combined method or the method comprising at least one filter,
- the sequences SEQ ID Nos 1 to 27,
- a purified and isolated cLysMAR element and/or fragment,
- a synthetic MAR sequence comprising natural MAR element and/or fragments assembled between linker sequences,

a sequence complementary thereof, a part thereof sharing at least 70% nucleotides in length, a molecular chimera thereof, a combination thereof and variants.

Preferably, some of the cells of the transgenic organisms have been transfected according the methods described herein.

Also envisioned in the present invention is a transgenic organism wherein its genome has stably incorporated at least one DNA sequence of

- a purified and isolated DNA sequence having protein production increasing activity,
- a purified and isolated MAR DNA sequence identifiable according to the method for identifying a MAR sequence using the described bioinformatic tool, the combined method or the method comprising at least one filter,
- the sequences SEQ ID Nos 1 to 27,
- a purified and isolated cLysMAR element and/or fragment,
- a synthetic MAR sequence comprising natural MAR element and/or fragments assembled between linker sequences,

a sequence complementary thereof, a part thereof sharing at least 70% nucleotides in length, a molecular chimera thereof, a combination thereof and variants.

Transgenic eukaryotic organisms which can be useful for the present invention are for example selected from the group comprising mammals (mouse, human, monkey etc) and in particular laboratory animals such as rodents in general, insects (drosophila,

etc), fishes (zebra fish, etc.), amphibians (frogs, newt, etc..) and other simpler organisms such as c. elegans, yeast, etc....

5 Yet another object of the present invention is to provide a computer readable medium comprising computer-executable instructions for performing the method for identifying a MAR sequence as described in the present invention.

10 The foregoing description will be more fully understood with reference to the following Examples. Such Examples, are, however, exemplary of methods of practising the present invention and are not intended to limit the scope of the invention.

## EXAMPLES

### Example 1: SMAR Scan® and MAR sequences

A first rough evaluation of SMAR Scan® was done by analyzing experimentally defined human MARs and non-MAR sequences. As MAR sequences, the previous results from the analysis of human MARs from SMART Db were used to plot a density histogram for each criterion as shown in Fig. 1. Similarly, non-MAR sequences were also analyzed and plotted. As non-MAR sequences, all Ref-Seq-contigs from the chromosome 22 were used, considering that this latter was big enough to contain a negligible part of MAR sequences regarding the part of non-MAR sequences.

The density distributions shown in Fig. 1 are all skewed with a long tail. For the highest bend, the highest major groove depth and the highest minor groove width, the distributions are right skewed. For the lowest melting temperature, the distributions are left-skewed which is natural given the inverse correspondence of this criterion regarding the three others. For the MAR sequences, biphasic distributions with a second weak peak, are actually apparent. And between MAR and non-MAR sequences distributions, a clear shift is also visible in each plot.

Among all human MAR sequences used, in average only about 70% of them have a value greater than the 75th quantile of human MARs distribution, this for the four different criteria. Similarly concerning the second weak peak of each human MARs distribution, only 15% of the human MAR sequences are responsible of these outlying values. Among these 15% of human MAR sequences, most are very well documented MARs, used to insulate transgene from position effects, such as the interferon locus MAR, the beta-globin locus MAR (Ramezani A, Hawley TS, Hawley RG, "Performance- and safety-enhanced lentiviral vectors containing the human interferon-beta scaffold attachment region and the chicken beta-globin insulator", *Blood*, 101:4717-4724, 2003), or the apolipoprotein MAR (Namciu, S, Blochinger KB, Fournier REK, "Human matrix attachment regions in-sulate transgene expression from chromosomal position effects in *Drosophila melanogaster*", *Mol. Cell. Biol.*, 18:2382-2391, 1998). Always with the same data, human MAR sequences were also used to determine the association between the four theoretical structural properties computed and the AT-content. Fig. 2 represents the scatterplot and the corresponding correlation coefficient  $r$  for every pair of criteria.

### Example 2: Distribution plots of MAR sequences by organism

MAR sequences from SMART DB of other organisms were also retrieved and analyzed similarly as explained previously. The MAR sequences density distributions for the mouse, the chicken, the sorghum bicolor and the human are plotted jointly in Fig. 3.

### Example 3: MAR prediction of the whole chromosome 22

All RefSeq contigs from the chromosome 22 were analyzed by SMAR Scan® using the default settings this time. The result is that SMAR Scan® predicted a total of 803 MARs, their average length being 446 bp, which means an average of one MAR predicted per 42 777 bp. The total length of the predicted MARs corresponds to 1% of the chromosome 22 length. The AT-content of the predicted regions ranged from

65,1% to 93.3%; the average AT-content of all these regions being 73.5%. Thus, predicted MARs were AT-rich, whereas chromosome 22 is not AT-rich (52.1% AT).

SMARTest was also used to analyze the whole chromosome 22 and obtained 1387 MAR candidates, their average length being 494 bp representing an average of one MAR predicted per 24 765 bp. The total length of the predicted MARs corresponds to 2% of the chromosome 22. Between all MARs predicted by the two softwares, 154 predicted MARs are found by both programs, which represents respectively 19% and 11% of SMAR Scan® and SMARTest predicted MARs. Given predicted MARs mean length for SMAR Scan® and SMARTest, the probability to have by chance an overlapping between SMAR Scan® and SMARTest predictions is 0.0027% per prediction.

To evaluate the specificity of SMAR Scan® predictions, SMAR Scan® analyses were performed on randomly shuffled sequences of the chromosome 22 (Fig. 4). Shuffled sequences were generated using 4 different methods: by a segmentation of the chromosome 22 into nonoverlapping windows of 10 bp and by separately shuffling the nucleotides in each window; by "scrambling" which means a permutation of all nucleotides of the chromosome; by "rubbling" which means a segmentation of the chromosome in fragments of 10 bp and a random assembling of these fragments and finally by order 1 Markov chains, the different states being the all the different DNA dinucleotides and the transition probabilities between these states being based on the chromosome 22 scan. For each shuffling method, five shuffled chromosome 22 were generated and analyzed by SMAR Scan® using the default settings. Concerning the number hits, an average of 3 519 170 hits (sd: 18 353) was found for the permuted chromosome 22 within nonoverlapping windows of 10 bp, 171 936,4 hits (sd: 2 859,04 ) for the scrambled sequences and 24 708,2 hits (sd: 1 191,59) for the rubbled chromosome 22 and 2 282 hits in average (sd: 334,7) for the chromosomes generated according to order 1 Markov chains models of the chromosome 22, which respectively represents 185% (sd: 0.5% of the mean), 9% (sd: 1.5%), 1% (sd: 5%) and 0.1% (sd: 15%) of the number of hits found with the native chromosome 22. For the number of MARs predicted, which thus means contiguous hits of length greater than 300, 1 997 MARs were predicted with the shuffled chromosome 22 within windows of 10 bp (sd: 31.2), only 2.4 MARs candidates were found in scrambled sequences (sd: 0.96) and none for the rubbled and for the sequences generated according to Markov chains model, which respectively represents 249% and less than 0.3% of the number of predicted MARs found with the native chromosome 22. These data provide indications that SMAR Scan® detects specific DNA elements which organization is lost when the DNA sequences are shuffled .

#### **Example 4: Analysis of known matrix attachment regions in the Interferon locus with SMAR Scan®**

The relevance of MAR prediction by SMAR Scan® was investigated by analyzing the recently published MAR regions of the human interferon gene cluster on the short arm of chromosome 9 (9p22). Goetze et al. (already cited) reported an exhaustive analysis of the WP18A10A7 locus to analyze the suspected correlation between BURs (termed in this case stress-induced duplex destabilization or SIDD) and *in vitro* binding to the nuclear matrix (Fig. 9, lower part). Three of the SIDD peaks were in agreement with the *in vitro* binding assay, while others did not match matrix attachment sites. Inspection of the interferon locus with SMAR Scan® (Fig. 9, top part) indicated that three majors peaks accompanied by clusters of SATB1, NMP4 and MEF2 regulators binding sites

correlated well with the active MARs. Therefore, we conclude that the occurrence of predicted CUEs and binding sites for these transcription factors is not restricted to the cLysMAR but may be a general property of all MARs. These results also imply that the SMAR Scan® program efficiently detects MAR elements from genomic sequences.

5

**Example 5: Accuracy of SMAR Scan® prediction and comparison with other predictive tools**

10 The accuracy of SMAR Scan® was evaluated using six genomic sequences for which experimentally determined MARs have been mapped. In order to perform a comparison with other predictive tools, the sequences analyzed are the same with the sequences previously used to compare MAR-Finder and SMARTest. These genomic sequences are three plant and three human sequences (Table 1) totalizing 310 151 bp and 37 experimentally defined MARs. The results for SMARTest and MAR-Finder in Table 1  
15 come from a previous comparison (Frisch M, Frech K, Klingenhoff A, Cartharius K, Liebich I and Werner T, In silico pre-diction of scaffold/matrix attachment regions in large genomic sequences, *Genome Research*, 12:349-354, 2001.).  
MAR-Finder has been used with the default parameters excepted for the threshold that has been set to 0.4 and for the analysis of the protamine locus, the AT-richness  
20 rule has been excluded (to detect the non AT-rich MARs as was done for the protamine locus).

Sequence, description and reference	Length (kb)	Experimentally defined MARs positions (kb)	SMARTest prediction positions (kb)	MAR-Finder prediction positions (kb)	SMART Scan prediction positions (kb)
<i>Oryza Sativa</i> putative ADP-glucose pyrophosphorylase subunit SH2 and putative NADPH dependant reductase A1 genes (U70541), [4]	30.034	0.0-1.2 5.4-7.4  17.3-18.5 20.0-23.1	- <b>6.5-7.0</b> 15.2-15.7 16.2-16.6 <b>17.6-18.3</b> <b>19.6-20.1</b> <b>20.7-21.3</b> 23.6-23.9 25.0-25.4 27.5-27.9	- - 15.7-15.9 - <b>17.5-18.4</b> <b>19.8-20.4</b> <b>21.3-21.5</b> 23.9-24.2 24.7-25.1 -	- - 15.6-16 - <b>17.6-18.2</b> <b>21.6-22</b> - 23.4-23.8 - -
<i>Sorghum bicolor</i> ADP-glucose pyrophosphorylase subunit SH2, NADPH-dependant reductase A1-b genes (AF010283), [4]	42.446	0.0-1.5 7.1-9.7  22.4-24.7  32.5-33.7 41.6-42.3	- - 21.3-21.9 <b>22.9-24.0</b> - 27.3-27.6 - -	- - - <b>23.2-24.2</b> - 26.9-27.5 - -	- <b>7.4-7.7</b> 21.5-21.8 <b>22.9-23.2</b> <b>23.6-24.0</b> 27.3-27.6 <b>33.4-33.9</b> -
<i>Sorghum bicolor</i> BAC clone 110K5 (AF124045), [37]	78.195	~0.9 ~5.8 ~6.3 ~9.3 ~15.0 ~18.5 ~21.9 ~23.3 ~25.6 ~29.1 ~34.6  ~44.1 ~48.5  ~57.9 ~62.9 ~67.1 ~69.3 ~73.7	- - - - 15.1-15.8 - <b>21.7-22.0</b> - - - - -  44.1-44.5 47.9-49.5 - - 63.1-63.7 - - - 74.3-74.7	- - - - - - - - - - - -  - 47.9-49.4 - - - - -	- - - - - - <b>21.4-21.9</b> - - <b>29.2-29.5</b> - 39.0-40.0  48.1-48.6 48.8-49.3 - - - 74.3-74.6
Human alpha-1-antitrypsin and corticosteroid binding globulin intergenic region (AF156545), [35]	30.461	2.6-6.3  22.0-30.4	5.5-6.0 - 25.7-26.2 27.5-27.8 - -	3.0-3.2 5.1-6.0 24.9-25.3 25.5-25.8 26.2-26.4 27.5-28.2	5.4-5.8 - <b>25.8-26.4</b> - - -
Human protamine locus (U15422), [24]	53.060	8.8-9.7 32.6-33.6 37.2-39.4 51.8-53.0	- - - -	8.0-8.9* 33.9-34.8* 33.9-34.8* -	- - - -
Human beta-globin locus (U01317), [21]	75.955	1.6-3.0 15.6-19.0  44.7-52.7 60.0-70.0	- <b>18.0-18.4</b> - 34.4-34.9 - 56.6-57.1 59.8-60.3 65.6-66.0	- 15.5-16.0 18.0-18.4 - 50.6-50.8 56.5-57.2 58.1-58.5 63.0-63.6	<b>2.3-2.6</b> <b>15.3-15.6</b> - - - - <b>62.8-63.1</b> -

			67.6-67.9 68.8-69.1	68.7-69.3 -	66.3-66.7 -
Sum(kb)	310.151	at least 56.1	14.5	13.8	9.5
Total numbers :		37	28	25	22
Average kb /predicted MAR			11.076	12.406	14.097
True positives [number of experimentally defined MAR found]			19[14]	20[12]	17[14]
False positives			9	5	5
False negatives			23	25	23
Specificity			19/28= 68%	20/25= 80%	17/22= 77%
Sensitivity			14/37= 38%	12/37= 32%	14/37= 38%

Table 1: Evaluation of SMAR Scan® accuracy

- 5 Six different genomic sequences, three plant and three human sequences, for which experimentally defined MARs are known, were analyzed with MAR-Finder, SMARTest and SMAR Scan®. True positive matches are printed in bold, minus (-) indicates false negative matches. Some of the longer experimentally defined MARs contained more than one in silico prediction, each of them was counted as true positive match.
- 10 Therefore, the number of true in silico predictions is higher than the number of experimentally defined MARs found. Specificity is defined as the ratio of true positive predictions, whereas sensitivity is defined as the ratio of experimentally defined MARs found. \* AT-rich rule excluded using MAR-Finder.
- 15 SMARTest predicted 28 regions as MARs, 19 (true positives) of these correlate with experimentally defined MARs (specificity: 68%) whereas 9 (32%) are located in non-MARs (false positives). As some of the longest experimentally determined MARs contains more than one in silico prediction, the 19 true positives correspond actually to 14 different experimentally defined MARs (sensitivity: 38%). MARFinder
- 20 predicted 25 regions as MARs, 20 (specificity: 80%) of these correlate with experimentally defined MARs corresponding to 12 different experimentally defined MARs (sensitivity: 32%). SMAR Scan® predicted 22 regions, 17 being true positives (specificity: 77%) matching 14 different experimentally defined MARs (sensitivity: 38%).
- 25 As another example, the same analysis has been applied to human chromosomes 1 and 2 and lead to the determination of 23 MARs sequences (SEQ ID N° 1 to 23). These sequences are listed in Annex 1 in ST25 format.

#### **Example 6: Analyses of the whole genome using the combined method (SMAR Scan®-pfsearch)**

- 30 In order to test the potential correlation between the structural features computed by SMAR Scan® and the S/MAR functional activity, the whole human genome has been analyzed with the combined method with very stringent parameters, in order to get
- 35 sequences with the highest values for the theoretical structural features computed, which are called "super" S/MARs below. This was done with the hope to obtain predicted MAR elements with a very potential to increase transgene expression and recombinant protein production. The putative S/MARs hence harvested were first
- 40 analyzed from the bioinformatics perspective in an attempt to characterize and classify them.

## 6.1 S/MARs predicted from the analysis of the whole human genome

As whole human genome sequence, all human RefSeq (National Center for Biotechnology Information, The NCBI handbook [Internet]. Bethesda (MD): National Library of Medicine (US), Oct. Chapter 17, The Reference Sequence (RefSeq) Project, 2002 (Available from <http://www.ncbi.nih.gov/entrez/query.fcgi?db=Books>) contigs (release 5) were used and analyzed with the combined method, using SMAR Scan® as filter in the first level processing, employing default settings except for the highest bend cutoff value, whereas a stringent threshold of 4.0 degrees (instead of 3.202 degrees) has been used for the DNA bending criterion.

In the second level processing, predicted transcription factors binding have been sought in the sequences selected from the previous step without doing any filtering on these sequences.

The analysis by the combined method of the whole human genome came up with a total of 1757 putative "super" S/MARs representing a total of 1 065 305 bp (0.35% of the whole human genome). Table 2 shows for each chromosome: its size, its number of genes, its number of S/MARs predicted, its S/MARs density per gene and its kb per S/MAR. This table shows that there are very various gene densities per S/MAR predicted for the different chromosomes (standard deviation represents more than 50% of the mean of the density of genes per S/MAR predicted and the fold difference between the higher and the lower density of genes per S/MAR is 6,5). Table 2 also shows that the kb per S/MAR varies less than the density of genes per S/MAR (standard deviation represents 25% of the mean of kb per S/MAR and the fold difference between the higher and the lower kb per S/MAR is 3.2).

Chromosome	Number of genes per chromosome	Size of the chromosome (millions bp)	Number of S/MARs predicted	Density of genes per S/MAR	Kb per S/MAR
1	2544	230	85	29.9	2705
2	1772	241	143	12.3	1685
3	1406	198	101	13.9	1960
4	1036	190	118	8.7	1610
5	1233	180	116	10.6	1551
6	1247	170	94	13.2	1808
7	1383	160	179	7.7	1754
8	942	145	77	12.2	1883
9	1100	119	48	22.9	2479
10	1003	133	71	14.1	1873
11	1692	132	67	25.2	1970
12	1278	131	78	16.3	1679
13	506	97	70	7.2	1385
14	1168	88	36	32.4	2444
15	895	83	35	25.5	2371
16	1107	81	41	27	1975
17	1421	80	37	38.4	2162
18	396	75	51	7.7	1470
19	1621	56	36	45.02	1555
20	724	60	28	25.8	2142
21	355	34	18	19.7	1888
22	707	34	28	25.2	1214
X	1168	154	170	6.8	905
Y	251	25	30	8.3	833
<b>Sum</b>	26 955	3 050	1 757	457	433 12
<b>Mean</b>	1 123	127	73	19	1 804
<b>Sd</b>	510	72.8	45	10	462

Table 2: Number of S/MARs predicted per chromosome. The number of genes per chromosome

corresponds to the NCBI human genome statistics (Build 34 Version 3) (National Center for Biotechnology Information, The NCBI handbook [Internet]. Bethesda (MD): National Library of Medicine (US), Oct. Chapter 17, The Reference Sequence (RefSeq) Project, 2002 (Available from <http://www.ncbi.nih.gov/entrez/query.fcgi?db=Books>) based on GenBank annotations.

Chromosome sizes are the sum of the corresponding human RefSeq (National Center for Biotechnology Information, The NCBI handbook [Internet]. Bethesda (MD): National Library of Medicine (US), Oct. Chapter 17, The Reference Sequence (RefSeq) Project, 2002 (Available from <http://www.ncbi.nih.gov/entrez/query.fcgi?db=Books>) (release 5) contig lengths

## 6.2 Bioinformatics analysis of "super" MARS for transcription factor binding sites

The 1757 predicted "super" S/MARs sequences obtained previously by SMAR Scan® were then analyzed for potential transcription factors binding sites. This has been achieved using RMatch™ Professional (Kel AE, Gossling E, Reuter I, Cheremushkin E, KelMargoulis OV, Wingender E, MATCH: A tool for searching transcription factor binding sites in DNA sequences, *Nucleic Acids Res.* 31(13):35769, 2003), a weight matrixbased tool based on TRANSFAC (Wingender E, Chen X, Fricke E, Geffers R, Hehl R, Liebich I, Krull M, Matys V, Michael H, Ohnhauser R, Pruss M, Schacherer F, Thiele S, Urbach S, The TRANSFAC system on gene expression regulation, *Nucleic Acids Research*, 29(1):2813, 2001). Match™ 2.0 Professional has been used with most of the default settings Match™ analysis was based on TRANSFAC Professional, release 8.2 (20040630). The sums of all transcription factors binding prediction on the 1757 sequences analyzed according to Match™ are in Table 3. Based on this table, only the transcription factors totalizing at least 20 hits over the 1757 sequences analyzed were considered for further analyses.

Hereafter are some of the human transcription factors that are the most often predicted to bind on the 1757 putative S/MAR sequences and their Match description: Cdc5 (cell division control protein 5) a transcriptional regulator/repressor, Nkx3A a homeodomain protein regulated by androgen, POU1F1 (pituitaryspecific positive transcription factor 1) which is specific to the pituitary and stimulates cells proliferation. Thus, in addition to SATB1, NMP4 and MEF2, other transcription factors can participate in the activity of MARs.

AP1	1	AR	2	Bach2	1	Brn2	1
C/EBP	20	C/EBPgamma	5	CDP CR3	1	COMP1	2
CREBP1	34	Cdc5	858	Cdx2	35	Evi1	472
FOX	78	FOXD3	79	FOXJ2	244	FOXP3	29
Freac7	272	GATA1	2	GATA3	142	GATA4	125
HFH1	12	HFH3	1	HLF	275	HNF1	337
HNF3alpha	23	HNF3beta	71	HP1	2	Lhx3	22
MEF2	114	MRF2	57	Myc	18	NKX3A	849
Nkx25	2	Oct1	191	PBX	5	POU1F1	483
POU3F2	11	POU6F1	29	Pax3	3	Pax6	20
Pit1	505	SRF	8	TEF	2852	TFIIA	14
TTF1	1	V\$MTATA_B	4	VBP	53	Vmw65	1
XFD1	65	XFD2	418	XFD3	2		

Table 3 is a summary of all transcription factors binding prediction (totalizing 20 hits or more) on the 1757 sequences analyzed.

### 5 6.3 Bioinformatics analysis of predicted "super" MARs for dinucleotide frequencies

10 Various computer analysis were performed in order to easily identify "super" S/MAR sequences using an explicit criterion that could be identified without computing. Among those, a di-nucleotide analysis was performed on the 1757 superMARs, computing each of the 16 possible dinucleotide percentage for each sequence considering both strands in the 5' > 3' direction.

15 A summary (min., max., median, mean, 25th percentile and 75th percentile) as well as the histograms of each dinucleotide percentage over the 1757 S/MAR sequences are respectively presented in Table 4. A similar analysis was performed on randomly selected sequences from the human genome, representing randomly selected non-S/MAR sequences (which might however contain some MARs). Table 5 represents respectively a summary of the dinucleotide content analysis for these sequences.

Table 4: Dinucleotide percentages over the 1757 S/MAR sequences

20

	AA %	AC %	AG %	AT %
Minimum	0.000	0.0000	0.0000	18.50
25th percentile	4.234	0.9372	0.1408	32.11
Median	7.843	2.2408	0.4777	34.68
Mean	7.184	3.2117	1.0865	34.32
75th percentile	10.110	4.7718	1.5096	36.94
Maximum	17.290	12.9479	8.1230	50.00
	CA %	CC %	CG %	CT %
Minimum	0.0000	0.00000	0.0000	0.0000
25th percentile	0.9695	0.00000	0.0000	0.1408
Median	1.9776	0.00000	0.0000	0.4777
Mean	2.6977	0.14123	0.2709	1.0865
75th percentile	3.7543	0.09422	0.1256	1.5096
Maximum	10.4061	4.24837	7.4410	8.1230
	GA %	GC %	GG %	GT %
Minimum	0.00000	0.0000	0.00000	0.0000
25th percentile	0.08696	0.0000	0.00000	0.9372
Median	0.32616	0.0000	0.00000	2.2408
Mean	0.63347	0.2104	0.14123	3.2117
75th percentile	0.83333	0.1914	0.09422	4.7718
Maximum	5.77889	9.8795	4.24837	12.9479
	TA %	TC %	TG %	TT %
Minimum	28.63	0.00000	0.0000	0.000
25th percentile	33.48	0.08696	0.9695	4.234
Median	35.22	0.32616	1.9776	7.843
Mean	35.29	0.63347	2.6977	7.184
75th percentile	37.14	0.83333	3.7543	10.110
Maximum	50.00	5.77889	10.4061	17.290

25 Considering the results of the predicted S/MAR elements and of the nonS/MAR sequences in the summary tables, noticeable differences can be noticed in the AT et TA dinucleotide contents between these two groups of sequences. AT and TA represent respectively at least 18,5 % and 28.6 % of the dinucleotide content of the predicted S/MAR sequences, whereas the minimum percentages for the same dinucleotides in

nonS/MAR sequences are respectively 0.3 % and 0%. Similarly, the maximum CC and GG content in S/MAR sequences is 4.2 %, whereas in nonS/MAR sequences the percentages for these two dinucleotides can amount up to 20.8 %.

The correlation between AT and TA dinucleotide percentages and the DNA highest bend as computed by SMAR Scan® is depicted in Fig. 17 for the predicted S/MAR sequences and in Fig. 18 for the nonS/MAR sequences. The different scatterplots of these figures show that the TA percentage correlates well with the predicted DNA bend as predicted by SMAR Scan®.

Table 5: Dinucleotide percentages over the 1757 nonS/MAR sequences summary

	AA %	AC %	AG %	AT %
Minimum	0.000	1.735	1.512	0.3257
25th percentile	7.096	4.586	6.466	5.1033
Median	9.106	5.016	7.279	6.8695
Mean	8.976	5.054	7.184	7.0108
75th percentile	10.939	5.494	7.969	8.7913
Maximum	17.922	13.816	12.232	23.1788
	CA %	CC %	CG %	CT %
Minimum	3.571	0.8278	0.0000	1.512
25th percentile	6.765	4.1077	0.4727	6.466
Median	7.410	5.5556	0.8439	7.279
Mean	7.411	5.9088	1.2707	7.184
75th percentile	8.010	7.2460	1.5760	7.969
Maximum	15.714	20.8415	12.6074	12.232
	GA %	GC %	GG %	GT %
Minimum	1.319	0.4967	0.8278	1.735
25th percentile	5.495	3.2615	4.1077	4.586
Median	6.032	4.4092	5.5556	5.016
Mean	6.065	4.7468	5.9088	5.054
75th percentile	6.602	5.8824	7.2460	5.494
Maximum	10.423	16.0000	20.8415	13.816
	TA %	TC %	TG %	TT %
Minimum	0.000	1.319	3.571	0.000
25th percentile	3.876	5.495	6.765	7.096
Median	5.625	6.032	7.410	9.106
Mean	5.774	6.065	7.411	8.976
75th percentile	7.464	6.602	8.010	10.939
Maximum	24.338	10.423	15.714	17.922

Four of the novel super MARs were randomly picked and analyzed for AT and TA dinucleotide content, and compared with the previously known chicken lysMAR, considering windows of 100 base pairs (Table 6).

Surprisingly, Applicants have shown that all of the super MARs have AT dinucleotide frequencies greater than 12%, and TA dinucleotides greater than 10% of the total dinucleotides analysed in a window of 100base pairs of DNA. The most efficient MARs display values around 34% of the two dinucleotide pairs.

Table 6. Summary of %AT and TA dinucleotide frequencies of experimentally verified MARs

CLysMAR (average of CUEs)	AT% : 12.03	TA% : 10.29	SEQ ID No
P1_68	AT% : 33.78	TA% : 33.93	SEQ ID No
P1_6	AT% : 34.67	TA% : 34.38	SEQ ID No

P1_42	AT% : 35.65	TA% : 35.52	SEQ ID No
Mean value for all human "super"MARs	AT% : 34.32	TA% : 35.29	
Mean value for all human non-MARs	AT% : 7.01	TA% : 5.77	

#### 6.4 Analysis of orthologous intergenic regions of human and mouse genomes

- 5 In order to get an insight on S/MAR evolution, orthologous intergenic regions of human and mouse genomes have been analysed with SMAR Scan®. The data set used is composed of 87 pairs of complete orthologous intergenic regions from the human and mouse genomes (Shabalina SA, Ogurtsov AY, Kondrashov VA, Kondrashov AS, Selective constraint in intergenic regions of human and mouse genomes, *Trends*
- 10 *Genet*, 17(7):3736, 2001) (average length ~12 000 bp) located on 12 human and on 12 mouse chromosomes, the synteny of these sequences was confirmed by pairwise sequence alignment and consideration of the annotations of the flanking genes (experimental or predicted).
- 15 Analysis of the 87 human and mouse orthologous intergenic sequences have been analysed with SMAR Scan® using its default settings. Analysis of the human sequences yielded a total of 12 S/MARs predicted (representing a total length of 4 750 bp), located on 5 different intergenic sequences.
- 20 Among the three human intergenic sequences predicted to contain a "super" S/MAR using SMAR Scan® stringent settings, one of the corresponding mouse orthologous intergenic sequence is also predicted to contain a S/MAR (human EMBL ID: Z96050, position 28 010 to 76 951 orthologous to mouse EMBL ID: AC015932, positions 59 884 to 89 963). When a local alignment of these two orthologous intergenic sequences is
- 25 performed, the best local alignment of these two big regions correspond to the regions predicted by SMAR Scan® to be S/MAR element. A manual search for the mouse orthologs of the two other human intergenic sequences predicted to contain a "super" S/MAR was performed using the Ensembl Genome Browser (<http://ensembl.org>). The mouse orthologous intergenic sequences of these two human sequences were
- 30 retrieved using Ensembl orthologue predictions (based on gene names), searching the orthologous mouse genes for the pairs of human genes flanking these intergenic regions.
- 35 Because SMAR Scan® has been tuned for human sequences and consequently yields little "super"MARs with mouse genomic sequences, its default cutoff values were slightly relaxed for the minimum size of contiguous hits to be considered as S/MAR (using 200 bp instead of 300 bp). Analysis by SMAR Scan® of these mouse sequences predicted several S/MARs having high values for the different computed structural features. This finding suggests that the human MAR elements are conserved across
- 40 species.

#### **Example 7 : Dissection of the chicken lysozyme gene 5'- MAR**

- 45 The 3000 base pair 5'-MAR was dissected into smaller fragments that were monitored for effect on transgene expression in Chinese hamster ovary (CHO) cells. To do so, seven fragments of ~400 bp were generated by polymerase chain reaction (PCR). These PCR-amplified fragments were contiguous and cover the entire MAR sequence when placed end-to-end. Four copies of each of these fragments were ligated in a head-to-tail orientation, to obtain a length corresponding to approximately half of that of

the natural MAR. The tetramers were inserted upstream of the SV40 promoter in pGEGFPControl, a modified version of the pGL3Control vector (Promega). The plasmid pGEGFPControl was created by exchanging the luciferase gene of pGL3Control for the EGFP gene from pEGFP-N1 (Clontech). The 5'-MAR-fragment-containing plasmids thus created were co-transfected with the resistance plasmid pSVneo in CHO-DG44 cells using LipofectAmine 2000 (Invitrogen) as transfection reagent, as performed previously (Zahn-Zabal, M., et al., "Development of stable cell lines for production or regulated expression using matrix attachment regions" *J Biotechnol*, 2001. 87(1): p. 29-42.). After selection of the antibiotic (G-418) resistant cells, polyclonal cell populations were analyzed by FACS for EGFP fluorescence.

Transgene expression was expressed at the percentile of high expressor cells, defined as the cells which fluorescence levels are at least 4 orders of magnitude higher than the average fluorescence of cells transfected with the pGEGFPControl vector without MAR. Fig. 5 shows that multimerized fragments B, K and F enhance transgene expression, despite their shorter size as compared to the original MAR sequence. In contrast, other fragments are poorly active or fully inactive.

### **Example 8 : Specificity of B, K and F regions in the MAR context**

The 5'-MAR was serially deleted from the 5'-end (Fig.6, upper part) or the 3'-end (Fig.6, lower part), respectively. The effect of the truncated elements was monitored in an assay similar to that described in the previous section. Figure 6 shows that the loss of ability to stimulate transgene expression in CHO cells was not evenly distributed.

In this deletion study, the loss of MAR activity coincided with discrete regions of transition which overlap with the 5'-MAR B-, K- and F-fragment, respectively. In 5' deletions, activity was mostly lost when fragment K and F were removed. 3' deletions that removed the F and b elements had the most pronounced effects. In contrast, flanking regions A, D, E and G that have little or no ability to stimulate transgene expression on their own (Fig. 5), correspondingly did not contribute to the MAR activity in the 5'- and 3'-end deletion studies (Fig. 6).

### **Example 9: Structure of the F element**

The 465 bp F fragment was further dissected into smaller sub-fragments of 234, 243, 213 bp and 122, 125 and 121 bp, respectively. Fragments of the former group were octamerized (8 copies) in a head-to-tail orientation, while those of the latter group were similarly hexa-decamerized (16 copies), to maintain a constant length of MAR sequence. These elements were cloned in pGEGFPControl vector and their effects were assayed in CHO cells as described previously. Interestingly, fragment FIII retained most of the activity of the full-length F fragment whereas fragment FII, which contains the right-hand side part of fragment FIII, lost all the ability to stimulate transgene expression (Fig. 7). This points to an active region comprised between nt 132 and nt 221 in the FIB fragment. Consistently, multiple copies of fragments FI and FIB, which encompass this region, displayed similar activity. FIIA on its own has no activity. However, when added to FIB, resulting in FIII, it enhances the activity of the former. Therefore FIIA appears to contain an auxiliary sequence that has little activity on its own, but that strengthens the activity of the minimal domain located in FIB.

Analysis of the distribution of individual motifs within the lysozyme gene 5'-MAR is shown in Fig. 8 A, along with some additional motifs that we added to the analysis. Most of these motifs were found to be dispersed throughout the MAR element, and not

specifically associated with the active portions. For instance, the binding sites of transcription factors and other motifs that have been associated with MARs were not preferentially localized in the active regions. It has also been proposed that active MAR sequences may consist of combination of distinct motifs. Several computer programs (MAR Finder, SMARTest, SIDD duplex stability) have been reported to identify MARs as regions of DNA that associate with the DNA matrix. They are usually based on algorithms that utilizes a predefined series of sequence-specific patterns that have previously been suggested as containing MAR activity, as exemplified by MAR Finder, now known as MAR Wiz. The output of these programs did not correlate well with the transcriptionally active portions of the *cLysMAR*. For instance, peaks of activity obtained with MAR Finder did not clearly match active MAR sub-portion, as for instance the B fragment is quite active in vivo but scores negative with MAR Finder (Fig. 8B, compare the top and middle panels). Bent DNA structures, as predicted by this program, did not correlate well either with activity (Fig. 8B, compare the top and bottom panels). Similar results were obtained with the other available programs (data not shown).

The motifs identified by available MAR prediction computer methods are therefore unlikely to be the main determinants of the ability of the *cLysMAR* to increase gene expression. Therefore, a number of other computer tools were tested. Surprisingly, predicted nucleosome binding sequences and nucleosome disfavoured sequences were found to be arranged in repetitively interspersed clusters over the MAR, with the nucleosome favouring sites overlapping the active B, K and F regions. Nucleosome positioning sequences were proposed to consist of DNA stretches that can easily wrap around the nucleosomal histones, and they had not been previously associated with MAR sequences.

Nucleosome-favouring sequences may be modelled by a collection of DNA features that include moderately repeated sequences and other physico-chemical parameters that may allow the correct phasing and orientation of the DNA over the curved histone surface. Identification of many of these DNA properties may be computerized, and up to 38 different such properties have been used to predict potential nucleosome positions. Therefore, we set up to determine if specific components of nucleosome prediction programs might correlate with MAR activity, with the objective to construct a tool allowing the identification of novel and possibly more potent MARs from genomic sequences.

To determine whether any aspects of DNA primary sequence might distinguish the active B, K and F regions from the surrounding MAR sequence, we analyzed the 5'-MAR with MAR Scan®. Of the 38 nucleosomal array prediction tools, three were found to correlate with the location of the active MAR sub-domains (Fig. 9A). Location of the MAR B, K and F regions coincides with maxima for DNA bending, major groove depth and minor groove width. A weaker correlation was also noted with minima of the DNA melting temperature, as determined by the GC content. Refined mapping over the MAR F fragment indicated that the melting temperature valley and DNA bending summit indeed correspond the FIB sub-fragment that contains the MAR minimal domain (Fig. 9B). Thus active MAR portions may correspond to regions predicted as curved DNA regions by this program, and we will refer to these regions as CUE-B, CUE-K and CUE-F in the text below. Nevertheless, whether these regions correspond to actual bent DNA and base-pair unwinding regions is unknown, as they do not correspond to bent DNA as predicted by MAR Wiz (Fig.9B).

#### **Example 10 : Imprints of other regulatory elements in the F fragment**

Nucleosome positioning features may be considered as one of the many specific chromatin codes contained in genomic DNA. Although this particular code may contribute to the activity of the F region, it is unlikely to determine MAR activity alone, as the 3' part of the F region enhanced activity of the minimal MAR domain contained in the FIB portion. Using the MatInspector program (Genomatix), we searched for transcription factor binding sites with scores higher than 0.92 and found DNA binding sequences for the NMP4 and MEF2 proteins in the 3' part of the F fragment (Fig. 8B). To determine whether any of these transcription factor-binding sites might localize close to the B and K active regions, the entire 5'-MAR sequence was analyzed for binding by NMP4 and MEF2 and proteins reported to bind to single-stranded or double-stranded form of BURs. Among those, SATB1 (special AT-rich binding protein 1) belongs to a class of DNA-binding transcription factor that can either activate or repress the expression of nearby genes. This study indicated that specific proteins such as SATB1, NMP4 (nuclear matrix protein 4) and MEF2 (myogenic enhancer factor 2), have a specific distribution and form a framework around the minimal MAR domains of *cLysMAR* (Fig. 10). The occurrence of several of these NMP4 and SATB1 binding sites has been confirmed experimentally by the EMSA analysis of purified recombinant proteins (data not shown).

#### **Example 11 : Construction of artificial MARs by combining defined genetic elements**

To further assess the relative roles of the various MAR components, the *cLysMAR* was deleted of all three CUE regions (Fig. 11, middle part), which resulted in the loss of part of its activity when compared to the complete MAR sequence similarly assembled from all of its components as a control (Fig. 11, top part). Consistently, one copy of each CUE alone, or one copy of each of the three CUEs assembled head-to-tail, had little activity in the absence of the flanking sequences. These results strengthen the conclusion that optimal transcriptional activity requires the combination of CUEs with of flanking sequences. Interestingly, the complete MAR sequence generated from each of its components, but containing also BglII-BamHI linker sequences (AGATCC) used to assemble each DNA fragment, displayed high transcriptional activity (6 fold activation) as compared to the 4.8 fold noted for the original MAR element in this series of assays (see Fig. 5).

We next investigated whether the potentially curved DNA regions may also be active in an environment different from that found in their natural MAR context. Therefore, we set up to swap the CUE-F, CUE-B and CUE-K elements, keeping the flanking sequences unchanged. The sequences flanking the CUE-F element were amplified by PCR and assembled to bracket the various CUEs, keeping their original orientation and distance, or without a CUE. These engineered ~1.8 kb MARs were then assayed for their ability to enhance transgene expression as above. All three CUE were active in this context, and therefore their action is not restricted to one given set of flanking sequences. Interestingly, the CUE-K element was even more active than CUE-F when inserted between the CUE-F flanking sequences, and the former composite construct exhibited an activity as high as that observed for the complete natural MAR (4.8 fold activation). What distinguishes the CUE-K element from CUE-F and CUE-B is the presence of overlapping binding sites for the MEF-2 and SatB1 proteins, in addition to its CUE feature. Therefore, fusing CUE-B with CUE-F-flanking domain results in a higher density of all three binding sites, which is likely explanation to the increased activity. These results indicate that assemblies of CUEs with sequences containing binding sites

for proteins such as NMP4, MEF-2, SatB1, and/or polyPpolyQ proteins constitute potent artificial MAR sequences.

### **Example 12 : Expression vectors**

Three expression vectors according to the present invention are represented on Figure 12.

**Plasmid pPAG01** is a 5640 bp pUC19 derivative. It contains a 2960 bp chicken DNA fragment cloned in *Bam*H1 and *Xba*I restriction sites. The insert comes from the border of the 5'-end of the chicken lysozyme locus and has a high A/T-content.

**Plasmid pGEGFP** (also named pSV40EGFP) control is a derivative of the pGL3-control vector (Promega) in which the luciferase gene sequence has been replaced by the EGFP gene sequence from the pEGFP-N1 vector (Clontech). The size of pGEGFP plasmid is 4334bp.

**Plasmid pUbCEGFP** control is a derivative of the pGL3 with an Ubiquitin promoter.

**Plasmid pPAG01GFP** (also named pMAR-SV40EGFP) is a derivative of pGEGFP with the 5'-Lys MAR element cloned in the MCS located just upstream of the SV40 promoter. The size of the pPAG01EGF plasmid is 7285bp.

### **Example 13 : Effect of the additional transfection of primary transfectant cells on transgene expression**

One day before transfection, cells were plated in a 24-well plate, in growth medium at a density of  $1.35 \times 10^5$  cells/well for CHO-DG44 cells. 16 hours post-inoculum, cells were transfected when they reached 30-40% confluence, using Lipofect-AMINE 2000 (hereinafter LF2000), according to the manufacturer's instructions (Invitrogen). Twenty-seven microliters of serum free medium (Opti-MEM; Invitrogen) containing 1.4  $\mu$ l of LF2000 were mixed with 27  $\mu$ l of Opti-MEM containing 830 ng of linear plasmid DNA. The antibiotic selection plasmid (pSVneo) amounted to one tenth of the reporter plasmid bearing the GFP transgene. The mix was incubated at room temperature for 20 min, to allow the DNA-LF2000 complexes to form. The mixture was diluted with 300  $\mu$ l of Opti-MEM and poured into previously emptied cell-containing wells. Following 3 hours incubation of the cells with the DNA mix at 37°C in a CO<sub>2</sub> incubator, one ml of DMEM-based medium was added to each well. The cells were further incubated for 24 hours in a CO<sub>2</sub> incubator at 37°C. The cells were then transfected a second time according to the method described above, except that the resistance plasmid carried another resistance gene (pSVpuro). Twenty-four hours after the second transfection, cells were passaged and expanded into a T-75 flask containing selection medium supplemented with 500  $\mu$ g/ml G-418 and 5  $\mu$ g/ml puromycin. After a two week selection period, stably transfected cells were cultured in 6-well plates. Alternatively, the cell population was transfected again using the same method, but pTKhygro (Clontech) and pSVdhfr as resistance plasmids. The expression of GFP was analysed with Fluorescence-activated cell sorter (FACS) and with a Fluorocan.

Fig.13 shows that the phenotype of the twice-transfected cells (hereafter called secondary transfectants) not only was strongly coloured, such that special bulb and filter were not required to visualize the green color from the GFP protein, but also contained a majority of producing cells (bottom right-hand side FACS histogram) as compared to the parental population (central histogram). This level of fluorescence corresponds to specific cellular productivities of at least 10 pg per cell per day. Indeed,

cells transfected only one time (primary transfectants) that did not express the marker protein were almost totally absent from the cell population after re-transfection. Bars below  $10^1$  units of GFP fluorescence amounted 30% in the central histogram and less than 5% in the right histogram. This suggested that additional cells had been transfected and successfully expressed GFP.

Strikingly, the amount of fluorescence exhibited by re-transfected cells suggested that the subpopulation of cells having incorporated DNA twice expressed much more GFP than the expected two-fold increase. Indeed, the results shown in Table 2 indicate that the secondary transfectants exhibited, on average, more than the two-fold increase of GFP expected if two sets of sequences, one at each successive transfection, would have been integrated independently and with similar efficiencies. Interestingly, this was not dependent on the promoter sequence driving the reporter gene as both viral and cellular promoter-containing vectors gave a similar GFP enhancement (compare lane 1 and 2). However, the effect was particularly marked for the MAR-containing vector as compared to plasmids without MAR- (lane 3), where the two consecutive transfections resulted in a 5.3 and 4.6 fold increase in expression, in two distinct experiments.

Type of plasmids	Primary transfection	Secondary transfection	EGFP fluorescence Fold increase
pUbCEGFP	4'992	14'334	2.8
pSV40EGFP	4'324	12'237	2.8
pMAR-SV40EGFP	6'996	36'748	<b>5.3</b>

Type of plasmids	Primary transfection	Secondary transfection	EGFP fluorescence Fold increase
pUbCEGFP	6'452	15'794	2.5
pSV40EGFP	4'433	11'735	2.6
pMAR-SV40EGFP	8'116	37'475	<b>4.6</b>

**Table 7.** Effect of re-transfecting primary transfectants at 24 hours interval on GFP expression. Two independent experiments are shown. The resistance plasmid pSVneo was co-transfected with various GFP expression vectors. One day post-transfection, cells were re-transfected with the same plasmids with the difference that the resistance plasmid was changed for pSVpuro. Cells carrying both resistance genes were selected on 500  $\mu$ g/ml G-418 and 5  $\mu$ g/ml puromycin and the expression of the reporter gene marker was quantified by Fluorocan. The fold increases correspond to the ratio of fluorescence obtained from two consecutive transfections as compared to the sum of fluorescence obtained from the corresponding independent transfections. The fold increases that were judged significantly higher are shown in bold, and correspond to fluorescence values that are consistently over 2-fold higher than the addition of those obtained from the independent transfections.

The increase in the level of GFP expression in multiply transfected cells was not expected from current knowledge, and this effect had not been observed previously.

Taken together, the data presented here support the idea that the plasmid sequences that primarily integrated into the host genome would facilitate integration of other plasmids by homologous recombination with the second incoming set of plasmid molecules. Plasmid recombination events occur within a 1-h interval after the plasmid DNA has reached the nucleus and the frequency of homologous recombination

between co-injected plasmid molecules in cultured mammalian cells has been shown to be extremely high, approaching unity (Folger, K.R., K. Thomas, and M.R. Capecchi, Nonreciprocal exchanges of information between DNA duplexes coinjected into mammalian cell nuclei. *Mol Cell Biol*, 1985. 5(1): p. 59-69], explaining the integration of multiple plasmid copies. However, homologous recombination between newly introduced DNA and its chromosomal homolog normally occurs very rarely, at a frequency of 1 in  $10^3$  cells receiving DNA to the most [ Thomas, K.R., K.R. Folger, and M.R. Capecchi, High frequency targeting of genes to specific sites in the mammalian genome. *Cell*, 1986. 44(3): p. 419-28.]. Thus, the results might indicate that the MAR element surprisingly acts to promote such recombination events. MARs would not only modify the organization of genes in vivo, and possibly also allow DNA replication in conjunction with viral DNA sequences, but they may also act as DNA recombination signals.

#### **Example 14 : MARs mediate the unexpectedly high levels of expression in multiply transfected cells**

If MAR-driven recombination events were to occur in the multiple transfections process, we expect that the synergy between the primary and secondary plasmid DNA would be affected by the presence of MAR elements at one or both of the transfection steps. We examined this possibility by multiply transfections of the cells with pMAR alone or in combination with various expression plasmids, using the method described previously. Table 3 shows that transfecting the cells twice with the pMAR-SV40EGFP plasmid gave the highest expression of GFP and the highest degree of enhancement of all conditions (4.3 fold). In contrast, transfecting twice the vector without MAR gave little or no enhancement, 2.8-fold, instead of the expected two-fold increase. We conclude that the presence of MAR elements at each transfection step is necessary to achieve the maximal protein synthesis.

**Table 8**

Primary transfection		Secondary transfection		
Type of plasmid	EGFP-fluorescence	Type of plasmid	EGFP-fluorescence	Fold increase
pMAR	0	pMAR	0	0
		pSV40EGFP	15'437	<b>2.3-2.5</b>
		pMAR-SV40EGFP	30'488	<b>2.6-2.7</b>
pMAR-SV40EGFP	11'278	pMAR-SV40EGFP	47'027	<b>4.3-5.3</b>
		pMAR	12'319	1.0-1.1
pSV40EGFP	6'114	pSV40EGFP	17'200	2.8
		pMAR	11'169	1.8-2.3

Interestingly, when cells were first transfected with pMAR alone, and then re-transfected with pSV40EGFP or pMAR-SV40EGFP, the GFP levels were more than doubled as compared to those resulting from the single transfection of the later plasmids (2.5 and 2.7 fold respectively, instead of the expected 1-fold). This indicates that the prior transfection of the MAR can increase the expression of the plasmid used in the second transfection procedure. Because MARs act only locally on chromatin structure and gene expression, this implies that the two types of DNA may have integrated at a similar chromosomal locus. In contrast, transfecting the GFP expression vectors alone, followed by the MAR element in the second step, yielded little or no improvement of the GFP levels. This indicates that the order of plasmid

transfection is important, and that the first transfection event should contain a MAR element to allow significantly higher levels of transgene expression.

If MAR elements favoured the homologous recombination of the plasmids remaining in episomal forms from the first and second transfection procedures, followed by their co-integration at one chromosomal locus, one would expect that the order of plasmid transfection would not affect GFP levels. However, the above findings indicate that it is more favourable to transfect the MAR element in the first rather than in the second transfection event. This suggests the following molecular mechanism: during the first transfection procedure, the MAR elements may concatemerize and integrate, at least in part, in the cellular chromosome. This integrated MAR DNA may in turn favour the further integration of more plasmids, during the second transfection procedure, at the same or at a nearby chromosomal locus.

#### **Example 15 : MARs as long term DNA transfer facilitators**

If integrated MARs mediated a persistent recombination-permissive chromosomal structure, one would expect high levels of expression even if the second transfection was performed long after the first one, at a time when most of the transiently introduced episomal DNA has been eliminated. To address this possibility, the cells from Table 3, selected for antibiotic resistance for three weeks, were transfected again once or twice and selected for the incorporation of additional DNA resistance markers. The tertiary, or the tertiary and quaternary transfection cycles, were performed with combinations of pMAR or pMAR-SV40EGFP, and analyzed for GFP expression as before.

**Table 9**

Tertiary transfection			Quaternary transfection		
Type of plasmid	EGFP-fluorescence	Fold increase	Type of plasmid	EGFP-fluorescence	Fold increase
pMAR	18368	2.2	pMAR pMAR-SV40EGFP	43'186 140'000	2.4 7.6
pMAR-SV40EGFP	16544	2.0	pMAR-SV40EGFP pMAR	91'000 33'814	5.5 2.0

**Table 9.** MARs act as facilitator of DNA integration.

The pMAR-SV40EGFP/ pMAR-SV40EGFP secondary transfectants were used in a third cycle of transfection at the end of the selection process. The tertiary transfection was accomplished with pMAR or pMAR-SV40EGFP, and pTKhygro as selection plasmid, to give tertiary transfectants. After 24 hours, cells were transfected again with either plasmid and pSVdhfr, resulting in the quaternary transfectants which were selected in growth medium containing 500 µg/ml G-418 and 5µg/ml puromycin, 300 µg/ml hygromycin B and 5µM methotrexate. The secondary transfectants initially exhibited a GFP fluorescence of 8300. The fold increases correspond to the ratio of fluorescence obtained from two consecutive transfections as compared to the sum of

fluorescence obtained from the corresponding independent transfections. The fold increases that were judged significantly higher are shown in bold, and correspond to fluorescence values that are 2-fold higher than the addition of those obtained from the independent transfections.

These results show that loading more copies of pMAR or pMAR-SV40EGFP resulted in similar 2-fold enhancements of total cell fluorescence. Loading even more of the MAR in the quaternary transfection further enhanced this activity by another 2.4-fold. This is consistent with our hypothesis that newly introduced MAR sequences may integrate at the chromosomal transgene locus by homologous recombination and thereby further increase transgene expression.

When the cells were transfected a third and fourth time with the pMAR-SV40EGFP plasmid, GFP activity further increased, once again to levels not expected from the addition of the fluorescence levels obtained from independent transfections. GFP expression reached levels that resulted in cells visibly glowing green in day light (Fig.14). These results further indicate that the efficiency of the quaternary transfection was much higher than that expected from the efficacy of the third DNA transfer, indicating that proper timing between transfections is crucial to obtain the optimal gene expression increase, one day being preferred over a three weeks period.

We believe that MAR elements favour secondary integration events in increasing recombination frequency at their site of chromosomal integration by relaxing closed chromatin structure, as they mediate a local increase of histone acetylation (Yasui, D., et al., SATB1 targets chromatin remodelling to regulate genes over long distances. *Nature*, 2002. 419(6907): p. 641-5.). Alternatively, or concomitantly, MARs potentially relocate nearby genes to subnuclear locations thought to be enriched in trans-acting factors, including proteins that can participate in recombination events such as topoisomerases. This can result in a locus in which the MAR sequences can bracket the pSV40EGFP repeats, efficiently shielding the transgenes from chromatin-mediated silencing effects.

**Example 16 : Use of MARs identified with SMAR Scan® II to increase the expression of a recombinant protein.**

Four MAR elements were randomly selected from the sequences obtained from the analysis of the complete human genome sequence with SMAR Scan® or the combined method. These are termed 1\_6, 1\_42, 1\_68, (where the first number represents the chromosome from which the sequence originates, and the second number is specific to the predicted MAR along this chromosome) and X\_S29, a "super" MAR identified on chromosome X. These predicted MARs were inserted into the pGEGFPControl vector upstream of the SV40 promoter and enhancer driving the expression of the green fluorescent protein and these plasmids were transfected into cultured CHO cells, as described previously (Zahn-Zabal, M., et al., *Development of stable cell lines for production or regulated expression using matrix attachment regions*. J Biotechnol, 2001. 87(1): p. 29-42). Expression of the transgene was then analyzed in the total population of stably transfected cells using a fluorescent cell sorter (FACS) machine. As can be seen from Fig. 19, all of these newly identified MARs increased the expression of the transgene significantly above the expression driven by the chicken lysosome MAR, the "super" MAR X\_S29 being the most potent of all of the newly identified MARs.

**Example 17: Effect on hematocrit of *in vivo* expression of mEpo by electrotransfer of Network system with and without Human MAR (1-68).**

5 The therapeutic gene encodes EPO (erythropoietin), an hormone used for the treatment of anemia. The EPO gene is placed under the control of a doxycycline inducible promoter, in a gene switch system described previously called below the Network system (Imhof, M. O., Chatellard, P., and Mermod, N. (2000). A regulatory network for efficient control of transgene expression. J. Gene. Med. 2, 107-116.). The  
10 EPO and regulatory genes are then injected in the muscle of mice using an *in vivo* electroporation procedure termed the electrotransfer, so that the genes are transferred to the nuclei of the muscle fibers. When the doxycycline antibiotic is added to the drinking water of the mice, this compound is expected to induce the expression of EPO, which will lead to the elevation of the hematocrit level, due to the increase in red blood  
15 cell counts mediated by the high levels of circulating EPO. Thus, if the MAR improved expression of EPO, higher levels of hematocrit would be expected.

*In vivo* experiments were carried out on 5 week-old C57BL6 female mice (Iffa Credo-Charles River, France). 30µg of plasmid DNA in normal saline solution was delivered by  
20 trans-cutaneous injections in the tibialis anterior muscle. All injections were carried out under Ketaminol (75 mg/kg) and Narcoxyl (10 mg/kg) anesthesia. Following the intramuscular injection of DNA, an electrical field was applied to the muscle. A voltage of 200 V/cm was applied in 8 ms pulses at 1Hz (Bettan M, Darteil R, Caillaud JM, Soubrier F, Delaere P, Branelec D, Mahfoudi A, Duverger N, Scherman D. 2000. "High-level protein secretion into blood circulation after electric pulse-mediated gene transfer into skeletal muscle". *Mol Ther.* 2: 204-10).

16 mice were injected by the Network system expressing EPO without the 1\_68 MAR and 16 other mice were injected with the Network system incorporating the MAR in 5' of  
30 the promoter/enhancer sequences driving the expression of the activator and EPO genes. In each group, half of the mice were submitted to doxycycline in drinking water from the beginning of the experiment (day 0 – the day of electrotransfer) and in the other half, doxycycline was put in drinking water starting at day 21.

35 Blood samples were collected using heparinated capillaries by retro-orbital puncture at different times after the injection of plasmids. Capillaries were centrifugated 10 minutes at 5000 rpm at room temperature and the volumetric fraction of blood cells is assessed in comparison to the total blood volume and expressed as a percentile, determining the hematocrit level.

40 As can be deduced from Fig. 16 The group of mice injected by MAR-network, induced from the beginning of the experiment, display a better induction of the hematocrit in comparison of mice injected by original network without MAR. After 2 months, haematocrits in "MAR-containing group" is still at values higher (65%) than normal  
45 hematocrit levels (45-55%).

More importantly, late induction (day 21) is possible only in presence of MAR but not from mice where the Network was injected without the MAR. Thus the MAR likely protects the transgenes from silencing and allows induction of its expression even after  
50 prolong period in non-inducing conditions.

Overall, the MAR element is able to increase the expression of the therapeutic gene as detected from its increased physiological effect on the hematocrit.

**CLAIMS**

1. A purified and isolated DNA sequence having protein production increasing activity characterized in that said DNA sequence comprises
  - a) at least one bent DNA element,
  - b) and at least one binding site for a DNA binding protein.
2. The purified and isolated DNA sequence of claim 1 characterized in that the bent DNA element contains at least 10% of dinucleotide TA, and/or at least 12% of dinucleotide AT on a stretch of 100 contiguous base pairs.
3. The purified and isolated DNA sequence of claim 2 characterized in that the bent DNA element contains at least 33% of dinucleotide TA, and/or at least 33% of dinucleotide AT on a stretch of 100 contiguous base pairs.
4. The purified and isolated DNA sequence of claims 1 to 2, characterized in that it comprises a MAR nucleotide sequence selected from the group comprising the sequences SEQ ID Nos 1 to 27, a sequence complementary thereof, a part thereof sharing at least 70% nucleotides in length, a molecular chimera thereof, a combination thereof and variants.
5. The purified and isolated DNA sequence of claims 1 to 2, characterized in that it comprises a cLysMAR element and/or fragment, a sequence complementary thereof, a part thereof sharing at least 70% nucleotides in length, a molecular chimera thereof, a combination thereof and variants.
6. The purified and isolated DNA sequence of claim 5, characterized in that said part thereof is a nucleotide sequence selected from the B, K and F regions.
7. The purified and isolated sequence of claims 1 to 6, characterized in that said DNA binding protein is a transcription factor.
8. The purified and isolated sequence of claim 7, characterized in that the transcription factor is selected from the group comprising the polyQpolyP domain proteins.
9. The purified and isolated sequence of claim 7, characterized in that the transcription factor is selected from the group comprising SATB1, NMP4, MEF2, S8, DLX1, FREAC7, BRN2, GATA 1/3, TATA, Bright, MSX, AP1, C/EBP, CREBP1, FOX, Freac7, HFH1, HNF3alpha, Nkx25, POU3F2, Pit1, TTF1, XFD1, AR, C/EBPgamma, Cdc5, FOXD3, HFH3, HNF3 beta, MRF2, Oct1, POU6F1, SRF, V\$MTATA\_B, XFD2, Bach2, CDP CR3, Cdx2, FOXJ2, HFL, HP1, Myc, PBX, Pax3, TEF, VBP, XFD3, Brn2, COMP1, Evl, FOXP3, GATA4, HFN1, Lhx3, NKX3A, POU1F1, Pax6, TFIIA and Vmw65 or a combination of two or more of these transcription factors.
10. A purified and isolated cLysMAR element and/or fragment having protein production increasing activity, a sequence complementary thereof, a part thereof sharing at least 70% nucleotides in length, a molecular chimera thereof, a combination thereof and variants.
11. The cLysMAR element and/or fragment of claim 10 consisting of at least one nucleotide sequence selected from the B, K and F regions.

12. A synthetic MAR sequence comprising natural MAR elements and/or fragments assembled between linker sequences.

13. The synthetic MAR sequence of claim 12, characterized in that the MAR comprises a cLysMAR element and/or fragment, a sequence complementary thereof, a part thereof sharing at least 70% nucleotides in length, a molecular chimera thereof, a combination thereof and variants.

14. The synthetic MAR sequence of claims 12 to 13, characterized in that the linker sequences are BgIII-BamHI linker.

15. A method for identifying a MAR sequence using a Bioinformatic tool comprising the computing of values of one or more DNA sequence features corresponding to DNA bending, major groove depth and minor groove width potentials and melting temperature.

16. The method for identifying a MAR sequence using a Bioinformatic tool according to claim 15, characterized in that said Bioinformatic tool contains algorithms, adapted to the use of profiles or weight-matrices, to compute values for one or more of said DNA sequence features corresponding to DNA bending, major groove depth and minor groove width potentials, and melting temperature.

17. The method for identifying a MAR sequence using a Bioinformatic tool according to claim 16, characterized in that said profiles or weight-matrices are based on dinucleotide recognition.

18. The method for identifying a MAR sequence using a Bioinformatic tool according to claim 17, characterized in that said Bioinformatic tool computes values for all of said DNA sequence features,

19. The method for identifying a MAR sequence using a Bioinformatic tool according to claim 18, characterized in that said Bioinformatic tool is SMAR Scan®.

20. The method for identifying a MAR sequence using a Bioinformatic tool according to claims 15-19, characterized in that the identification of one or more DNA sequence features further comprises a feature corresponding to one or more binding sites for DNA binding proteins.

21. The method for identifying a MAR sequence using a Bioinformatic tool according to claim 20, characterized in that said DNA binding protein is a transcription factor.

22. The method for identifying a MAR sequence using a Bioinformatic tool according to claim 21, characterized in that the transcription factor is selected from the group comprising polyQpolyP domain proteins or transcription factors.

23. The method for identifying a MAR sequence using a Bioinformatic tool according to claims 20 to 21, characterized in that the DNA binding protein is selected from the group comprising SATB1, NMP4, MEF2, S8, DLX1, FREAC7, BRN2, GATA 1/3, TATA, Bright, MSX, AP1, C/EBP, CREBP1, FOX, Freac7, HFH1, HNF3alpha, Nkx25, POU3F2, Pit1, TTF1, XFD1, AR, C/EBPgamma, Cdc5, FOXD3, HFH3, HNF3 beta, MRF2, Oct1, POU6F1, SRF, V\$MTATA\_B, XFD2, Bach2, CDP CR3, Cdx2, FOXJ2, HFL, HP1, Myc, PBX, Pax3, TEF, VBP, XFD3, Brn2, COMP1, Evl, FOXP3, GATA4,

HFN1, Lhx3, NKX3A, POU1F1, Pax6, TFIIA and Vmw65 or a combination of two or more of these transcription factors.

5 24. The method for identifying a MAR sequence using a Bioinformatic tool according to claims 15-23, characterized in that values for the identification of DNA bending are comprised between 3 to 5 °.

10 25. The method for identifying a MAR sequence using a Bioinformatic tool according to claim 24, characterized in that values for the identification of DNA bending are comprised between 3.8 to 4.4 °.

15 26. The method for identifying a MAR sequence using a Bioinformatic tool according to claims 15-25 characterized in that values for the identification of the major groove depth are comprised between 8.9 to 9.3 Å and values for the identification of minor groove width are comprised between 5.2 to 5.8 Å.

20 27. The method for identifying a MAR sequence using a Bioinformatic tool according to claims 26, characterized in that values for the identification of major groove depth are comprised between 9.0 to 9.3 Å and values for the identification of minor groove width are comprised between 5.4 to 5.7 Å.

25 28. The method for identifying a MAR sequence using a Bioinformatic tool according to claims 15-27, characterized in that the melting temperature is comprised between 55 to 75 ° C.

29. The method for identifying a MAR sequence using a Bioinformatic tool according to claim 28, characterized in that the melting temperature is comprised between 55 to 62 ° C.

30 30. The method for identifying a MAR sequence using a Bioinformatic tool of claims 15 to 29, characterized in that it further comprises at least one filter predicting DNA binding sites for DNA transcription factors.

35 31. The method for identifying a MAR sequence using a Bioinformatic tool according to claim 30, characterized in that the filter is applied before or after the Bioinformatic tool.

40 32. The method according to claims 30 to 31, characterized in that the filter detects clusters of DNA binding sites using profiles or weightmatrices.

33. The method according to claim 32, characterized in that the filter detects densities of clusters of DNA binding sites.

45 34. A method for identifying a MAR sequence characterized in that it comprises at least one filter detecting clusters of DNA binding sites using profiles or weightmatrices.

35. A purified and isolated MAR DNA sequence identifiable according to claims 15 to 33 or claim 34.

50 36. The purified and isolated MAR DNA sequence of claim 35, containing at least 10% of dinucleotide TA on a stretch of 100 contiguous base pairs.

37. The purified and isolated MAR DNA sequence of claim 36, containing at least 33% of dinucleotide TA on a stretch of 100 contiguous base pairs.
- 5 38. The purified and isolated MAR DNA sequence of claims 35 to 37, further containing at least 12% of dinucleotide AT on a stretch of 100 contiguous base pairs.
39. The purified and isolated MAR DNA sequence of claim 38, further containing at least 33% of dinucleotide AT on a stretch of 100 contiguous base pairs.
- 10 40. The purified and isolated MAR DNA sequence of any of claims 35 to 39, comprising a sequence selected from the sequences SEQ ID Nos 1 to 27, a sequence complementary thereof, a part thereof sharing at least 70% nucleotides in length, a molecular chimera thereof, a combination thereof and variants.
- 15 41. The purified and isolated DNA sequence of claim 40, comprising a sequence selected from the sequences SEQ ID Nos 24 to 27, a sequence complementary thereof, a part thereof sharing at least 70% nucleotides in length, a molecular chimera thereof, a combination thereof and variants.
- 20 42. The use of a purified and isolated DNA sequence comprising a first isolated matrix attachment region (MAR) nucleotide sequence which is a MAR nucleotide sequence selected from the group comprising:
- a purified and isolated DNA sequence of claims 1 to 9,
  - a purified and isolated MAR DNA of claims 35 to 41,

25 - the sequences SEQ ID Nos 1 to 27,

  - a purified and isolated cLysMAR element and/or fragment,
  - a synthetic MAR sequence of claims 12 to 14,
- a sequence complementary thereof, a part thereof sharing at least 70% nucleotides in length, a molecular chimera thereof, a combination thereof and variants for increasing
- 30 protein production activity in a eukaryotic host cell.
43. The use of the purified and isolated DNA sequence of claim 42, characterized in that said purified and isolated DNA sequence further comprises a promoter operably linked to a gene of interest.
- 35 44. The use of the purified and isolated DNA sequence of claims 42 or 43, characterized in that said purified and isolated DNA sequence further comprises at least a second isolated matrix attachment region (MAR) nucleotide sequence which is a MAR nucleotide sequence selected from the group comprising:
- a purified and isolated DNA sequence of claims 1 to 9,
  - a purified and isolated MAR DNA of claims 35 to 41,
  - the sequences SEQ ID Nos 1 to 27,
  - a purified and isolated cLysMAR element and/or fragment,
  - a synthetic MAR sequence of claims 12 to 14,

40 a sequence complementary thereof, a part thereof sharing at least 70% nucleotides in length, a molecular chimera thereof, a combination thereof and variants for increasing

45 protein production activity in a eukaryotic host cell.

45. The use of the purified and isolated DNA sequence of claim 44, characterized in that said first and at least second MAR sequences are located at both the 5' and the 3' ends of the sequence containing the promoter and the gene of interest.

50

46. The use of the purified and isolated DNA sequence of claim 44, characterized in that said first and or at least second MAR sequences are located on a sequence distinct from the one containing the promoter and the gene of interest.

5 47. The use of the purified and isolated DNA sequence of any of claims 42 to 46, characterized in that said purified and isolated DNA sequence is in the form of a linear DNA sequence as vector.

10 48. A method for transfecting a eukaryotic host cell, said method comprising  
a) introducing into said eukaryotic host cell at least one purified DNA sequence comprising at least one DNA sequence of interest and/or at least one purified and isolated DNA sequence consisting of a MAR nucleotide sequence or other chromatin modifying elements,  
15 b) subjecting within a defined time said transfected eukaryotic host cell to at least one additional transfection step with at least one purified DNA sequence comprising at least one DNA sequence of interest and/or with at least one purified and isolated DNA sequence consisting of a MAR nucleotide sequence or other chromatin modifying elements  
c) selecting said transfected eukaryotic host cell.

20 49. The method of claim 48, characterized in that said DNA sequence of interest is a gene of interest coding for a protein operably linked to a promoter.

25 50. The method of claims 48 and 49, characterized in that the selected transfected eukaryotic host cells are high protein producer cells with a production rate of at least 10 pg per cell per day.

30 51. The method of claims 48-50, characterized in that the MAR nucleotide sequence is selected from the group comprising:  
- a purified and isolated DNA sequence of claims 1 to 9,  
- a purified and isolated MAR DNA of claims 35 to 41,  
- the sequences SEQ ID Nos 1 to 27,  
- a purified and isolated cLysMAR element and/or fragment,  
35 - a synthetic MAR sequence of claims 12 to 14,  
a sequence complementary thereof, a part thereof sharing at least 70% nucleotides in length, a molecular chimera thereof, a combination thereof and variants.

40 52. The method of claims 48-50, characterized in that the MAR nucleotide is a purified and isolated sequence according to claims 1 to 9, a sequence complementary thereof, a part thereof sharing at least 70% nucleotides in length, a molecular chimera thereof, a combination thereof and variants.

45 53. The method of claims 48 to 52, characterized in that the defined time corresponds to intervals related to the cell division cycle.


54. The method of claim 53, characterized in that the defined time is the moment the host cell just has entered into a second cell division cycle.

50 55. A method for transfecting a eukaryotic host cell, said method comprising co-transfecting into said eukaryotic host cell at least one first purified and isolated DNA sequence comprising at least one DNA sequence of interest, and a second and isolated purified DNA comprising at least one MAR nucleotide selected from the group comprising:

- a purified and isolated DNA sequence of claims 1 to 9,
  - a purified and isolated MAR DNA of claims 35 to 41,
  - the sequences SEQ ID Nos 1 to 27,
  - a purified and isolated cLysMAR element and/or fragment,
  - 5     - a synthetic MAR sequence of claims 12 to 14,
- a sequence complementary thereof, a part thereof sharing at least 70% nucleotides in length, a molecular chimera thereof, a combination thereof and variants.
- 10     56.     A process for the production of a protein wherein
- a)     a eukaryotic host cell transfected according to claims 48 to 54 or claim 55, is cultured in a culture medium under conditions suitable for expression of said protein and
  - b)     said protein is recovered.
- 15     57.     A eukaryotic host cell transfected according to any one of claims 48 to 54 or claim 55.
- 20     58.     A cell transfection mixture or kit comprising at least one purified and isolated DNA sequence according to claims 1 to 9, 10 to 11, 12 to 14 or 35 to 41.
- 25     59.     A transgenic organism characterized in that at least some of its cells have stably incorporated at least one DNA sequence of claims 1 to 9, 10 to 11, 12 to 14 or 35 to 41.
- 30     60.     A transgenic organism characterized in that its genome has stably incorporated at least one DNA sequence of claims 1 to 9, 10 to 11, 12 to 14 or 35 to 41.
61.     The transgenic organism of claims 59 and 60 characterized in that some of its cells have been transfected according to the method of claims 48 to 54 or claim 55.
- 30     62.     A computer readable medium characterized in that it comprises computer-executable instructions for performing the method for identifying a MAR sequence of claims 15 to 33 and/or claim 34.


1/15

FIG.1




2/15

FIG.2




3/15

FIG.3



4/15

FIG.4



5/15

FIG.5





FIG.6



6/15

FIG.7





FIG.8



7/15

FIG.9


(A)

Melting temperature (°C)


Bend (degrees)

Major groove depth (Å)

Minor groove width (Å)



(B)



8/15

FIG.10






FIG.11



9/15

FIG.12



10/15

FIG.13






FIG.14




11/15

FIG.15




12/15

FIG.16




13/15

FIG.17




% TA dinucleotide vs Bent DNA




14/15

FIG.18



% TA dinucleotide vs Bent DNA



15/15

FIG.19





FIG.20



SEL PCT 012.ST25  
SEQUENCE LISTING

<110> Selexis S.A.

<120> HIGH EFFICIENCY GENE TRANSFER AND EXPRESSION IN MAMMALIAN  
CELLS BY A MULTIPLE TRANSFECTION PROCEDURE OF MAR SEQUENCES

<130> SEL PCT 012

<150> US 60/513,574

<151> 2003-10-24

<150> EP 04 002 722.9

<151> 2004-02-06

<160> 241

<170> PatentIn version 3.1

<210> 1

<211> 320

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

## SEL PCT 012.ST25

&lt;222&gt; (1)..(320)

&lt;223&gt; MAR of human chromosome 1, nt from 36686 to 37008

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(320)

&lt;223&gt; MAR of human chromosome 1, genomic contig; 36686 to 37008

&lt;400&gt; 1

ttatattatg ttgttatata tattatatta tgttattaga ttatattatg ttgttatatt 60

atataataat attatattat atattatata ttatattata taatatataa taatattata 120

taattatata ttacattata taatatataa taatattata taattatata ttacattata 180

taatataata taatattata taataatata taattatata atatataata atattatata 240

atattatata atattatata atatataaat atataataat atatattata ttatataata 300

gtatataata ttatataata 320

&lt;210&gt; 2

&lt;211&gt; 709

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(709)

&lt;223&gt; MAR of human chromosome 1, nt from 142276 to 142984

## SEL PCT 012.ST25

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(709)

&lt;223&gt; MAR of human chromosome 1, genomic contig; 142276 to 142984

&lt;400&gt; 2

tacaatatat ttctattat atatatttg tattatatat aatatacaat atattttcta 60

ttatatataa tatatttgg attatatata ttacaatata ttgtgatta tataatatat 120

aatacaatat ataatatatt gtattatata ttatataata caatatatta tatattgtat 180

tatatattat atataaact atataatata ttgtattata tattatatat aatactatat 240

aatatatatt attatatatt atatataata caatatataa tatattgtat tataatacaa 300

tgtattataa tgtattatat tgtattatat attatatata atacaatata taataatata 360

ttataatata taataataat ataataaat aataatatat attgtattat atattatata 420

atacaatata taatatattg tattatatat atttattac atataatata taatacatta 480

tataatatat ttgtattat atataatata ttattattg tattatagat aatatatttt 540

attatatatt atatataata caatatataa tatatttgg attgtatata atatataata 600

caatatataa tatattgtat tatatataat attaatatat ttgtattat atatttatat 660

tttatattat aattatggtt tgcattatat attcatatt atatatacc 709

&lt;210&gt; 3

&lt;211&gt; 409

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

## SEL PCT 012.ST25

&lt;222&gt; (1)..(409)

&lt;223&gt; MAR of human chromosome 1, nt from 1368659 to 1369067

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(409)

&lt;223&gt; MAR of human chromosome 1, genomic contig; 1368659 to 1369067

&lt;400&gt; 3

```

tatacataaa tacatatgca tatatatatt gtatatatac ataaatacat atgcatatac    60
attatgtata tatacataaa tacatatgca tatacattat gtatatatac ataaatacat    120
atgcatatac attatgtata tatacataaa tacatatgca tatacattat gtatatatac    180
ataaatacat atgcatatac attatgtata tatacataaa tacatatgca tatacattat    240
gtatatatac ataaatacat atgcatatat tatatacata aattatatta tatacataat    300
acataatacat atattatgtg tatatatata taaatacata tacatatatt atgtgtatat    360
atacatgata catatacata tattatgtat atatatacat aaatacata                409

```

&lt;210&gt; 4

&lt;211&gt; 394

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(394)

&lt;223&gt; MAR of human chromosome 1, genomic contig; 2839089 to 2839482

## SEL PCT 012.ST25

&lt;400&gt; 4

tatgtatata tacacacata tgtatatata cacacatacg tatatacgta tatatgtata 60  
 tatacacaca tatgtatata cgtatatatg tatatatata cacatatgta tatacgata 120  
 tatgtatata tacacacata tgtatatatg tatatatgta tatatacaca catatgtata 180  
 tatgtatata tacacacata tgtatatatg tatatatgta tatatacaca catgtgtata 240  
 tatatatata catatgtata tatgtatata tacacacata tgtatatatg tgtatgtata 300  
 tatacacaca tatgtatata tacacacata tatgtatata tacacacata cttatatata 360  
 cacatatata tgtatatata cacatatgta taca 394

&lt;210&gt; 5

&lt;211&gt; 832

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(832)

&lt;223&gt; MAR of human chromosome 1, genomic contig; 1452269 to 1453100

&lt;400&gt; 5

tatattacta tatatacaat atacatatta ctatatatac catgtattac tatatatatc 60  
 tactatatat attactatat atacaaaata tatattacta tatatacaat atacatatta 120  
 ctatatatac catatattac tatatatatc tactatatat attactatat atacaaaata 180  
 tatattacta tatatactat atattactgt atatacaata tatattacta tatatatact 240  
 atatattact atatatacac tatatattac tatatatata caatatatat attactatat 300

## SEL PCT 012.ST25

atacacaatg tatataacta tatatacaat atatattact atatatacta tatatattac 360  
 tatacatact atatattact ctatatatac aatatataa ttacaatata tactacatat 420  
 tactacatat actttatata ttactatata tactatataf tactgtatat acaatatata 480  
 ttactaaata tacacaatat atattactat atataca~~caa~~ tatatatatt actatatata 540  
 cacattatat atgactatat atacacacta tatatattac tatatataca caatatataa 600  
 ctatatatac acagtataca tattactata tataca~~caat~~ atatatatta ctatatatac 660  
 actatatatt actatatata cacaatatat attactctat gtatacacta tatatattac 720  
 tatatataca gaatatatat aactatatat acactat~~att~~ actatatata ctatatatta 780  
 ctatatgtac tatatatatt actatatata ctatatatta ctatatatac ac 832

&lt;210&gt; 6

&lt;211&gt; 350

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(350)

&lt;223&gt; MAR of human chromosome 1, genomic contig; 831495 to 831844

&lt;400&gt; 6

aatatataat atataaatat taatatgtat tatataat~~at~~ atattaatat attatattat 60  
 attactatat aaataatatt aatatattat attaaa~~at~~at taataaatat atcatattaa 120  
 atattatatt aattaaatat taataaatat attatatta~~a~~ tatatttata tattaaacct 180  
 ataacatatg catatactta ttatatata acatgcat~~gt~~ acttatttat atatacaata 240  
 tatatttata tattatataa tatattatat gtatttatat attatata~~tc~~ atatattata 300

## SEL PCT 012.ST25

tgtatttata tattatatat catataatat atatatttat attatatata 350

<210> 7

<211> 386

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(386)

<223> MAR of human chromosome 1, genomic contig; 1447225 to 1447610

<400> 7

acatttaatt taattatata ctgctatata taattaaatc tatatatcta tataacttat 60

aatttatatt aatttaatta tatatactat atagtatat atacatatat gtaattatat 120

atagtataat tatagtatat atgtatatat aatgtaagta aatatatagt atatatttat 180

atatactata tatttataca tatgtcttta tatatactaa tatatataca catatgtaat 240

atgtacatat ggcatatatt ttatagtgtat tatatacata tatgtaatat atatagtaat 300

atgtaaatat atagtacata ttttaattata tggtaatatata tacacatatata tgtaatatgt 360

gtatttatagt acatatttta tagtat 386

<210> 8

<211> 585

<212> DNA

<213> Homo sapiens

<220>

## SEL PCT 012.ST25

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(585)

&lt;223&gt; MAR of human chromosome 1, genomic contig; 4955365 to 4955949

&lt;400&gt; 8

atacacacat atacacatat gtacgtatat atactatata tacacacata tacacatatg 60  
 tacgtatata tactatatat acacacatat acacatatgt acgtatatat actatatata 120  
 cacacatata cacatatgta cgtatatata ctatatatac acacatatatac acatatgtac 180  
 gtatatatac tatatatata cacatatata catatgtacg tatatatatt atatacacac 240  
 atatacacat atgtacgtat atatactata tatacacaca tatacacata tgtacgtata 300  
 tatactatat atacacacat atacacatat gtacgtatat atactatata tacacacata 360  
 tacacatatg tacgtatata tactatatat acacacatat acacatatgt acgtatatat 420  
 actatatata cacacatata cacatatgta cgtatatata ctatatatac acacatatatac 480  
 acatatgtac gtatatatac tatatatata cacatatata catatgtacg tatatatact 540  
 atatataccc atacacatac gtatatacgt acatatatat acgta 585

&lt;210&gt; 9

&lt;211&gt; 772

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(772)

&lt;223&gt; MAR of human chromosome 1, genomic contig; 5971862 to 5972633

## SEL PCT 012.ST25

&lt;400&gt; 9

agtaaacata tatatagtaa atatatatag tgtatatata gtaaatatat atagtgcata 60  
tatatagtc atatatatag tgtatatata gtaaatatat agtgtatata tatagtaa 120  
atatatagtg tatatatagt aaatatatat agtaaata tatatactat atatagtaaa 180  
tatatatata ctatatatag taaatatata tatagtatat atatagtaaa tatatatata 240  
gtatatatat agtaaata tatatatag atatatatag aatatatata tagtatatat 300  
agtaaata tatagtatat atatagtaaa tatatatata gtatatatat agtaaata 360  
tatatagtat atatatatag aatatatata tagtatatat atagtaaata tatatagtat 420  
atatatatag aatatatata gtatatatat agtaaata tatagtatat atatagtaaa 480  
tatatataca ctgtatatat atagtaaata tatatact gtatatatat agtaaata 540  
tatatactgt atatatatag taaatatata tacactgtat atatatatag aatatatata 600  
cactgtatat acatagtaaa tatatataca ctgtatatat atagtaaata tatatact 660  
gtatatatat agtaaata tatatactgt atatacatag taaatatata tacagtgtat 720  
atacatagta aatatatata cagtgtatat acatagtaaa tatatataca gt 772

&lt;210&gt; 10

&lt;211&gt; 304

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(304)

&lt;223&gt; MAR of human chromosome 1, genomic contig; 6221897 to 6222200

&lt;400&gt; 10

## SEL PCT 012.ST25

atatataata tatataatta tattatatat aatatataat atatataatt atattatata 60  
 ttatatataa tatattatat attatatata taatatatat tatatattaa atatatatta 120  
 tatatataat atatattata tattaaatat atattatata tataatatat attatatata 180  
 atatatataa tatatattat atatatatta tatattatat atatatatta tatatatata 240  
 atatatataa tatatattat atataatata tattatatat atataatata tataatatat 300  
 atta 304

&lt;210&gt; 11

&lt;211&gt; 311

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(311)

&lt;223&gt; MAR of human chromosome 1, genomic contig; 9418531 to 9418841

&lt;400&gt; 11

tatatataat atttatatat aatattcatg tatttatata taaatattta tatatttata 60  
 tataaatatt tatatattta tatataaata ttatatatt tatatataat atttatacat 120  
 tatatataat atttatatat tatatataat atttatatat aatatttata tattatatat 180  
 aatatttata tatttatatg tataatatat attttatata tgtatgtata atatatattt 240  
 tatatatgta tgtataatat attttatata tgtatgtata atatattatt atatataata 300  
 tataatttat a 311

&lt;210&gt; 12

&lt;211&gt; 302

## SEL PCT 012.ST25

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(302)

&lt;223&gt; MAR of human chromosome 1, genomic contig; 15088789 to 15089090

&lt;400&gt; 12

atataatata tatattatat atataaatat atataaatat ataacatata tattatatat 60

aaatatatat aaatatataa catatatatt atatataata atatataata atatataaca 120

tatatattat atatataaat atatataaat atataacata tatattatat atataaatat 180

atattatata ttatatata taatatatat aaatatataa tatatattta tatatataat 240

atatataaat atataatata tatatttata tataatatat ataaatatat aatatataat 300

at 302

&lt;210&gt; 13

&lt;211&gt; 461

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(461)

&lt;223&gt; MAR of human chromosome 1, genomic contig; 6791827 to 6792287

## SEL PCT 012.ST25

&lt;400&gt; 13

tatataatat atattatata tacacatata taatatatat tatatatata catatataat 60  
 atatattata tatacacata tataatatat attatatata cacatatata atatattata 120  
 tatatacaca tatataatat atattatata tacacatata taatatatat tatatatata 180  
 catatataat atatattata tatacacata tataatatat attatatata cacatatata 240  
 atatattata tatatacaca tatgtaatat atattatata cacacatata atatattata 300  
 tatacacata tataatatat attatatata catatataat atatattata tatacacata 360  
 tataatatat attatatata cacatatata atatattata tatatacaca tataatatat 420  
 aatatataca catatataat atatattata tatatgcaca t 461

&lt;210&gt; 14

&lt;211&gt; 572

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(572)

&lt;223&gt; MAR of human chromosome 1, genomic contig; 163530 to 164101

&lt;400&gt; 14

atattataat tatatatatt atatataatt atataaaata tatattataa ttatatatat 60  
 ttatatataat atatattata taattaatat attatatata atatatatat tatatatata 120  
 atatattata tatatatatt atataatata tataatatat ataatatata atataatata 180  
 tatattatat ataatatata atatatatata tatattataa tataatatat ataatatata 240  
 atataatata tataatatat aatatatat ataatatata atatatatata tatataatat 300

## SEL PCT 012.ST25

aatatataat atataataa tataatataa tatataatat atataatata ttataatata 360  
 atatatataa tatataatat aatatatata atataata taatatataa tatataatat 420  
 atatttaata tatttattaa ttatttgta tatatttatt aatatataat atataatata 480  
 tttaatatat tataactata tattatatta taattatata tattatatat atacaattat 540  
 aattatatat tatatatact tataatatat at 572

<210> 15

<211> 357

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(357)

<223> MAR of human chromosome 1, genomic contig; 1842332 to 1842688

<400> 15

tatatctata tatatctata tatatataat atagataata tctatatata taatatagat 60  
 aatattatct atataataa tagataatat tatctatata taatatagat aatattatct 120  
 atatataaaa ttatattata tctatatata ttatatatat aaaattatat tatatctata 180  
 tataatatag ataatatcta tatataaata gataatatct atatatataa tatagatatt 240  
 atctatatta tagatataga taatattatc tatattatag atattatcta tatataatat 300  
 agataatatt atctatatta tatatataat atatctatat tatctataat attatct 357

<210> 16

<211> 399

<212> DNA

## SEL PCT 012.ST25

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(399)

<223> MAR of human chromosome 1, genomic contig; 2309560 to 2309958

<400> 16

attatatata atatataa tatattatat atatcaagca gcagatataa tatataatat 60

atataatata tataatatat attgtatatt atataatata taatatatat aatatatatt 120

gtatattata taatatataa tatatataat atatattgta tattatataa tatataatat 180

atataatata tattgtatat tatataatat ataatatatg taatatatta tgtaatatat 240

tatataatat atattatata ttatatataa tatatattat atataatata tattacataa 300

tatattacat atattacgta atatatgta tatattacat ataatatata acatatatta 360

cgtaatatat gtaatatatt acatataata tatacatta 399

<210> 17

<211> 394

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(394)

<223> MAR of human chromosome 1, genomic contig; 2231759 to 2232152

## SEL PCT 012.ST25

&lt;400&gt; 17

atatatactt ataaattata tacttatata tacttataaa ttatatactt atatatactt 60

ataaattata tacttatata tacttataaa ttatatactt atatatactt ataaattata 120

tacttatata tacttataaa ttatatactt atatatactt ataaattata tacttatata 180

tacttataaa ttatatactt atatataatt ataaattata tacttatata taattataaa 240

ttatatactt atatataatt ataaattata tacttatata taattataaa ttatatactt 300

atatataatt ataaattata tacatatata taattataaa ttatatacat atataattat 360

aaattatata catatatataat tataaattat atac 394

&lt;210&gt; 18

&lt;211&gt; 387

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(387)

&lt;223&gt; MAR of human chromosome 1, genomic contig; 7406524 to 7406910

&lt;400&gt; 18

tatattatat ataatatata ttatatataa tataaataat atatattata tataatatat 60

aaataatata taatatataa ataatatata atatataata tataaataat atataatata 120

taacatatata ataatatata taatatataa ataatatata taatatataa ataatatata 180

taatatataa aaatatataa tatataatac atatataaat aatatattat attatatatg 240

atacataata tattatatat aatatattat atgatacata atatattata tagaatatat 300

tatatgatac ataatatatt atatagaata tattatatga tacataatat attatatgat 360

SEL PCT 012.ST25

acataatata ttatatataa tatatta

387

&lt;210&gt; 19

&lt;211&gt; 370

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(370)

&lt;223&gt; MAR of human chromosome 1, genomic contig; 9399572 to 9399941

&lt;400&gt; 19

catatatata tatatacaca tatatacaca tatatatata catacatatg tacacatata 60

tatacacata tgtatacaca tatatacaca tatatacaca catatatata catatatata 120

cacatatata cacatatata cacatatata cacatatata catatatata catatatata 180

tatatacaca tatatatata atacacacat atatatacac atatatacac acatatatac 240

acatatatac acatatatat acacatatat acacatatat acatatatac acatatatat 300

acatatatac acatatatac atatatacac atatatacat atatacacac atatatacac 360

atacatatac

370

&lt;210&gt; 20

&lt;211&gt; 377

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

## SEL PCT 012.ST25

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(377)

&lt;223&gt; MAR of human chromosome 1, genomic contig; 12417411 to 12417787

&lt;400&gt; 20

attatatata ataatatata ttatatattt atatatataat tataataaat acatatataatt 60

atatatttat atataaatta tatataataa ataatatata ttatatatat ttataaatta 120

taataaatac atataattac atatatattat atataaatta tatataataa ataatatata 180

ttatatatat ttatatgtag attatatata aatatatata attatatatat ataataatat 240

atataattta tatatatata atatatatata ataatatat ataatttata tatataatta 300

tatatatata ataatatata taatatatat aatttatata tataattata tatataataa 360

atatatatata ttatat 377

&lt;210&gt; 21

&lt;211&gt; 1524

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(1524)

&lt;223&gt; MAR of human chromosome 1, genomic contig; 1643307 to 1644830

&lt;400&gt; 21

tataaatata tataaatata taaatatata taaatatata aatatatata aatatatata 60

aatatatataa aatatataa tatatatataa tatatatataa tatataaaaa cataaaaaata 120

## SEL PCT 012.ST25

tatataaata tatataaata tataaaaata tataaatata taaatatata aaaatatata	180
aatatataaaa tatatacata aatatatata aatatatata aatatataaaa aatatatata	240
aatatataaaa tatatataaaa tatatataaaa tatatataaaa tatataaaaaa tatatataaaa	300
tatataaata tataaaaata tatataaata tataaatata taaatatata taaatatata	360
aatatataaaa taaatatataag tatttatgaa tatatatgaa tatataaata tataaaaaaat	420
atatataaat atataaatat atataaatat ataaatatat acatatatac atatataaat	480
aaataaatat aagtatttat gaatatatat gaatatataa atatataaaa aatatatata	540
aatatataaaa tatatataaaa tataaatata taaaaatata taaaaatata tataaatata	600
taaatatata taaatatata aatatatata aatatatata aatatataaaa tatatataaaa	660
tatatataaaa tatataaata tataaatata tataaatata tataaatata taaatatata	720
aatataaata tataaatata tataaatata tataaatata taaatatata taaatatata	780
taaatatata taaatatata taaatatata aatatatata aatatatata taaatatata	840
taaatatata aatatataaaa tatataaaaaa tatataacaa tatataaata tatataaaaaa	900
tatataacaa tatataaata taaatatata taaaaatata taacaatata taaatatataa	960
tatatataaaa tatataaata taaatatataa aaatatatat aaatatataa atatataata	1020
atatataaat gtataaatat atataaaaaat atataacaat atataaatat ataaatatat	1080
aacaatatat aaatatataa aaatatataa caatatataa atataaatat atataaaaaat	1140
atataacaat atataaatat aaatatatat ataaatatat aaatatataat ataaaaaata	1200
tatataaata tataaatata tatataaata tatataaata tataaatgta taaatatata	1260
taaatatata aatatataaaa aatatataaaa tatatataaaa tatatataaaa tatataaata	1320
taaatatata aatatatata aatatataaaa tataaatata taaacatata taaatatata	1380
taaataaaca tatataaaga tatataaaga tataaagata tataaatata taaatatata	1440
aagatatata aatatataaaa gatatatataa tatataaaga tatataaata tataaagata	1500
tataaatata atatataaat atat	1524

## SEL PCT 012.ST25

&lt;211&gt; 664

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(664)

&lt;223&gt; MAR of human chromosome 1, genomic contig; 1398763 to 1399426

&lt;400&gt; 22

```

acacatatat atataaaata tatatatata cacacatata tataaaatat atatatatat 60
acacatatat ataaaatata tatatacaca catatatata aaatatatat atacacacat 120
atatataaaa tatatatata cacacatata tataaaatat atatatacac acatatatat 180
aaaatatata tatacacaca tatatataaa atatatatat acacacatat atataaaata 240
tatatatata cacatatata taaaatatat atatacacac atatataaa aatatatata 300
tacacacata tatataaaat atatatatat acacatatat aaaatatata tatacacaca 360
tatataaaat atatatatat acatatatat aaaatatata tatacacata tatataaaat 420
atatatacac acatatatat aaaatatata tatacacaca tatatataaa atatatatat 480
acacatatat ataaaatata tatatacaca tatatataaa atatatatat atacacatat 540
atataaaata tatatacaca catatatata aagtatatat atacacacat atatataaaa 600
tatatatata cacatatata taaaatatat atatacacat atatataaaa tatatatata 660
caca                                     664

```

&lt;210&gt; 23

&lt;211&gt; 1428

&lt;212&gt; DNA

## SEL PCT 012.ST25

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(1428)

&lt;223&gt; MAR of human chromosome 2, genomic contig; 17840365 to 17841792

&lt;400&gt; 23

```

aatftattat atattatata ttatatatat tatatatatt atatattata tatattatat   60
atattatata ttatatatat tatatattat atatttatat ataatatata tctaatatat   120
atattagata taatatatat ctaatatata tatattttat atatataata tatctctaata   180
atatatattt tatatgtata taatatatct ctaatatata tatattttat atgtatataa   240
tatatctcta atatatatat ttttatata taatatatct ctaatatata tattttatat   300
atataatata tatctaata atataatata tatattagat atatataaaa tatatatgat   360
atatttatta tatatataat atataatata taatatatat attatattat atacatatat   420
attatataca atatatatta tatatatttt atatacatia tatattatat atattttata   480
tacaatatat attatatatt ttatatacaa tatatattat atatatttta tatttttata   540
tacaatatat attatatata ttttatatat aatatatatt atatatattt tatataatat   600
atattatata tatttttatat ataatatata ttatataaat tatatataat atatattata   660
ataaattata atatttttta tatatataat atgtatttta tatataatat attataatat   720
atattttata tataatatat tataatatat attttatata taatatatta taatatatat   780
tttatattat aatatattat aatatatatt ttatatataa tatattataa tatatatttt   840
atatataata tattataata tatattttat atataatata ttataatata tatattataa   900
tatatatttt atatataata tattatcata tatatattaa atatatattt tatatataat   960
atattataat atatatatta taatatatat ttatatata atatattata atatatatat  1020

```

## SEL PCT 012.ST25

tataatatat attttatata taatatatta taatatatat ttatatata atatattata 1080  
 atatataatt tatatatata atattataat atatatttta tatataatat aatatatatt 1140  
 ttatatataa tatattataa tatatatatt atatataata tattataata tatattttat 1200  
 atataatata ttataatata tttttatat ataatatatt ataatatata ttttatatat 1260  
 aatatattat aatatatatt ttatatataa tatattataa tatatatatt atatataata 1320  
 tattataata tatattttat atataatata ttaattaaat ttattaattt attaattatt 1380  
 aatatttatt atattattaa ttaataatat ataaattatt aatatata 1428

&lt;210&gt; 24

&lt;211&gt; 4624

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(4624)

&lt;223&gt; MAR 1\_6 of chromosome 1

&lt;400&gt; 24

ggatcttaaa tctattttat ttatttattt ttcatgtggc caataccctc cacccttc 60  
 ttctgtctct tcaacttat tgtggttacc ttgaggctac ctgagacagt aggcttgggt 120  
 ggggaagtat gcattctaag tgtaaagttt gatgagcttt gacaaatgtc aacctatgta 180  
 ccagaacatt ttcatcacc ataaaatctc ccttgtgtca ctgcccagtc agtgtctatt 240  
 ctagtatcca actcctggct ccaagaaacc attgaactgt ttctgtcac tataaattag 300  
 atttgtcttt tctagagttt catgtaaatg gaatcataca ctaagtactc ttgtgcctg 360  
 gcttctgctc agcataatgt ttfgagaat cattcatgct gctgcatgtt tcagtagtt 420

## SEL PCT 012.ST25

catttttta aataggtgaa ttgtaactca ttctgtgaat ataccatatt ctgtcttcca 480  
tttatctgtt agtggatctt taggtcgttt ctagtgttg gctattgcaa ataaagctgc 540  
tgtaaataatt aatgcacaag tttccatgt tcatatgttt catttcactt aggaaaatac 600  
ctaagagagg aattgcacat attaaaaaaa ttttaaaaac tactaagctg ttctccaaaa 660  
tgggtgtaca atttttattc ccaagagcaa tatgagtgtt taattgctcc acattctcac 720  
caacacttgg tgcttgttag ttttatttc attgttttca ttgttatgtc tgtgaggcag 780  
cattgatgtg catgtctctg agtgtcatct tagcgggtgat gctgagcatc agttcacgtc 840  
cttataggcc gttgtatat ctgcttttg aaatgtctgt tcaaattctt tgcctatttt 900  
aaattgagtt gtgttcgtct tcttaggatt aagtaatgag ttaaaaatat ttctgatata 960  
aatctttcat tatatatttc taatgcttcc tcatctatag tttattttct catattcttt 1020  
aactgtatct ttgaagagc aaattttact ttgattatg cccaatttat caagttttta 1080  
tatggctctt ttgattatgc ccataatcac attagacttt gcctaacca agtttgacaga 1140  
gattttttct tttatgcttt tatctagaaa tttgtagtt ttaggtttta aaaaagttta 1200  
atttatttat ttgagacagg gtattgctct ttacatatac tggagtgcag tgatgcaatc 1260  
atggctcact gcagccctcaa cctcttgggc tcaagcgggt ctcccatctc agagtcctga 1320  
gtagctggcc aggtgcatgc cagcttcaat gtgttttca ttgcatttc cctgataatt 1380  
attgacgttg agcatttttt tcatatatca gttagctatt tgtacgtctt cttttgagaa 1440  
acatctattc gggctctttg cccattttta agtcagatgg ttgtttgtc agctattgag 1500  
ttgtttgagt tcttgtata ttctggatat taccatcttg tcagatgcac agtttgcaaa 1560  
tttttttt ctattttgta ggttgtctct ttctctgttg ttctctccgg tatgcagaag 1620  
tttttagtg tgatgtaatt tcatttgtct gtttttgctt ttgtgcctg tactttctta 1680  
ttcttatcca aaaaattctt atctagatca atgtcacgaa gagtttctcc tctgttttct 1740  
tcgagtagtt tttataatt ttgggtatac atttaagtct ttaatctatt tggaattgat 1800  
tttgcataat ggtgagagat cagagtctaa ttcatactt ttggatgtgg aaagctagtt 1860  
tttcagcac catttattga agagactgtc tcttctccaa tgtgtgttct ttgtgccttc 1920

## SEL PCT 012.ST25

gtcaaaaatc agttggctgt gcggtggattt atttctgtgt tctctatttt gttccattgg 1980  
 tctagtftta gccttaaatt taggtctgca atttttttt tttgtatat ggtgtgaagt 2040  
 aagagtcaaa gttcattatt ttcatatgg atatgtaatt actccagtac catcatttag 2100  
 ttgaatgga ctgtccttic tccatggaat tacatgggca tctttgtct gaaaccaatt 2160  
 atgtatgttt acgtatgtgt atgtttatgc atatgttata ggtttaatat atattaatat 2220  
 atataatata taatatataa atattaatat gtattatata atatatatta atatattata 2280  
 ttatattact atataaataa tattaatata ttatattaaa atattaataa atatatcata 2340  
 ttaaataa tattaattaa atattaataa atatattata ttaatatatt tatatattaa 2400  
 acctataaca tatgcatata cttatttata tataacatgc atgtacttat ttatatatac 2460  
 aatatatatt tatatattat ataatatatt atatgtattt atatattata tatcatatat 2520  
 tatatgtatt tatatattat atatcatata atatatatat ttatattata tatattatat 2580  
 gatataaat attatataat gtattaatat atattaaacc tatatttata attctggact 2640  
 cactatttg ttccattgggt gtctgtgtgt atctaaccct atgccaataa tgtactatct 2700  
 taattacat agctttatag taagcttga aatcagatag tgtattttt atcattgttt 2760  
 ttaaaataa tagtttatct tttatttga atttgaatc agctagtcag tttctgcaa 2820  
 aagcttactg ggattttgct tggaattatg ttacatctgt agcatgtact atccaatatt 2880  
 ctagccttta tccacatgtg gctattaagg tttaaattaa ttaaattaa atttaattaa 2940  
 ttaaattaa aacttaataa ttggttcctc attcacacta ccatatgtca agtgttcaat 3000  
 agccacatat ggtcaatgtc ttggaaaagt caatacagta catttccatt attgcagtaa 3060  
 gttctgtcaa acagcactat cgtagaccga ttaggagaga actgacttaa cagtattgga 3120  
 tgctccagtc aatgaacatc tttttttt tcatatttt cagtagtctc tgcagtatat 3180  
 tatagatttc agttacata tttgcatat attttattaa atgtataacg gtagaagtac 3240  
 tattattgga tgatgtgtc tatagatgta ttttaggtca agtttgtga tagtgtgtt 3300  
 taaatctcgt atacctctg attttttat ttactgttc ttgaattac tgagacagga 3360  
 atgttatatc ctttaactata tttgtgaatt tattcacttc ttccctcagt tctgttaact 3420

## SEL PCT 012.ST25

ttgcttagg tgcttttaa aaatgaaact ttcaatctct gcctttaat ttagcattt 3480  
 agaccattta cattcaatgt aattatcaat atcagtttat ttaagtctga agttgtgcaa 3540  
 ttttcctct acctatatta taaatcttc tatatacaaa acacatgcta tgtttctgc 3600  
 atatgtttta aatgacaccc ggaaagcatt gacactattt ttgcttagg ttatctttca 3660  
 aagatgttaa aaatgagaaa gaaatattct gcatttatcc atacacttat tatttgcaaa 3720  
 ggtttttta aataccttg ttagatttc agttaccaac ttgtattcc ttcagctga 3780  
 agaacttaca atttctgta ggacaggctct ctgacaacaa attatctcag ctttcttg 3840  
 tctaaaaaag ttattgcctt ttttttaaa atatatcttc actggatatt gaatttagg 3900  
 tgataatctt tttttttg ttagcattt aaatatgtct tctaattgcc tcttgcttc 3960  
 atagttctg atgagaagtc tactgttatt agtatctctt tgtgtgtgc tctctttt 4020  
 ccctctctgc tattatggct attttttt tttttttt ttttggtcac tgggtgcagc 4080  
 aatttaatta tgggtgcct tggatgttt tgtgtgtgtg tgtgtgtgtg tgtgtgtgtg 4140  
 tgtgtgtgtg tagctgatgt tcttgagct ttagaatctg tgagtttga gtttcatca 4200  
 attattttt ctttcatc ctttattta ctcatgttcg tgtttatt tatatttta 4260  
 agaatttgt gcgtattgt aataactgtt taaatgtcat ttgtgaattc cattgcttct 4320  
 aggtaggatt ctattgacag atatttttc cctgacgaga ggtcatactt tcctattct 4380  
 tcatgtatct agtggtttt ggtgaatac tggatattt gaatttatg ggagtgtga 4440  
 attctacaat attcctaaa aatgtgttg atttgttt agcagatagc tatcttactt 4500  
 gaagatcaat tcatatttt ttgatgttca ttttcatc tattaaagaa taggtccatg 4560  
 gtagagtta ctgatatcaa ccttctggt gtctctaata aatgcaacat attcaataag 4620  
 atcc 4624

&lt;210&gt; 25

&lt;211&gt; 3616

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

## SEL PCT 012.ST25

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(3616)

&lt;223&gt; MAR 1\_68 of chromosome 1

&lt;400&gt; 25

gactctagat tataccaacc tcataaaata agagcatata taaaagcaaa tgctcttata 60  
ttgcagatcc ctgaactgag gaggcaagat cagtttgga gttgaagcag ctggaatctg 120  
caattcagag aatctaagaa aagacaaccc tgaagagaga gaccagaaaa cctagcagga 180  
gtttctcaa acattcaagg ctgagggata aatgttacat gcacaggggtg agcctccaga 240  
ggcttgcca ttagcaactg ctacagtttc attatctcag ggatcacaga ttgtgctacc 300  
tattgcctac catctgaaaa cagttgcttc ctatattca tccagtttaa tatttattta 360  
aaccaagaag gftaatctgg caccagctat tccgttgga gtggatgtga aagtaccaat 420  
tccattctgt ttactatta actatccttt gccttaatat gtatcagtag gtggcttggt 480  
gctaggaaat attaaatgaa tggcatgttt catagggtgt gtttaaagtt gtttttgag 540  
ttaaatcttt cttaataat actttctgat gtcaaaaaca cttagaagtc atgggtgtga 600  
acatctatat agggttggat ctaaaatagc ttctaacct ttctaacca ctgttttgt 660  
ttgttgttt ttaactaagc atccagttg ggaaattctg aattagggga atcataaaag 720  
gtttcatttt agctgggcca cataaggaaa gtaagatatc aaattgtaa aatcgtaag 780  
aactctatc ccactgaag tgggggttag gtgcctcttc tctgtgctcc ctaaacatcc 840  
tattttatct gtatatatat atattcttcc aaatatccat gcatgggaaa aaaaatctga 900  
tcataaaaaat atttaggct gggagtggtg gctcacgcct gtaatccag cactttggga 960  
ggctgaggtg ggcggatcat gaggtaaga gatcgagacc atcctgacca atatggtgaa 1020  
accccatctc tactaaagat acaaaaactat tagctggacg tggtggcacg tgctgtagt 1080

## SEL PCT 012.ST25

cccagctact cgggaggctg aggcaggaga acggcttgaa cccaggaggt ggagggtgca 1140  
 gtgagctgag atcgcgccac tgcactccag cctgggcgac agagcgagac tctgtctcaa 1200  
 aaaaaaata tatatatata tatatatata catatatata taaaatatat atatatacac 1260  
 acatatatat ataaaatata tatatatata cacatatata taaaatatat atatatacac 1320  
 acatatatat aaaatatata tatacacaca tatatataaa atatatatat acacacatat 1380  
 atataaaata tatatatata cacatatata taaaatatat atatacacac atatataaa 1440  
 aatatatata tacacacata tatataaaat atatatatat acacatatat ataaaatata 1500  
 tatatacaca catatatata aaatatatat atacacacat atatataaaa tatatatata 1560  
 cacacatata tataaaatat atatatacac acatatataa aatatatata tacacacata 1620  
 tataaaatat atatatacac atatataaa aatatatata tacacatata tataaaatat 1680  
 atatacacac atatataaa aatatatata tacacacata tatataaaat atatatatat 1740  
 acatatatat aaaatatata tatacacata tatataaaat atatatatat acacatatat 1800  
 ataaaatata tatacacaca tatatataaa gtatatatat acacacatat atataaaata 1860  
 tatatatata catatatata aaatatatat atacacatat atataaaata tatatatata 1920  
 catatatata aaaatatata tatatatatt ttaaaatatt ccaattgtct cactttgtgg 1980  
 atgagaaaaa gaagtagtta gaggtcaagt aactggcct acatctttc tcaagattgt 2040  
 aaactcctag tgagcaataa ccacatcttc atttctttg tataaaacaa gaaagttag 2100  
 catgaaaaag gtactcaatt acaaatgtgt tggattgaat tgaagaccct tggaagggga 2160  
 tttgtacct gaggatctct ttctttggc catatgttc aatggacaaa atttagcctt 2220  
 cgaaggcagg cggattgag gtaataacta cctttaccac ttgatagcta tgtgaccttg 2280  
 gccatgtggt ttcaacagtc tgaacctcat ttctctgtg tatgtgtggt cctccttaca 2340  
 agttgtgaa aaatgtgaag tccttagcca tgatagcca atataacagg cttaatgata 2400  
 ataggtttat gttctttcc ttatatctc cagataagca ctgtccaagt ttgaggtgtt 2460  
 ttgaggtctc gcctgattg gattgttga gttatgcta ttcttgaat tcttgagct 2520  
 gttctgaagc agtgtatcat gaacaaaaac atccccagtt cagtccaaac ccttggttac 2580

## SEL PCT 012.ST25

atatcattct tatgccaagt tataaccagt ttgagagtg tccctctgtt attgcattta 2640  
 agtttcagcc tcacacagaa attcagcagc caatttctaa gccctaagca taaaatctgg 2700  
 ggtggggggg ggggatggcc tgaagagcag cattatgaat agcaccatta taattaatga 2760  
 tctctcagga agatttaca tcacaggtag cagataaaac aaatagtact gcttctgcac 2820  
 ttccctctct ttattcgtct atgaaatttt atgggaaatc agtcagtgta aaaatgtaag 2880  
 ctctaatct tcccagaaa tctacctca ttgatgaat acttgaggg aatgaattag 2940  
 agcattttt tctttatag tctactcgc atttacgaag tgaggacggg agcttaggct 3000  
 gcctggccaa ctgatgagaa ggtcagaggc attttagag acctctgttg tcttcattc 3060  
 atgttcattt tccacaaggc aagtaatttc caacaaatca gtgtcttcat tagtaataag 3120  
 attattaaca acaataatag tcatagtaac tattcagtgta gagtcatta tataacaggc 3180  
 attctacaag gtactttata tacatctgag taaacctcac acaattctac agggaggtat 3240  
 ttctatcccc atttaacaaa taaggaaacg aagtccaagt aaattaactt gccaaggctc 3300  
 acacagatag tacctggcag aacaggaatt taaacctaaa ttgtccaac tccaaaagca 3360  
 gccttctatt tgtataaat gctgcctctc attatcacat atttattat taacaacaac 3420  
 aaacatacca attagcttaa gatacaatac aaccagataa tcatgatgac aacagtaatt 3480  
 gttatactat tataataaaa tagatgtttt gtatgttact ataacttga atttgaatag 3540  
 aaatttgcatt tctgaaagc atgttcctgt catctaatat gattctgtat ctattaaaat 3600  
 agtactacat ctagag 3616

&lt;210&gt; 26

&lt;211&gt; 4660

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

## SEL PCT 012.ST25

&lt;222&gt; (1)..(4660)

&lt;223&gt; MAR 1\_42 of chromosome 1

&lt;400&gt; 26

gatcccttga ggtcagtagt ttaagaccag cctgaccaac atggtgaaac ccatctctac 60  
taaaaataca aaaattagcc aggcgtgggtg gcgggggcct gtaaccccag ctactcagga 120  
ggctgaggca caagaatctc ttgaaccggg gaggcggagg ttgcagttag ctgagattgt 180  
gtcactgcac tccagcctgg gcaacagtgc cagactctgc cttaaaaaaa aaaaaaaaaa 240  
aaaaaaggcc gggcgcggtg gctgacgcct gtaatcccag cactttggga ggccgaggcg 300  
ggtggatcat gaggtcagga gatcgagacc acagtgaaac cccgtctcaa ctaaaaatac 360  
aaaaaattag ccgggcgcggtg tgggtggcggtg ttgtagtccc agctactcag gaggctgagg 420  
caggagaatg gcgatgaacct gggaggcgga gcttgcagtg agccgagatg gcaccactgc 480  
actccagcct gggcgaaaga gtgagactcc gtctcaaaaa aaaaaaaaaa ttagctgggt 540  
atggtgggtg gtgcctgtaa tcccagctac tcgggagggt gaggcaggag aatcccttga 600  
acctgggagg cggagggtgc agtgatctgc catcctgtca ctgcatcact acactccagc 660  
ctgggtgaca gagcgagact ctgtctcaa aaaaaaaaaa aaaaaaaaaa ctgggtgtgg 720  
tggtatgcac cagctgtagt cccagctact tgggaggctg agttgggggg attgcctgag 780  
ccaggagggt cgagggttca gggagccatg attatgccac tgcactccag cctggggccac 840  
agagtgaaac ctctgtcaa aaacaaaaaa acaaaaaaac acagtgtgtt agatcttgct 900  
agacttggtg atataattaa gaggccatta tgggcagaac tgtgccccct tccaaaattc 960  
atatataaat atatataaat atatataaat atataaatat atataaatat ataaatatat 1020  
ataaatatat aaatatatat aaatatataa atatataata atatataata atatataaaa 1080  
atatataaat atatataaat atatataaat atataaaaa ataaaaaatat atataaatat 1140  
atataaatat ataaaaaat ataaatatat aaatatataa aaatatataa atatataaat 1200  
atatacataa atatataata atatataata atatataaaa atatataata atatataaat 1260

## SEL PCT 012.ST25

atatataaat atatataaat atatataaat atataaaaat atatataaat atataaatat 1320  
 ataaaaatat atataaatat ataaatatat aaatatatat aaatatataa atatataaat 1380  
 aaatataagt atttatgaat atatatgaat atataaatat ataaaaaata tatataaata 1440  
 tataaatata tataaatata taaatatata catatataca tatataaata aataaatata 1500  
 agtatftatg aatatatatg aatatataaa tatataaaaa atatatataa atatataaat 1560  
 atatataaat ataaatatat aaaaatatat aaaaatatat ataaatatat aaatatatat 1620  
 aaatatataa atatatataa atatatataa atatataaat atatataaat atatataaat 1680  
 atataaatat ataaatatat ataaatatat ataaatatat aaatatataa atataaatat 1740  
 ataaatatat ataaatatat ataaatatat aaatatatat aaatatatat aaatatatat 1800  
 aaatatatat aaatatataa atatatataa atatatatat aaatatatat aaatatataa 1860  
 atatataaat atataaaaat atataacaat atataaatat atataaaaat atataacaat 1920  
 atataaatat aaatatatat aaaaatatat aacaatatat aaatataaat atatataaat 1980  
 atataaatat aaatatataa aatatatata aatatataaa tatatatataa tatataaatg 2040  
 tataaatata tataaaaaata tataacaata tataaatata taaatatata acaatatata 2100  
 aatatataaa aatatataac aatatataaa tataaatata tataaaaaata tataacaata 2160  
 tataaatata aatatatata taaatatata aatatataaata taaaaaatat atataaatat 2220  
 ataaatatat atataaatat atataaatat ataaatgtat aaatatatat aaatatataa 2280  
 atatatataa atatataaat atatataaat atatataaat atataaatat aaatatataa 2340  
 atatatataa atatataaat ataaatatat aaacatatat aaatatatat aaataaacat 2400  
 atataaagat atataaagat ataaagatat ataaatatat aaatatataa agatatataa 2460  
 atatataaag atatataaat atataaagat atataaatat ataaagatat ataaatatataa 2520  
 tatataaata tataaagata tataaatata atataaaaat atataaatat atattaaaaa 2580  
 tatatacata taaatatatg tatatftttt tgagatgggg tctcgctcag ccaccacgc 2640  
 tggagtgcag tggcagcagc tcggctcact gcaaccactg tctctcgggt ccaagcaatt 2700  
 ctgtctcagc ctccaagta gctgggatta caggcacctg ccatcatgcc cggctaattt 2760

## SEL PCT 012.ST25

ttgtatttta gtagagatgg agtttcacca tgttggccag gttggtctcg aattcctgac 2820  
ctcaggtgat ctgccggcct cggcctccca gtgctgggat tacaggcatg agtcaccacg 2880  
cccgcccta tatatatttt tgagacaagc tctgtgtctc ccaggctgga gtgcagcagc 2940  
atgatcatga ctactgtag cctagacctc cagggctcaa gtgattctcc cacctcagcc 3000  
tcccaagtag ctgggattac aggcattgcac caccacaccc agctaatttt tgttttgttt 3060  
tgttttgag acagaatctc tctctgtcac ccaggctgga gtgcagtggg gtgatctcag 3120  
ctcagtgcaa cctccacctc ctgggttcaa gtgattctca tgcctcagcc tctgagtag 3180  
ctgggactac aggcgtgagc caccacgccc tgataaattt tgtatttttt ttctcagatg 3240  
gagtctcact ctgtcatact caggctggag tgcagtggcg tgattttggg ttattgcaac 3300  
ctctgcttcc tgggttcaag cgattctcct gcctcagcct ccagagtagc tgggattaca 3360  
ggcgctgcc accatgccca cctagctaac ttttttttt tttttttga gatagagtct 3420  
cactctgtca ccaggctgg agtgcaatgg ggcgatattg gctcactaca acctccacct 3480  
cccagggtca agcgattctc ctgcctcagc ctctgagta gctgggatta cagggtgggtg 3540  
ccaccacgcc agactaatat ttgtattatt agtagagacg gggtttcacc acattgggtca 3600  
ggctggtttc gaactcctgg cctcagggtga tctgcctgcc tcggcctccc aaagtgctgg 3660  
gattacaggc atgagccact gcggctggcc caatttttgc attttttgg tagagacggg 3720  
ggtttcacta tgcttccag gctggtctca aactcctgga ctcaagcgat ctgcctgtct 3780  
cagcctccca aagtgcaggg attacagtca tgagccacca ctgcacggcc ccaaaattta 3840  
ttattttat tattattatt attttttga tggagtctcc ttctgttgc agattggaat 3900  
gcagtgccac gatctcagct cactgcaacc tccccctct gggatcaagt gattcttttt 3960  
ttttaagac tctgtctcaa aaaaaaagaa aaaaaaaaaa aatatatata tatatatata 4020  
tatacacgaa ttttgggcca ggcacagtgg ctatgcctg taatcccagc actttgggag 4080  
ggccgaggtg ggtggatcac aaggtcagga gtttgagacc agcctggcca atatggtgaa 4140  
acctgtctc tactaaaaaa tacaaaaatt agctgggcgt ggtggcacga gcctgtaatc 4200  
ctggctactc tggaggctaa ggcaggagaa tcgcttgaac cggggaggca gaggttgcag 4260

## SEL PCT 012.ST25

tgagccagga tcgcatcact gcactccagc ctgggtaaca gagcaagact ctgtctcaaa 4320  
 aaacaaacaa aacaaaacaa aacaaaataa ataacggtgc aaaattgaat atgccttttt 4380  
 gactctctaa atgcctcaga tccatttacc ctggggattt gtcctttcta gccccaccac 4440  
 catctcccct ctggaagact gctgacctat aaggataaag accagactct tgagcaggca 4500  
 cttaggggtct tctgcccacat ccctatcccc aactccccct cagtaatttt ggctactagt 4560  
 atttctccac atctgaggct atcgtgggtc tcccttcagt ggcatgaag gacaagggtg 4620  
 gagaagtttg ccctcgtgag tctgatgagg gattgggtgg 4660

&lt;210&gt; 27

&lt;211&gt; 3354

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(3354)

&lt;223&gt; MAR X\_S29 of chromosome X

&lt;400&gt; 27

gatcccttta taaaaccaca atataatgga gtgctataat ttcaaacagt gtttggctcg 60  
 ctggcagagt ggtcattcta acagcagtc cagtagagta gaaataagac tgcagtatat 120  
 ctaaggcaaa aagctgaggt ttcaggagct tgaaggtaaa gaggaagaaa gaaatgggaa 180  
 tgggaattgg aaagacaaat atcgtaaga gaaaattgct ttaggagag gggaaagaat 240  
 ctatgtgtac ttaagactat ggaatcaatc ccatttaagc tgggaaacta gtttcatata 300  
 taactaataa atttatttta cagaatatct atttacctga tctaggcttc aagccaaagg 360  
 gactgtgtga aaaacatca gttctgtcat attcctaaaa aaaaattaaa aagttaaaaa 420

## SEL PCT 012.ST25

taaataaata ataaaacttc ttttctttca aaataatcaa ggtgcttatt cacatccatt 480  
 ccaatttggg gaaatactta ttttctatg attagcgaag agaaaagtaa ctgcatctc 540  
 aattcaagtt gatacatgtc actttaaga ggtcaactaa tatttgctag ttgagctaac 600  
 catataggct ttaaatactt tcatagtaga aagaaaatga aaatcattag tgaactgtat 660  
 aaaatagatc atactttttg aaagaatcag actgaagttt ccgaaaaaaa gaagtaagct 720  
 tcaatgaaaa ggtaagtga ttagcatctt actcagcatc tactatggac ttaacaccta 780  
 acagtagata atctgaaggc aaacatattt gtataggggac tgcagaatga tagatgataa 840  
 atatcatctc ttctatttga atgaatattt ttcaaactc ttacacaca gtggtttgct 900  
 atggaaagat tttagtaga ttaaacaat ctgaagatgg agttagaaag cttaggctat 960  
 gttttgagca caacatataa tttctctgtg attgtttctt catctttcaa atgaggttac 1020  
 tgtgaagatt aaatgagata actaaatgat gataaaataa tgtaacttta gcagcacctt 1080  
 atttaactctg tgcaacaact ctgtgaagt agtagggctc agcttcagtc acttctctgc 1140  
 catttattaa ctaagatagt ttggaaagtt acctatctct tcagctgtaa aatgatgagg 1200  
 atcataccta tttatgggg ctgcttttag gtacaaatat acaggcaagc actttgttaa 1260  
 tactaaagca ttacaccaat tagttttact cttttccatt cacacatgaa attaatgtaa 1320  
 tcagaattct gtagattacc taaatcttct gtaacacgt gatatgcagt tcagggttaa 1380  
 tgtcagttga gttaccaaag cacatacata ctaccacccc tatccaaatc tacaagcctc 1440  
 ccagttgtc ttactattt tggttaaatt aatatgaatt cctagatgaa aatttcactg 1500  
 atccaaatga aataaaaaat atattacaaa actcacacct gtaatctcaa cattttggga 1560  
 ggccaaggca ggtagatcac ttgaggccag gagttcaaga ccagcctgat caacatggtg 1620  
 aaaccctgtc tctactaaaa atacaaaaat tagccagggtg tgggtggcatg tgctgtagt 1680  
 cctacctact cgggagggtg aggcacaaga atcgcttgaa tgtgggaggt ggagggtgca 1740  
 gtgacctgag atcggtccac tgcactccag cctaggcaac agagtgagat catgtgtcat 1800  
 atatatatat atatatatat atatatatat acacacacac acatatatat 1860  
 atacacatat atatacgtat atatatatat gtatatatat acatatatat acatatatat 1920

## SEL PCT 012.ST25

atatacgtat atatatacgt atatatatat caatgtaa atttgggaa atttggtatg 1980  
 aatagtcttc cctgtgaaca cagatcataa aatcatatat caagcagaca aataagtagt 2040  
 agtcacttat atgcttatac ttgtaactta aagtaaaaga attacaaaag catatgacaa 2100  
 agactaattt taagatatcc taatttaa atgtttctaa aagtgtgtat accattttac 2160  
 ctatcatatg aataatttag aaacatgttt ataaaattaa tgcctaaatc cattcaaaag 2220  
 ttttgtaatg cagatcacc acaacaaca agaatcctag cctattaaaa aagcaacacc 2280  
 acctacatat aatgaaatat tagcagcatc tatgtaacca aagttacaca gtgaatttgg 2340  
 gccatccaac actttgagca aagtgttgaa ttcatcaaat gaatgtgtaa tcatttactt 2400  
 actaatgcc aatacactta aggtaatctt aagtagaaga gatagagttt agaatttttt 2460  
 aaatttatct ctgtttgtaa agcaatagac ttgaataaat aaattagaag aatcagtcatt 2520  
 tcaagccacc agagtatttg atcgagattt cacaaactct aactttctga taccatttct 2580  
 cccaaaaacg tgtaacctcc tgcgatagg aacaaccac tgcagggatg ttctctgtgg 2640  
 aaaaaggaaa ttcttttgc attggtttca gacctactg gttacaagaa aaaccaaagg 2700  
 ccattgcaca atgctgaagt actttttca aatttaaaat ttgaaagttg ttcttaaaat 2760  
 ctatcattta tttaaaata cggatgaatg agaaagcata gatttgataa agtgaattct 2820  
 ttctgcaat ctacagacac ttcaaaaat cactacagac actacagaca ctacagaaaa 2880  
 tcataaataa acaagtgcata gtatcaatat tttaacaaa aaatggcatt cttagaattt 2940  
 ttataggct agaaggtttg tacaactaa tctgccacgg attttaa atgagtgaat 3000  
 aaatttatatt gcaaaaaaaaa tcaggttaca gagaactggc aaggaagact cttatgtaaa 3060  
 acacagaaaa catacaaac gtattttta gacaataaa aacagaactt gtacctcaga 3120  
 tgatactgga gattgtgttg acatattagc attatcactg tcttgctaaa acataaaaat 3180  
 aaaaagatgg aagatgaaat tacaatacaa atgatgattt aaacatataa aaggaaaata 3240  
 aaaattgttc tgaccaacta ctaaaggaag acctactaaa gatatgcat ccagcacatt 3300  
 gccactctac atgtgtgtc taaaccagca gcatagggat cctctagcta gagt 3354

&lt;210&gt; 28

## SEL PCT 012.ST25

&lt;211&gt; 677

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(677)

&lt;223&gt; MAR of chromosome 1 genomic contig; 12803267..12803943

&lt;400&gt; 28

```

ttatatagta tatataatag tatatattat atagtataca taatagtata tattatatag   60
tatacataat agtatatatt ataatataca taattgtata tatcatatag tatacattat   120
agtatatatc atatagtata cattatagta tatatcatat agtatacatt atatagtata   180
tatcatatag tatacattat agtatatc atatagtata cattatagta tatatcatat   240
agtatacgtat atagtatata tcatatagta tacgtaatag tatatatcat atagtatacg   300
taatagtata tatcatatag tatacgtaat agtatatc atatagtata cgtaatagta   360
tatatcatat agtatacgtat atagtatata tcatatagta tacgtaatag tatatatcat   420
atagtatacg taatagtata tatcatatag tatacgtaat agtatatc atatagtata   480
tattatatag tatatatcat atagtatata ttatatagta tatatcatat agtatatatt   540
atatagtata tatcatatag tatatattat atagtatata tcatatagta tatataatag   600
tatatatcat atagtatata taatagtata tatcatatag tatatatact atactatatt   660
atatatagta tacataa

```

677

&lt;210&gt; 29

&lt;211&gt; 332

&lt;212&gt; DNA

## SEL PCT 012.ST25

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(332)

<223> MAR of chromosome 1 genomic contig; 13079684..13080015

<400> 29

```

ttaattatat tatatatatt atataattat atattaatat atattaatta tattatatat   60
attatataat tatatattaa tatatattaa ttatattata tatattatat aattatatat   120
taatatatat taattatatt atatatatta tataattata tattaatata tattaattat   180
attatatata ttatatatta taattatata ttatataatt ataatatata tgттаатата   240
atatatataa ttaatatata attaaaacta tttaattata tgtatattat atataaatatg   300
tattatttaa ataataaata tattatttat at                                     332

```

<210> 30

<211> 479

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(479)

<223> MAr of chromosome 1 genomic contig; 15682296..15682774

<400> 30

## SEL PCT 012.ST25

acaagtacat atatatatag tatatatata caagtacata tatatagtat atatatatat 60  
acaagtacat atatatatagta tatatatata tacaagtaca tatatatagt atatatatat 120  
acaagtacat atatatatagta tatatatata caagtacata tatatatagt atatatatat 180  
acaagtacat atatatatagta tatatatata caagtacata tatatatagt atatatatat 240  
acaagtacat atatatatagta tatatatata caagtacata tatatatagt atatatatat 300  
acaagtacat atatatatag tatatatata tacaagtaca tatatatata gtatatatat 360  
atacaagtac atatatatag tatatatata tatatacaag tacatatata tagtgtatat 420  
atatatatat aagtacatat atatacttgt attagtatat atatatatat atacaagta 479

<210> 31

<211> 531

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(531)

<223> MAr of chromosome 1 genomic contig; 15694611..15695141

<400> 31

tataatatat ataatacata atagatatat tatattatat aatagatata taattataaa 60  
cataataata tataatgaat ataataataa ataaatataa taaaatatat aatatatcta 120  
ttatgtatta tatattatat atgtttatat ataataaat tatatatggt tatatataat 180  
ataattatat atgtttatat ataataaat tatattatat atattataga tataatatat 240  
aatatactat atattataga tataatatat aatatactat atattataga tataatatat 300  
aatatactat atattataga tataatatat aatatactat atattataga tataatatat 360

## SEL PCT 012.ST25

aatatactat atattataga tataatatat aatatatatt atatattata gatataatat 420

ataatatatt atatattata tctatatata atatattgta tattatatat aatatattgt 480

atattatata taatatattg tatattatat ataatatatt gtatattata t 531

<210> 32

<211> 378

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(378)

<223> MAR of chromosome 1 genomic contig; 886276..886653

<400> 32

ttatattata tatcttacct aaattatata tatatattac ataaattata tacaatataa 60

attatataca atataattta tatataaaat ataaattata taaataattt atatataaaa 120

tataaattat ataaataatt tatatataaa atataaatta tgtataaaat ttatatataa 180

aatataaatt gtgtataaaa ttatatataa aatataaatt gtgtataaaa ttatatata 240

aaatataaat tatatataat ttatatatta taatataaat tatatataat atatatcata 300

aaatataaat tatatataat atatatcata agatataaat tatatataat atatatcata 360

agatataaaa tatataat 378

<210> 33

<211> 595

<212> DNA

<213> Homo sapiens

## SEL PCT 012.ST25

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(595)

&lt;223&gt; MAR of chromosome 1 genomic contig; 3326732..3327326

&lt;400&gt; 33

```

aaaatatata aatatatata aaaatatata aaaatatata aatatatata aaaatatata   60
aatatatata aatatatata aaaatatata aatatatata aatatatata aaaatatata   120
atatatataa aatatatata aatatatata aatatatata aaaatatataa tatatatataa   180
aatataaata tatataaata tatataaaaa tataaatata tataaatata tataaatata   240
taaatatata taaatatata taaatatata aatatatata aatatatata aatatatata   300
aatatataaa tatataaaaa tatatatataa tatataaata tatataaata tataaatata   360
taaaaatata tataaatata taaatatata taaatatata taaatatata tataaatata   420
tataaatata tatatatata aatatatata aatatatata taaatatata taaatatata   480
tatatatata taaatatata taaatatata taaatatata tataaatata tataaatata   540
tataaatata tatataaata tatataaata tatatatataa tatatatataa tatat      595

```

&lt;210&gt; 34

&lt;211&gt; 738

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(738)

## SEL PCT 012.ST25

<223> MAR of chromosome 1 genomic contig; 4485716..4486453

<400> 34

ataatagata atatatatta tatgatagat atataatata ttatataata tataatatat 60  
 tatatatcta tcatataata tatataatat ataatatatt atatatctat catataatat 120  
 aatatatata atataataa tatatcatat tatattgtat ataatatata tcatattata 180  
 ttgtatataa tatatatcat attatatatt atataatata tatcatatta tattgtatat 240  
 aatatatatt atattatatt gtatataata tatatcatat tatattgtat ataatatata 300  
 tcatattata ttgtatataa tatatatcat attatatatt atataatata tatcatatta 360  
 tattgtatat aatatatatt atattatatt gtatataata tatatcatat tatattgtat 420  
 ataatatata tcatattata ttgtatataa tatatatcat attatatatt atataatata 480  
 tatcatatta tattgtatat aatatatatt atattatatt tattatattg tatataatat 540  
 atattatata ttatctatta tattgtatat aatatatatt atattatatt tattatattg 600  
 tatataatat atattatata ttatctatta tattgtatat aatatataat aaatatagta 660  
 tatataatag ataatatata gtatatatga tatattatat atactatata ttatatatca 720  
 tatatactat atactata 738

<210> 35

<211> 386

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(386)

<223> MAR of chromosome 1 genomic contig; 5423067..5423452

## SEL PCT 012.ST25

&lt;400&gt; 35

taa~~a~~atata a~~a~~aatata t~~a~~aaaatata a~~a~~aatat~~t~~ta tataaatata t~~a~~aaaatatt 60

tatataaata tataaatata taaatataa tttatataaa tatataaata tataaatata 120

taa~~a~~atata tttatataaa tatataaata tatatttata taaatataa aatataata 180aa~~a~~atataa atatatattt atataaatat at~~a~~aatatat at~~a~~aaaatata taaatataa 240tattttatat aaatata~~t~~aa atatatataa aatata~~t~~aaa tatatatatt ttatataaat 300atataaatat atata~~a~~aaata tataaatata tatattttat atattttat atataaatac 360atata~~t~~ttca tatatcacat atatga 386

&lt;210&gt; 36

&lt;211&gt; 584

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(584)

&lt;223&gt; MAR of chromosome 1 genomic contig; 5805559..5806142

&lt;400&gt; 36

taa~~a~~atattt t~~a~~aaaatata atattttata atata~~a~~att tatattataa tgtgtacata 60

atatatatta taatataata tatataatac tgtatattat attatatata ttataatata 120

tattattata tattatatta tatataaat aatatatatt ataatatatt atattataca 180

tattataatg tattataata tatattatat tatatattat aatatatatt atattatata 240

ttataatata tattatatta tatattataa tatatattat attatattat atatattata 300

## SEL PCT 012.ST25

atacatatta taatacatat tatataatat attataatat gtattataat acatattata 360  
 taatatatta taatatatta tatataataa tatattataa tacatattat atataatata 420  
 tattatgtat attatatata atatatatta caatgtatat tatgtatatt atatatatta 480  
 tatatcatat aatatatatt atatatataa tgatatataa tatatattat ataatatatt 540  
 atatgatata tataatatgt attacatgta atatatatca taat 584

<210> 37

<211> 345

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(345)

<223> MAR of chromosome 1 genomic contig; 10802644..10802988

<400> 37

tgtatatata tactatatat atactatata tatagtgtat atatatacta tatatatact 60  
 atatatatag tgtatatata tactatatat atagtatata gtatatatag taatatatat 120  
 atatagtata tatatacact atatatagta tatatagtat atatatattg tgtatatagt 180  
 atatatatag tgtatatata gtatatatat attgtatata tagtatatat attgtgtata 240  
 tatagtatat atatagtata tatagtatat atagtatata tatagtatat atatactata 300  
 tatatagtat atatatattg tatatatata ctatatatat agtat 345

<210> 38

<211> 474

<212> DNA

## SEL PCT 012.ST25

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(474)

<223> MAR of chromosome 1 genomic contig; 13496468..13496941

<400> 38

atattatata taatataatt atatctataa ttatatatta tatataatat aattatatat 60

ctataattat atattatata taatatatat tatatataat atataattat atataattta 120

tataatataa tatataatat ataattatat ataattatat aatataatat ataatatata 180

attatataata atttatataa tataatatat aatatataat tatatatatt tatataatat 240

aattatatat aatatataat tatatataat ttatatataa taattatata taatatataa 300

ttatatataa ttatatataat ataattatat ataattatat atttatata atttatataa 360

tataattata tataatatatat aattatatat aatatataat tatatataat tatatataat 420

atataattat atataattta tataatataa ttatatatta tatatatatt atat 474

<210> 39

<211> 483

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(483)

<223> MAR of chromosome 1 genomic contig; 2509163..2509645

## SEL PCT 012.ST25

<400> 39  
 caaaatacat aatatataat agtattatat aatagtatgt atagttataa tatatagtat 60  
 aattacaata tatgatatgg ttatatatt atatatagta taatataata taacataata 120  
 ctattataat atataaacta tataatatat actattataa tatatgaact attataatat 180  
 ataaactata tataatatat aatatgtact attataatat ataaactatt ataataatat 240  
 atataaacta ttataataca taaactatta taatatatat aatactatgt atacatatat 300  
 tacattatgt acatactaca ttatgtatta tgtatgtata tatacacaaa atacataata 360  
 tataatagta ttatataata gtatatatag ttataatata tagtataatt acaatatata 420  
 atatgggtta tatattatat atagtataat acaatataac ataatactat tatatataaa 480  
 cta 483

<210> 40

<211> 641

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(641)

<223> MAR of chromosome 1 genomic contig; 2776349..2776989

<400> 40  
 tgttatatat atataacata gatattatat atacatgtta tatatataac atagatatta 60  
 tatatacatg ttatatatat aacatagata ttatatatat aacatagata ttatatatac 120  
 atgttatata taacatagat attatatata catgttatat ataacagata ttatatatac 180

## SEL PCT 012.ST25

atgttatata taacatagat attatatatg tatgttatat ataacataga tattatata 240  
 gtttatataa tatataacat atgtttaaca tatataatat ataacatgtt tatataatat 300  
 ataacataat tatagttat atatgatata aaacatatat attatatacg ttatatgtaa 360  
 tatataacat atattgtata cgttatatgt aatatataac atatattgta tacgttatat 420  
 gtaatatata acatatattg tatacgttat atgtaatatata taacatatat tgcatacgtt 480  
 atatgtaata tataacatat attgtatacgt ttatatgtaa tatgtaacat atattgtata 540  
 cgttatatgt aatatgtaat atataatata tataacatgt atatataaca tatatgtata 600  
 taacatatat ataacatatata taacatatat gttatattat a 641

<210> 41

<211> 745

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(745)

<223> MAR of chromosome 1 genomic contig; 2858703..2859447

<400> 41

atatttatat atgtaataat atataatata ttatatatgta ttgtatatg taataatata 60  
 tatataataa aatatgtaat aatatataat atatttatat ataaatatat tatatttatat 120  
 atatattatt atatttataa tataatatat atttatatta tatattataa atatattatta 180  
 tataatatat attataaata tatatttatat aatatatatt ataaatatat attatattat 240  
 atattataaa tatatattat ataatatata ttataaatat atatttatata atatattatta 300  
 taaatatata ttatattat aatatatatt ttgtatatatt atatattata tattataaat 360

## SEL PCT 012.ST25

attattatat ttataatata ttatatattt tatatataat atatgatata tattataaat 420  
 atatcttata aatatatata ttatatata tatattataa atatataaat ataaatatat 480  
 aatataatat aatataatat aataaatata atatataata tatataatat ataataaata 540  
 taataaatat aaatatatca tataaatata aatataaata taaatatatc atataaatat 600  
 atatatttat atgatataatt atagtatata taaatatatt tatatattat aaaatattta 660  
 tataatatat aattataata ttttatata tataaattaa ctaatatata taaactaata 720  
 taatatataa tgtaataata tagta 745

<210> 42

<211> 307

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(307)

<223> MAR of chromosome 1 genomic contig; 945522..945828

<400> 42

catatataat atatattacc tatgttatat aggtcatata taacataaat atattacata 60  
 tatgtaatat atattaaata taaatatata acatatatgt gtaactatat atgtaaatat 120  
 gtacatatac atatatgtaa atatataata tatatttaca ttatattata taatatatat 180  
 ttacattata ttttatata tacattatat atatttacet tataaatatt tatataatat 240  
 atatttacet tatattacat tatataaaat acaatatatt acattataat acattataac 300  
 agataaaa 307

<210> 43

## SEL PCT 012.ST25

<211> 357

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(357)

<223> MAR of chromosome 1 genomic contig; 3402743..3403099

<400> 43

aatattatat taaatataat atattaatat ttaatatatt taatataata ttaaataaat 60

atattataaa taaattataa tatataaata tatattatgt atttatgtat aatatataaa 120

aattatatat aatatatata tttttataaa tatataaata tataataaat aaatatatta 180

aataaataat aatatattaa atattaatat attaaatatt atatattaaa tataatatgt 240

aatatgaaat atattaaata ttatatatta aatataatat ataatttgaa atatattaaa 300

tattatatat taaatataat atataatatg aaatatatta aatattatat attaaat 357

<210> 44

<211> 323

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(323)

<223> MAR of chromosome 1 genomic contig; 3485830..3486152

## SEL PCT 012.ST25

<400> 44  
 atatttatag actatatatt tatatatatta gtgtatttgt atactatata tttatatagt 60  
 tagtatattt gtatactata tatttatata tttagtatat ttgtatacta tatatttata 120  
 tatttgaat atttgatac tatatatatta tatatttagt atatttgtat actatatatt 180  
 tagtatattt gtatactata tatttatata tttagtatat ttgtatacta tatatttata 240  
 tatttagtat atttatatac tatatactta tatatttagt atatttatat actatatact 300  
 tatatatatta gtatatttat ata 323

<210> 45

<211> 498

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(498)

<223> MAR of chromosome 1 genomic contig; 3548336..3548833

<400> 45  
 aattattact atattgttaa tataattatt atataatata atataattat atcactatta 60  
 ttatattata gtattaatat aatagtgtat aacattaata taatatagta ttaatataat 120  
 agcgtataac attaataata tatagtatta atataatagc gtataacatt aatataatat 180  
 agtattaata taatagtgtat tattaatata atatagtatt aatatataat attaataata 240  
 tatatcaata taatagtata taatataata taatatatca atataatagt atataatata 300  
 atataatata tcaatataat agtatataat ataataatata atatcaatat aatagtatat 360

## SEL PCT 012.ST25

aatattaata taatataata tcaatataat agtatataat attaatatat taatataata 420  
 gtatataata ttaatgtaat ataatatata cataatgtat ataataatat aatagtatat 480  
 aataactaata taatataa 498

&lt;210&gt; 46

&lt;211&gt; 400

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(400)

&lt;223&gt; MAr of chromosome 1 genomic contig; 4595109..4595508

&lt;400&gt; 46

aaatatatta tattatatat tatatatatt tcaatataact ataatatata ttatatatgt 60  
 ttaatacaat atataatatt tacatatatt cccatttatt tatataacat atattatatg 120  
 atattatata ttactccata taatataata tattatacat aatatattac tcagtataat 180  
 acataatata tataatatat tactcggtat aatatataat attatatgtt atgcaatata 240  
 atatatataa ttatatataa tacattattc aatataatat ataatatatt atataataca 300  
 ttattcaata taatatataa tacactattc aatataatat acaatatatt atataataca 360  
 ttattcaata taatatatat tatataatat atatatttat 400

&lt;210&gt; 47

&lt;211&gt; 403

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

## SEL PCT 012.ST25

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(403)

&lt;223&gt; MAr of chromosome 1 genomic contig; 7205509..7205911

&lt;400&gt; 47

agtatatata tgtgtatata tatgagtata tatatgtgta tatatatgag tatatatatg 60

tgtatatata tgagtatata tatgtgtata tatatgagta tatatatgtg tatatatatg 120

agtatatata tgtgtatata tatgagtata tatatgtgta tatatatgag tatatatatg 180

tgtatatata tgagtatata tatgtgtata tatgagtata tatatgtgta tatatgagta 240

tatatatatg tgtatatatg tgagtatata tatgtgtata tatatgagta tatatgtgta 300

tatatatgag tatacatatg tgtatatata tgagcatata tgtgtatata tatgagtata 360

tatatgtgta tatatatgag tatatatgtg tatatatatg agt 403

&lt;210&gt; 48

&lt;211&gt; 309

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(309)

&lt;223&gt; MAR of chromosome 1 genomic contig; 7507280..7507588

&lt;400&gt; 48

## SEL PCT 012.ST25

tataaaatat atattattta tatattatat ataaaatata tattatatta tatattatag 60  
 atataataaa taaataatat ataatatatt atataattat ttatacataa ttatatataa 120  
 ttatatgtaa ttgtacaatt atatataatt atatacaatt atacacataa ttatatacaa 180  
 ttatacaatt atatacataa ttatatatat aatatacata attatatatt aattatacaa 240  
 ttatatatat aattatatat aattatacaa ttatatatat aattattatg tatattatat 300  
 tatataata 309

<210> 49

<211> 516

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(516)

<223> MAR of chromosome 1 genomic contig; 3581085..3581600

<400> 49

atatatatat atatatatat atttatatat atatatatatta atatatatatta tatataaaaa 60  
 tatataaaaat ttatatatat aatttatata tataaaaaata tataaaaattt atatatatata 120  
 tttatatata taaaaatata taaaatttat atatataatt tatatatata aaaatatata 180  
 aaatttatat atataattta tatatatataa aatatataaaa atttatatat ataatttata 240  
 tatataaaaa tatataaaaat ttatatatat aatttatata tataaaaaata tataaaaattt 300  
 atatatatata tttatatata taaaatatat aaatttatata tataattata tatataatat 360  
 aaaatttatat atataattat atatataata taaaattata tatataatta tatatatatat 420  
 ataaaattat atatatatig tatatatata aaatatataa aatttatata tataaaaatat 480

## SEL PCT 012.ST25

aaaatataca taaaaataaa tatatataat ttatat 516

<210> 50

<211> 534

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(534)

<223> MAR of chromosome 1 genomic contig; 3084851..3085384

<400> 50

atataatata tatgactata tttttatat tatattctat ttcaataaaa tatttatatt 60  
 ttattatata ttataatata taattatata tgtaataata tataatataat aatatatatt 120  
 ttatattata ttttatattt atttttatat ttatattat attttattat atatattata 180  
 atatataatt atatatgcaa taatatatta tatattataa tatataatta tatatgcaat 240  
 aatatattat atattataat atataattat atatgcaata atatattata gattataata 300  
 tataattata tatgcaataa tatattatat attatatatt agataatata ttaatatata 360  
 ttataacata taatatataa catataatat ataatatatt atctaataata taatataaca 420  
 tataatataat aatatattat ataatatatt attacatata taatatattg taatatataa 480  
 tattacatat atottcaaaa agagttatgt gtatataata catatatata ccat 534

<210> 51

<211> 583

<212> DNA

<213> Homo sapiens

## SEL PCT 012.ST25

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(583)

&lt;223&gt; MAR of chromosome 1 genomic contig; 160087..160669

&lt;400&gt; 51

ttttatata aaatatataa aatatattat atataaatat attatatata atatatttat 60  
 atattataca atatatttat atattatata taatatattt tatataatat acataatata 120  
 ttttatatat tatatataat atattttata tataatgtac aatatatttt atatattata 180  
 tataatatat tttatatata ctatacaata tttttatat attatatatt ttatatatat 240  
 ttttcagtga acatatatat tttatatata atatatatac catatataat atattttata 300  
 tataatatat ataccatata taatatattt tatatataat atgtatatca tatatagtat 360  
 attttatata taataggtat accatatata atatatttta tatataatag gtataacata 420  
 tataatatat tttatatata atatgtatac catatataat atattttata tattatagat 480  
 accatatgta atatacttta tatataatat agataccata tgtaatatat tttatatata 540  
 atatagatac catatgtaat atactttata tataatatag ata 583

&lt;210&gt; 52

&lt;211&gt; 314

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(314)

## SEL PCT 012.ST25

<223> MAR of chromosome 1 genomic contig; 4350424..4350737

<400> 52

```
tatgtgtata taaatatatg tatatatgtg tatataaata tatataaata tatgtatata    60
tgtatatata catatatatta tatataaata tatgcatata tttatatata aaatatatgc   120
atatatgtat atatataaaa tatatacata tatgtatata tataaaatat atacatatat   180
gtatatatat aaaatatata catatatgta tatatataaa atatatacat atatgtatat   240
atataaaata tatacatata tgtatatata taaaatatat acatatattt atatatataa   300
aataccaagt ctta                                     314
```

<210> 53

<211> 828

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(828)

<223> MAR of chromosome 1 genomic contig; 8443267..8444094

<400> 53

```
tattatataa ttatatatac tatataatta tataatatat agttatatag tatatataat    60
atatataata tatactatag tatatataat atatataata tatactatag tatatataat   120
atataattat atataatata tataatatag tatatattat atatatatta tatatatata   180
atatatatat aatatatata atatagtata tataatatat aattatatat aatatataat   240
atagtatata taatatataa tatatatata attatatact ataatatata taatatataa   300
```

## SEL PCT 012.ST25

ttatatatta tatactatag tatatattat tatatataat agatataata tataataatta 360  
 ttatataata tagtatatat aatatataat tatatataat agatataata taatataatt 420  
 atatataata tagtatatat aatatataat tatattatat tatatataat atataattat 480  
 aatatataat tatattatat aatatatata atatataatt atattatata attatattat 540  
 ataatatata taatatataa ttatattata taatatatat aatatataat tatattatat 600  
 aatatatata atatataatt atattatata atatataata tatataatta tattatataa 660  
 tatatataat atataattat atattatata taatatagta tatataatat gtaattatat 720  
 atcatataat atataacatt gtatataata tataattaca tattatataa tgtatataat 780  
 atataattat atacattata taatatagta tataattata tattatgt 828

&lt;210&gt; 54

&lt;211&gt; 573

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(573)

&lt;223&gt; MAR of chromosome 1 genomic contig; 8703190..8703762

&lt;400&gt; 54

tatattatat ataaaatata catataatat acctataata tacatataat atataatata 60  
 tattatgtac atataatata catataatat atataatata taatgtacat ataataatata 120  
 tataatatat gttatatatt atatataaaa tataggatat atataatata gaatatatat 180  
 actatattgt atatataaga tatataatat atagtatata tactatataa tatataatat 240  
 atagtatata taatatataa tatagaatat atatacaata tataatatag aatataggat 300

## SEL PCT 012.ST25

atatatagaa tatacatata taatatgtat atattatata ttatattata tattatataa 360  
 aaatatataa tatataatat aaaaatatat tatatattat ataataaaa atatattata 420  
 tattatatat tatataatat aaaatatatt atatattata tattatatat aaaaatatatt 480  
 atatattata tattatatat aaaaatatat tatatattat atattatata taaaaatata 540  
 ttatatatta tatataaaaa tatatattat tac 573

<210> 55

<211> 597

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(597)

<223> MAR of chromosome 1 genomic contig; 8819076..8819672

<400> 55

acatatctta tatataaaat atataaatat acacatattt tatatataat atatattata 60  
 tatatgaaat atacacatat ttttatatat ataatatata tattatatat aatatatgca 120  
 tatattatat ataaaatata tatattatat ataaaatatg catatattat atataatata 180  
 tataatataa aatatataat atatattata tattatatat aatatatatt atatataata 240  
 catatatata atatataata tatataaaat ataatatata tattatataa tatatatata 300  
 aatatatata atatatatat aatatatata ttatatataa aatatatatt atatgtaaaa 360  
 tatataatat atataatata tatattatat gtaaaaatata tattatatat aaaaatatata 420  
 atatatataaa tatatattat atataaaata tataatatat aaaatatata atatatatata 480  
 aatatataat atatataaat atatattata tataaaatat ataatatata taaatatata 540

## SEL PCT 012.ST25

ttatatataa aatatataat atatataaat atatattata tataaaatat atattat 597

<210> 56

<211> 646

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(646)

<223> MAr of chromosome 1 genomic contig; 759619..760264

<400> 56

taatatatat aatatatatt atataataat atataatata tattatatta taatatataa 60  
 tatattatat aataatatat attatataat atataataat atatataata catattattt 120  
 aataatatat aatatatatt atataataat atataatata tattatataa taatatatat 180  
 tatattatat aatatataat atatataata tatattatat aataatatat aatatatatt 240  
 atagaatgat atattagata ttatataatt atatatataa tattatatat tatataataa 300  
 tatataatat atattatata attatataa taatattata tattatataa ttatatataa 360  
 tatattatat aattatatat ataattatt atattatata attatataa atatatatta 420  
 tataattata tatataatac tatatattat ataattatat ataatactat atattatata 480  
 atttatataa ttatatatat tatatattat ataattatat atattatata ttatataata 540  
 acatatatat tatatattat ataataacat atatattata tattatataa tacatatata 600  
 ttatatatta tataatacat tattatataa tatataatat atatta 646

<210> 57

<211> 752

## SEL PCT 012.ST25

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(752)

&lt;223&gt; MAR of chromosome 1 genomic contig; 1226710..1227461

&lt;400&gt; 57

taaacatata tataaatata tataaatata tatataaata tatataaata tataaatata	60
taaatatata tgaatatata aatatatata aatatatatg aatatataaa tatatatata	120
aatatatata aatatatata taaatatata taaatatata taaatatata taaatatata	180
taaataaata tataaatata tataaatata taaatatata tataaatatg taaataaata	240
tatataaata tataaatata tataaatata tataaatata tatagaaata tatatagaaa	300
tatatataaa tatatataga aatatataga aatatatata gaaatatata taaatatata	360
taaatataga aatatatata aatatatata aatatatata gaaatatata atatataaa	420
atatatataa atatataaat atatataaa atatatatat aaatatatat aaatatatat	480
aaatatatat aaatatatat aaatatatat attaatatat aaatctatat taatatatat	540
taatataaa atctatatta atatatatta atatatatat taatatatat taatatataa	600
atatatatat taatatataa atatataaa atatatatgt aaatatatat ataaatatat	660
ataaatatat atataaatac atataaatat atatataaat atatataaat atatataaa	720
atatataaa atatatatat aaatatatat aa	752

&lt;210&gt; 58

&lt;211&gt; 300

&lt;212&gt; DNA

## SEL PCT 012.ST25

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(300)

<223> MAR of chromosome 1 genomic contig; 1119049..1119348

<400> 58

taatatacat ttatataat atagtgaata tatattttat atatatgtaa tatatatatt 60

atataatata tgtaatatat attttatata tatgtaatat atattttata taatatatgt 120

aatatatatt ttatataata tatgtaatat atattttata taatatatgt aatatatatt 180

ttatataata tatgtaatat atattttata taatatatgt aatatatatt ttatataata 240

tatgtaatat atattttata taatatatgt aatatatatt ttatatatat gtaatacata 300

<210> 59

<211> 617

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(617)

<223> MAR of chromosome 1 genomic contig; 3603613..3604229

<400> 59

aaaatataat atatatata tataatatat ataatatatt atatatataa tatataatat 60

## SEL PCT 012.ST25

ataatatata taataaaata tacataatat ataattgata ataaaatata cataatatat 120  
 aatatataat aaatatataa tatataatat ataataaaat atataatata taatatataa 180  
 taaaatatat aatatattat atataataaa atatataata tattatatat aataaaatat 240  
 ataatatatt atatataata aaatatataa tatattatat ataataaaat atataatata 300  
 ttatatataa taaaatatat aatatattat atataataaa atatataata tataatatat 360  
 aataatatat ataatatata atatatataa taaaatatat ataatatata atatatataa 420  
 taaaatatat aatatataat atatataata aaatatatat gatataatat atatataata 480  
 aaatatatga tatataatat atataataaa atatataata tataatataa tatataatat 540  
 atatactaaa aaatatataa tatataataa aaaatatata atatataata tatataatat 600  
 ataataaaat atatata 617

<210> 60

<211> 674

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(674)

<223> MAR of chromosome 1 genomic contig; 2592460..2593133

<400> 60

taagcttata tatatatata agcttatata tatatatata agcttatata tatatagaaa 60  
 gcttatatat atatagaaag cttatatata taagaagctt atatataaaa gcttatgtat 120  
 aaatatatat aaatatattt atttatgctt atagatacat atataaatat atttatttat 180  
 atttatatat aaacatatat ttatatatat ttatatataa tttatttatt atataaataa 240

## SEL PCT 012.ST25

atatataata aataataaat atatataata tatttattgt attatttata taaatttatt 300  
 aatataatat ataataaaat aataattata taaatatata aatatctata aatatatata 360  
 aatatatata atatctataa atatatataa atataaatat atataaatatc tataaatata 420  
 gataaatata aatatatata atatctataa atatagataa atataaatat atataactat 480  
 atataaatat atataactat atataaatat atatataaat atatataact atatatataa 540  
 ctatatatat aaatatatat aactatatat ataaatatat atataaatat atataactat 600  
 atatataaat atatataact atatataaat atatatataa atatatataa ctatatatat 660  
 aaatatatat ataa 674

&lt;210&gt; 61

&lt;211&gt; 1694

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(1694)

&lt;223&gt; MAR of chromosome 1 genomic contig; 2891680..2893373

&lt;400&gt; 61

atatgtaata catatattat atatgcatat atacatgcat atgtatatac atatattata 60  
 tatgcatata tacatgcata tgtatataca tatataaagt atgattatat ataatatata 120  
 catgtatatg tatatacatg tatatattat attatatatt atttatacat attattatgt 180  
 ctatatataa tataatatat acatattaat aatataatac ataataaat ataatatatt 240  
 atataatata taatatataa taatatatta tataatacat aatataatat aatatattat 300  
 atgatacata atataatata atatattata tgatacataa tataatataa tacatattaa 360

## SEL PCT 012.ST25

taatatatta ttattattaa tataatatat acatatattat atacatacat atatattata 420  
 ttatatataa tatacatata atataatatg taatattata tataatataa tacataatat 480  
 aatacatatt aataatatat tattaataag ataatatata tgtatctata atatatacat 540  
 atatgtatat gtatgtatat attatagata tacatgttta tacatgtata tattatagat 600  
 atatacatgt atatacatgt atatattata gatatacata tgtatatacg tatatattat 660  
 agatatacat gtatatatgt atatatatta tagatataat atatacaaga atataagaat 720  
 atatataata taatatataa tacacataat acgtatatat tatatatata tgtatattat 780  
 atatgtacat atatacatgt atattatata tacatgtata ttatatatac atgcatatta 840  
 tatatatttt tatatataat atccatgtat attatgtata ttgtgtata ttatatatac 900  
 atgtatatta tatatacatg catattatat atatttttat atataatatac catatatatt 960  
 atgtatattt gtgtatatta tatatacaca tatattatat atacatggat attatatata 1020  
 cacatatatt atatatacat atatattata tatacacata tattatatat acatgtatat 1080  
 tatatatata cgtatattat atatacacac gtatattata tatacacgta tattatatat 1140  
 acacacgtat attatatata cacgtatatt atatatacac acgtatatta tatatacacg 1200  
 tatattatat atacacacgt atattatata tacacgtata ttatatatac acacgtatat 1260  
 tatatatata cgtatattat atatacacac gtatattata tatacacgta tattatatat 1320  
 acacacgtat attatatata cacgtatatt atatatacac acgtatatta tatatacatg 1380  
 tatattatat atacatgtat attatatata cacatgtata ttatatatac atgtatatta 1440  
 tatatacaca tgtatattat atatgcatgt atattatata tacacatgta tattatatat 1500  
 acacatgtat attatatata catatatatt atatatacat gtatattatg tatacatata 1560  
 tattatatat acatgtatat tatagataca tatatattaa atatacatgt atattatgta 1620  
 tacatatata ttaaataatac atgtatattg tatatacata tatattatat acatgtatat 1680  
 tacatgtata cata 1694

&lt;210&gt; 62

&lt;211&gt; 587

## SEL PCT 012.ST25

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(587)

&lt;223&gt; MAR of chromosome 1 genomic contig; 3432560..3433146

&lt;400&gt; 62

gaattatata tatatagctg aattatatac atatataata tatacaatat atattatata 60  
tttatatatg atatatacaa tatatattac atattatata tacaatatat aatatataat 120  
atataatatt atatattata tattgtatat aatatatatt atataacatt atataatata 180  
taatattata tattatatat tgtatataat atatattata taacattata taatatatac 240  
tattatatat tataatatat aatatataat aatatataat agtatatatt atatatattg 300  
tatatattat atataaatat ataatatata atatatatta tataatatat attatataat 360  
atatattatt atatattata tatttatata taatatatat tatatatatt atattttata 420  
tataaatata taatatataa taatatataa tttaatatat ataatatata caatatataa 480  
tatataatat attaatatat ataatatata caatatataa tatataatat ataatatata 540  
atataaatta ttatatataa tatatattat atatagctga attatat 587

&lt;210&gt; 63

&lt;211&gt; 313

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

## SEL PCT 012.ST25

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(313)

&lt;223&gt; MAR of chromosome 1 genomic contig; 3805392..3805704

&lt;400&gt; 63

tatataatat gtatattatg taatatttta tatagcatat atgtatatta tatataatct 60

tttatatata gtatataata tgtatattat atattatata attatataat tatgtattat 120

ataaaatata ttatataata tataattata tattttttga aatatagatt atataataa 180

tatatggcag tgagctgaga tataatatat attatctata ctatataata tatattatat 240

atactctata ttatatatgt atatattata tataatatat acatatataa tgtgtatata 300

ttatatataa taa 313

&lt;210&gt; 64

&lt;211&gt; 349

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(349)

&lt;223&gt; MAR of chromosome 1 genomic contig; 4521378..4521726

&lt;400&gt; 64

ttatatacac tatataatat gattttatat atacttatat acactatata tgtatttata 60

tataattata tacactatat aatatgtatt tatatataat tatatacact atataatatg 120

tatttatata taattatata cactatataa tatgtattta tatataattg tatacactat 180

## SEL PCT 012.ST25

ataatgtata tttatatata attgtatata ctatataatg tatatttatg tataattgta 240

tacactatat aatgtatatatt tatgtataat tgtatacact atataatgta tatttatgta 300

taattgtata taccatataa tgtatatfta tgtataattg tatatacca 349

<210> 65

<211> 500

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(500)

<223> MAR of chromosome 1 genomic contig; 3240166..3240665

<400> 65

ttaatatata atatatafta tatatttata tattaatata taatatatat ttatatataa 60

tatatattat atatttatat tacatatatt tataigttaa tatatatftt atatatttat 120

atattttata tatttatata ttatatattt atatattata ttatatatt atatatttat 180

attatatatt tatatattat atttatatat tatatatfta tatttatatat ttatatattg 240

tatatttata ttatatattt atatattgta ttatatatt atatatttat atactatata 300

tatttatata tatttatatat ttatatatta tatatatfta tatatatatt atatttatat 360

attatatata ttatatata ttatatattt atatattata tatatttata tatatttatat 420

atatttatat atatttatata ttatatata atatatafta tatattttat ctatatattt 480

atatattaat atatattata 500

<210> 66

<211> 866

## SEL PCT 012.ST25

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(866)

&lt;223&gt; MAR of chromosome 1 genomic contig; 409429..410294

&lt;400&gt; 66

```

atatatataa tatattatat atattatata ttatatatat aatacatata ttatatatat   60
aatatataat acatatatta tatatattat atattatata taatatataa tacatatatt   120
atatataata tataatatat aatatattat ataataaat tatataatta tataatataa   180
tataatatat aatattatat aattatataa tatatataat tatattatat attataaata   240
ttatataata tatatattac aaatatatat tatatatatt ataaatatta tataacatat   300
atattatata atatatataa tatataatat atataaaaat ataatatata agatatatat   360
aatatatgat atatatgata tataatatat gatatatatg atatatataa tatatgatat   420
atatgatata tatgatatat ataatatatg atatatatga tatatatgat atatgatata   480
tatgatatat gatatatatg atatatatga tatatgatat atatgatata tatgatatat   540
gatatatatg atatatatga tatatgatat gatatatata atatatgata tgatatatat   600
aatatatgat atatatgata tatgatatgt aatatatatg atatattata tataatatat   660
aatatataca taatatataa tatataatat ataatatata taatatgtga tatatataat   720
atatgatata tgatatatga tatatattat ataatatata taatatatat tatatataat   780
atatattata taatatatat aatatatatt atatataata tataagatat aagatataat   840
atatataata tataatatat ataata                                     866

```

&lt;210&gt; 67

## SEL PCT 012.ST25

<211> 335

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(335)

<223> MAR of chromosome 1 genomic contig; 614754..615088

<400> 67

acccaatata tgtgtatata tgtatgtata tatacatata catacatata tatatgtaca 60

tacatatata catacatata tatatatgta catacatata tacatacata catatatata 120

tataacatat atacacacat atatacagat atacatatat acatacatat atacatataa 180

catatatata tacatatata catataaacac atacatacat acatatatac atacaacata 240

tatacatata tatatacata tgtatacata catatatgta tacatatatg tatacatata 300

tgtatacata tatgtatata tatattgtta tatat 335

<210> 68

<211> 455

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(455)

<223> MAR of chromosome 1 genomic contig; 1299520..1299974

## SEL PCT 012.ST25

&lt;400&gt; 68

ggatatatat attattagtt gttatattat tatatattat atatattatt atatataata 60  
 tattatatca tatatattat tatatataat atattatatac atatatatta ttatataata 120  
 tattatatca tatatattat tatatataat atatattata tatattatta tatataatat 180  
 atattatata tattattatg tataatatat atattatata ttatttatat atatataaat 240  
 tatataataa tatataatta attatacata tatacatata taagtataca tataatatat 300  
 ttatatagta tatataaata tatatacaat atatttatat attatatatt atatataaat 360  
 atatacaata tatttatatc atatatttta tatatgatac atataatata tattatatat 420  
 gatataaat atatatcata tatgatatat aacat 455

&lt;210&gt; 69

&lt;211&gt; 404

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(404)

&lt;223&gt; MAR of chromosome 1 genomic contig; 1970778..1971181

&lt;400&gt; 69

atatataata tgtataatat ataatatata tcatatattg ttctatgtat attacatata 60  
 atatgcatta tatatttatat attgcatata atatgcatta tatatttatat attgcatata 120  
 atatgcatta tatatttatat attgcatata atatgcatta tatatttatat attgcatata 180  
 atatgcatta tatatttatat attgcatata atatgcatta tatatttatat attgcatata 240

## SEL PCT 012.ST25

atatgcatta tatattatat aatatatata catataatat atataattta tatatattta 300

tatatattta catttattat atatttatta tatataaata tatttttata tattacttat 360

atattatata taatatatat aatatatata ttatatataa tata 404

<210> 70

<211> 605

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(605)

<223> MAR of chromosome 1 genomic contig; 3562918..3563522

<400> 70

tatatatata aaatacatat atattatata tattatatat aatacatata ttatatatta 60

tatataatac acgtatataa tatataatat ataatacata taatatatat gatataaat 120

acataataa tatatgatat ataatacata tataatatat atgatatata atacatatat 180

aatatatatt atatataata catatataat atatattata tataatacat atataatata 240

tattatatat aatacatata taatatatat tatatataat acatatataa tatatattat 300

ataatacata tataatatat attatataat acatgtatat aatatatatt atatataata 360

catatatatt atataatata tgtatataat atatattata tataatacat atatattata 420

tattatatat taatatattt atataatagt aatatataat attaatatat tatatatatt 480

aatattatat ataatacata tattatatat aatataaata tatataatac atatataata 540

cacatattat atataatata tatattatat ataatatata tattatatat aatatatatg 600

taata

605

## SEL PCT 012.ST25

&lt;210&gt; 71

&lt;211&gt; 317

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(317)

&lt;223&gt; MAR of chromosome 1 genomic contig; 189743..190059

&lt;400&gt; 71

tattttttat atttatatat tatatatatt ttatatgta atatattata tataaaatta 60

tataatttta ctacatataa tatataaaat tatataattt tactacatat aatatataaa 120

attatataat ttactatat ataatatata aaattatata attttatata taatatatat 180

tataatatat atttatgca atatatatta tatattatat tataatatat tgtatatattt 240

tgtatataaa atatataata tataatatat ttatagacaa taatatataa tataatatat 300

aaaattttat atataaa 317

&lt;210&gt; 72

&lt;211&gt; 522

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(522)

## SEL PCT 012.ST25

<223> MAR of chromosome 1 genomic contig; 229111..229632

<400> 72

gatatatata tttatatata taaaagatat atattattta tatataaaga tatatattta 60  
 tatatatata agatatatat ttttatata tataaaagat atatatttat atatatgata 120  
 tatattattt atatatataa aagatatata tttatatata tgatatatat ttttatata 180  
 taaaagatat atataaaaga tatatattat ttatatatat aaaagatata tatataaaag 240  
 atatatatta tttatatata taaatgatat atattattta tatataaaag atatatatta 300  
 tttatatata aaagatatat attatttata tatataaaag atatacatat aaaagatata 360  
 ttttatata taaaagatat atatatttat atataaaaga tacatatatt tatatatata 420  
 aaagatatat atatttttat atataaaata tatattatat atataaaaga tatatatata 480  
 tatatatatc ttttatatat aaaagatata tataaatata ta 522

<210> 73

<211> 1110

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(1110)

<223> MAR of chromosome 1 genomic contig; 1138030..1139139

<400> 73

tatgtatgta tacataatat attatatatg tatattatgt atacataata tattatatat 60  
 gtatattatg tatacataat atattatata tgtatattat gtatacataa tatattatat 120

## SEL PCT 012.ST25

attatatgta tattatgtat acataatata ttatatatta tatgtatatt atgtatacat 180  
 aatatattat atattatatg tatattatgt atacataata tattatatat tatatgtata 240  
 ttatgtatac ataatatatt atatattata tgtatattat gtatacataa tatattatat 300  
 attatatgta tattatgtat acataatata ttatatatta tatgtatatt atgtatacat 360  
 aatatattat atattatatg tatattatgt atacataata tattatatat tatatgtata 420  
 ttatgtatac ataatatatt atatattata tgtatattat gtatacataa tatattatat 480  
 attatatgta tattatgtat acataatatt tatatattat atgtatatta tgtatacata 540  
 atatattata tattatatgt atattatgta tacataatat gtacacataa tattatata 600  
 ttatatgtat attatgtata cataatatt atatattata tgtatattat gtatacataa 660  
 tattatata ttatatgtat attatgtata cataatatt atatattata tgtatattat 720  
 gtatacataa tattatata ttatatgtat attatgtata cataatatat tatatattat 780  
 atgtatatta tgtatacata atatattata tattatatgt atattatgta tacataatat 840  
 attatatatt atagtatat tatgtataca taatattat atattatatg tatattatgt 900  
 atacataata tattatatat tatatgtata ttatgtatac ataatatatt atatattata 960  
 tgtatattat gtatacataa tatattatat attatatatg tatattatgt atacataata 1020  
 tattatatat tatatatgta tattatgtat tatattatat attatgtata ttatagatta 1080  
 tgtatgcata cataatatgt attgtatatt 1110

&lt;210&gt; 74

&lt;211&gt; 521

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(521)

## SEL PCT 012.ST25

<223> MAR of chromosome 1 genomic contig; 2863407..2863927

<400> 74

```

aatatatata aatatataaa tatatatataa tatatatata tataaatata taaatatata    60
tatgtaaata tatgtaaata tatgtaaata tatgtatatg tatatatatg taaatgtatg   120
taaatatata taaatatatg taaatatata taaatatatg taaatatata aatatatata   180
actatatata aatatatata aatatataaa tataaatata tataaatata tataaatata   240
taaataaata catataaata tataaataaa tacatatata tatatatata tatataaaaa   300
tatatatataa tatatatata aatatataaa catatatata tatataaata tatataaata   360
tataaatata taaaatatat aaatatatat aaatatataa atatatatata atatagataa   420
atatagataa atatataaat atatataaat atataaatat agataaatat ataaatatat   480
aaatatataat atataaaaat atatataaat atataaaaat a                        521

```

<210> 75

<211> 560

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(560)

<223> MAR of chromosome 1 genomic contig; 5712303..5712869

<400> 75

```

ataataattat atatataatta tatattatat ataattatat attatatata atgtataatt    60
atatattata tataatatat ataaatatat atatttttta tataaatata ttatatattt   120

```

## SEL PCT 012.ST25

atatattata tataaattta tatatatataa ttttatata ttatatatat ttatatatta 180  
 tatattgtat atatttatat attacatatt gtatatattt atatattata tattatatat 240  
 ttatatatta tatatttatat atttatatat tatatatatt atatatttat atatttatata 300  
 taaattttt atatataata tataaatata tattatataa tataaatttg tatatatata 360  
 atatatttat attatatata aaatatttat attatatata aaatataata taaatatata 420  
 catataatat atatattata tatttataat tatatatatt atataataca tataatatat 480  
 aatatataat acatatatat catatatgaa atatatatca tatattatac atatttatata 540  
 taacatatat attatatatc 560

<210> 76

<211> 479

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(479)

<223> MAR of chromosome 1 genomic contig; 8578812..8579290

<400> 76

tatggtatac atatagtata tatggggtac atatatggta tatatatggg ttatatatat 60  
 gatatatatt atatatgtat atggtatata tatggtatat atattataca tgcatatggt 120  
 atgtatatgg tatatatatg atatatacat atggtgtata tatatgttat atatgatata 180  
 tataagggtat atatatggta tatataagggt atatatagta tatatatgggt atatataagg 240  
 tatatattgt atatatatgg tatatataag gtatatatat tgtatatatg gtatatatat 300  
 ggtttatata tatggtgtgt atatatgggt ttatatataca cactttatat actatatatt 360

## SEL PCT 012.ST25

atatacacac tatatataat atatattata tatagttaaa tatatggat atgcaattag 420

atatatggta tatgtaatta tatatatggg atatagatgg tgtatatatg gtatatata 479

<210> 77

<211> 477

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(477)

<223> MAR of chromosome 1 genomic contig; 8579294..8579770

<400> 77

tatagtatat atacacacta taggtaatat actacatatt atatacacac tataaataaa 60

atatataata tataatattt tctatatagt atatattata tattgtatat actatatata 120

atataacta tagacagtag atactttata tactatagac agtatatact atatactgta 180

tacactatag acagtatata ctatatactg tatacagtat atgtagtgta tatgtagtgt 240

atataatata tagtatatat tatctatact atatacagta tatatagtgt atacataata 300

tatattatat attatatata ctatatacag tatacatagt gtatatgtag tgtataatat 360

atataatgtg tatataaaat atatatacta tatataatat atattatata taatatatac 420

actatatata ctatagatac actatatatt cactatatat actatatata ctatata 477

<210> 78

<211> 331

<212> DNA

<213> Homo sapiens

## SEL PCT 012.ST25

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(331)

&lt;223&gt; MAR of chromosome 1 genomic contig; 8580024..8580354

&lt;400&gt; 78

actatatggt atatacataa gatatagtat ataccatata ttatatcat tatatatagt 60

gtatactata tataatgtat ataatatata gtatatatac actatatata ctatgtatat 120

atacactata tatactatgt atatatacac tatatatact atgtatatat acactatata 180

tactatgtat atatacacta tatatactat gtatatatac actatatata ctatgtatat 240

atacactata tatactatgt atatatacac tatatatact atgtatatat agtgtatata 300

tactgtatat gttatagtggt atatatagta t 331

&lt;210&gt; 79

&lt;211&gt; 410

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(410)

&lt;223&gt; MAR of chromosome 1 genomic contig; 8580705..8581114

&lt;400&gt; 79

tatagtctat attatataca gtctatataa tatatagtat atactatata tacttttcct 60

## SEL PCT 012.ST25

cattctgact atatactata tatatactat atatagtata ttagtgat atatacacca 120  
 tatatactat atatagtata cataccatat atagtatact atacatacca tatatagtat 180  
 acataccgta tatagtatac tatacttacc atatatagta tacatactat atataatata 240  
 tctgggtgat atatacacta tatatactat atatactata tatagtatat gtacactatt 300  
 tatagtattt atagtatata tactgtatat atagtatga gtatatatac tatatattat 360  
 gtagactata tataatatag actatgtgta gagtatatat actatatata 410

&lt;210&gt; 80

&lt;211&gt; 433

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(433)

&lt;223&gt; MAR of chromosome 1 genomic contig; 12979167..12979599

&lt;400&gt; 80

atatataata tatatatgct ctatatataa aatatatcat atataaaat atatatgata 60  
 tttttatat attaaatata taattatata taaatatata tttatatata aatatattat 120  
 ttcaatatat ataaatatat ttaaatatat ttaaatagaat tattaaatat ataaatatat 180  
 aattatattt aatatataaa tatatattaa atatatatt atatttaata tatataaata 240  
 tatattaaat atataattat atatttatat atttattata tataaatata ttttgttct 300  
 aaataaatat atattctaaa tatataatat tttatattat ataatatata atataaaata 360  
 tataataaat atataatata taaataaata aatatttatt ataaaatata tataaatatt 420  
 aaatatatat taa 433

## SEL PCT 012.ST25

&lt;210&gt; 81

&lt;211&gt; 385

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(385)

&lt;223&gt; MAR of chromosome 1 genomic contig; 16336644..16337028

&lt;400&gt; 81

```

tttatataaa tatctatata aataaatata taaatatata aatataaata tatataaata   60
tataaataaa tatataaata tatataaata taaatatata tataactatg aatttatatt   120
tatataaata tatatctata tgaatataaa tatatattta tataaatata aatatatata   180
taaatatata tatttatata gatataaata tatatatataa tatatatatt tatatagata   240
taaatatata tctatatatg aatatatatc tataggaata taaatatata tctatataaa   300
tataaatata tataagtata aatatatata aatatatatc tatataaata taaatatata   360
tataaatata aatatatata taaat                                     385

```

&lt;210&gt; 82

&lt;211&gt; 363

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

## SEL PCT 012.ST25

&lt;222&gt; (1)..(363)

&lt;223&gt; MAR of chromosome 1 genomic contig; 20624448..20624810

&lt;400&gt; 82

tatatatata gttatatata tatttatata tatagttata tatatatattt tatatagtta 60

tatatatagt tatatatata gttatatata tatagttata tatatagtta tatatatagt 120

tatatatata tagttatata tatagttata tatatagtta tatatatagt tatatatata 180

tagttatata tatagttata tatatatagt tatatatata gttatatata tatagttata 240

tatatagtta tatatatagt tatatatata gttatatata tagttatata tatatagtta 300

tatatatata gttatatata tatagttata tatatagtta tatatatata gttatatata 360

tag 363

&lt;210&gt; 83

&lt;211&gt; 310

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(310)

&lt;223&gt; MAR of chromosome 1 genomic contig; 566025..566334

&lt;400&gt; 83

tatatataat atatattgta tatattatat attgtatata taatatatat tgtatatatt 60

atatattgta tatataatat atattgtata tattatatat tgtatatata atatatatat 120

tgtatatatt atatattgta tatataatat atatattgta tatattatat attgtatata 180

## SEL PCT 012.ST25

taatatatat attgtatata ttatatattg tatatataat atatatattg tatatattat 240

atattgtata tataatatat atattgtata tattatatat agtatatatt atatatagta 300

tatataatat 310

<210> 84

<211> 1236

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(1236)

<223> MAR of chromosome 1 genomic contig; 1171429..1172664

<400> 84

aaagtattat atgtattata tgtatatgta ttatatatta catatgtatt atataata 60

tatattatat attattatat attatatatt atatattatt atttatataa tgtattatat 120

attatatagt atatatagta tatataatgt attatatatt atatagtata tatagtatat 180

ataatgtatt atatatagta tatataatgt attatatagt atataacta tataatgtat 240

tacatattat gtatagtata tgtaatgtat tatatattat atagtatatg taatgtatta 300

tatgtattat atagtatata ttatatatga tgtattattt agtatatata atatatatga 360

tgtattatat aacatatata atatatatga tgtattatat agcatgtata gtatatatga 420

tgtattatat agcatgtata gtatatatga tgtattatat atagcatgta tagtatatat 480

gatgtattat atatagcatg tatagtatat atgatgtatt atatatagca tgtatagtat 540

atatgatgta ttatatatag catgtatagt atatatgatg tattatatat agcatgtata 600

gtatatatga tgtattatat atagcatgta tagtatatat gatgtattat atatagcatg 660

## SEL PCT 012.ST25

tatagtatat atgatgtatt atatagca tgtatagat atgatgta ttatatatag 720  
catgtatagt atatatgatg tattatatat agcatgtata gtatatatga tgtattatat 780  
attatatatg gtatatatga tgtattatat attatatatg gtatatatga tgtattatat 840  
attatatatg gtatatatga tgtattatat attatatata atatatatga tgtattatat 900  
attatatata atatatatga tgtattatat atgatgtatt atatataata tatatgatgt 960  
attatatata ttattatcta ttatatacga tgtattatat gcaagttatt atgtataata 1020  
tataatgtat tatatattat ataagtata atatataaat atataaatat ataattatgt 1080  
ataaatatag aaatatatac attatacatt atatacatta taatgtataa tatataaata 1140  
tattatatat aaatgtatac attatatata aatatattat atacattata tataaaatat 1200  
gtatatagtt attatacctt atatatacta aacagt 1236

&lt;210&gt; 85

&lt;211&gt; 309

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(309)

&lt;223&gt; MAR of chromosome 1 genomic contig; 1925173..1925481

&lt;400&gt; 85

atatatttat ataaatatat ttatatataa tatatattat ataattattat aatatatgtt 60  
atattatata tatatttatac aatatataat atatattata tatattttat acaatatata 120  
atatatatta tatatatattt atataatata taatatatat tatatatatt ttatacaata 180  
tataatatat attatatatt atataatata tattatatat attttatata atatatataa 240

## SEL PCT 012.ST25

tatatattat acaatatata atgtatatca ttatattata taatgtatat catattatat 300

aatgtatat 309

<210> 86

<211> 312

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(312)

<223> MAR of chromosome 1 genomic contig; 4396756..4397067

<400> 86

cacagtgtat atatagtata tatactgtat atatactgtg tatatacact gtatatacac 60

agtgtatata cagtatatat actatatata cactgtgtat atatagtata tataaattot 120

aggaatatat atactatata tatactatat atataaattc taggaatata tacacactat 180

atatacacta tatatacaca tatatacact atatataffa tacacatata ttatatatat 240

acactatata tacacgagat atataacata tacactatat actatacata acatatatac 300

tatatatact at 312

<210> 87

<211> 398

<212> DNA

<213> Homo sapiens

<220>

## SEL PCT 012.ST25

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(398)

&lt;223&gt; MAR of chromosome 1 genomic contig; 56057..56454

&lt;400&gt; 87

atatatatta catattatat atataatata tattatataa tatatattat attatataat 60

atataatata aatataatat aaattatatt atataatata taatataaat ataataataa 120

ttatataaat ataatatata ttttattata taatataata tatattatat aaatataata 180

tataaattat ataataatat atatattata taatataata tattttatta tataaatata 240

tattatatta tataatatat attttattat ataatatata ttatatattt atagaatata 300

atatatattt tattatataa tatatattat ataatatata ttatatattt atataacata 360

tattattata taaaatatgt ataatatata ttatataa 398

&lt;210&gt; 88

&lt;211&gt; 391

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(391)

&lt;223&gt; MAR of chromosome 1 genomic contig; 56984..57374

&lt;400&gt; 88

tactataata catattatat ataattattat atactatata ttactatatt attatattat 60

atataattaa actatattat agtatataat atataatata tactatatgt aatattacta 120

## SEL PCT 012.ST25

tgatactgat attatattat atataattaa attatattat attaatatat aaattatata 180  
 taatacataa tatataaatt atattatatt atttatatat aatgtatgcc atataattta 240  
 tatataatgc atttatata atttatatat aatgcattaa atataaatta tatataatgc 300  
 atttatata atttatata atgcattata tataatttat attaatata taaatttata 360  
 tttaatatat ttatatatta tatataataa a 391

&lt;210&gt; 89

&lt;211&gt; 309

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(309)

&lt;223&gt; MAR of chromosome 1 genomic contig; 469547..469855

&lt;400&gt; 89

atatatatgt aatatatatg ttatatatgt aatatatatg ttatgttata tatgttatat 60  
 atatgttata tataatatat atgttatata tacgttatat gttatatata tgttatatat 120  
 aatatatggt atatatatgt tatatgttat atatgttata tataatatat gttatatata 180  
 atatatgtta tatatgttat atataatata tgttatatat atttatata atatatgtta 240  
 tatatattat atataatata taatatatgt gatataaat ataaaatata tgtgatatat 300  
 atttatatat 309

&lt;210&gt; 90

&lt;211&gt; 441

&lt;212&gt; DNA

## SEL PCT 012.ST25

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(441)

&lt;223&gt; MAR of chromosome 1 genomic contig; 546190..546630

&lt;400&gt; 90

atacacaaca tatgtgtata tatatagtat atatacaca catatgtgta tatatatagt	60
atatatacac aatatatgtg tatatatata gtatatatac acaatatatg tgtatatata	120
gtataaatat atactatata tagtatatat agtataaata taaactatat atagtatata	180
catagtataa atatatacta tatatagtat atacatagta taaatatata ctatatatag	240
tatatacata gtataaatat atactatata tagtatatac atagtataaa tatatactat	300
atatagtata tacatagtat aaatatatac tatatatagt atatacatag tataaatata	360
tactatatat agtatataca tagtataaat atatactata tatagtatat acatagtata	420
aatatatact atatatagta t	441

&lt;210&gt; 91

&lt;211&gt; 1367

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(1367)

&lt;223&gt; MAR of chromosome 1 genomic contig; 124643..126009

## SEL PCT 012.ST25

<400> 91  
 atatttatat gatataaat atatataata ttatatataa tattatatat gatataaac 60  
 attatataat attatatatg atatataatta tatatattat atatgatata taatatatat 120  
 aatatttatat atgatattat atatcatata taatatataa aatatttatat atgatatata 180  
 atatataataa tattatatat attatatata ttatatatca tatataatat tctaaatata 240  
 taatattata tgatatataa gattatatac attatatata atatataata ttatatatga 300  
 tatataatat tatatacaatt atatataata tataatgtat ataattat atattatata 360  
 ttatattat atacaatgta tataatatta tatatcatat atatttatat tatatacaat 420  
 gtatataata ttatatatca tatataatat tatatacaat gtatataata tatatttatat 480  
 atatttatat tatatacaat gtatataata tatatttatat atatttatat tatatacaat 540  
 gtatataata tatatttatat atatttatat tatatacaat gtatacaata ttatatatta 600  
 tatatttatat atttatatta tatacaatgt atatattata tattatatat ttattatata 660  
 tacaatgtat atatttatata ttatatatt atatttatata caatgtatat attatatatt 720  
 atatatttat attatataca atgtatatat tatatattat atatttatat tatatacaat 780  
 gtatatatta tatatttatat atttatatta tatataatgt atgtaatt atatattata 840  
 tatttatatt atatataatg tatgtaatat tatatattat atatttatat tatataaat 900  
 gtatgtaata ttatatatta tatatttata ttatatataa tgtatgtaat attatatatt 960  
 atatatttat attatatata atgtatgtaa tattatatat tatatattta tattatatat 1020  
 aatgtatgta atatttatata ttatatatt atatttatata taatgtatgt aatatttatat 1080  
 attatatatt tatatttatat ataatgtatg taatattata tattatatat ttattatata 1140  
 tataatgtat gtaattat atatttatata ttattatatt atataatgta tgtaattatta 1200  
 tatatttatat atttatatta tatataatgt atataatt atatttatata tatttatatt 1260  
 gtatataata ttatatatta tatatttata ttatatataa tatatattat atatttatat 1320  
 tgtatataat attatatatt atatatttat attatatata atgtata 1367

## SEL PCT 012.ST25

&lt;210&gt; 92

&lt;211&gt; 458

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(458)

&lt;223&gt; MAR of chromosome 1 genomic contig; 58908..59365

&lt;400&gt; 92

tatatgatat atatgatata tatgggatat atatgatata tatgatatat atggtatata	60
tatgatatat agtatatatg atatatatgg tatatatatg atatatagta tatatgatat	120
atatggtata tatgatatat agtatatatg atatatatgg tatatatggt atatatatga	180
tatatgatat atatgatata tatgatatat gatatatatg atatatatga tatatatggt	240
atatatgata tatatggtat atatggtata tatatgatat atatgatata tatggtatat	300
atatgatata tatgatatat atggtatata tatgatatat atgatatata tggatatat	360
atgatatata tgatatatat ggtatatata tgatatatat gatatatatc atatatatgg	420
tatatatatg atatatatga tatatatcat atatatgg	458

&lt;210&gt; 93

&lt;211&gt; 330

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

## SEL PCT 012.ST25

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(330)

&lt;223&gt; MAR of chromosome 1 genomic contig; 306867..307196

&lt;400&gt; 93

ataatatata aatatatg atatatatct atatatatca tatataaata tatatgatat	60
atatctatat atatcatata taaatatata tgatatataa atatatatga tatatatcta	120
tatatatcat atataaatat atatgatata taaatatata tgatatatat ctatatatat	180
catatataaa tatatatgat atatatctat atatcatata taaatatata tgatatatat	240
ctatatatat catatataaa tatatatgat atctatctat atatatcata tataaatata	300
tatgatatct atctatatat atcatatata	330

&lt;210&gt; 94

&lt;211&gt; 353

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(353)

&lt;223&gt; MAR of chromosome 1 genomic contig; 636899..637251

&lt;400&gt; 94

tatgtatata tatacacata tacgtatata tatacatata tacacatata cgtatatata	60
tacgtatata tacatatgta tatgtatacg tatacacaca tatgtatatg tatacgtata	120
cacacatata cgtatatatg tatacgtata cacacatata cgtatatgta tacatatata	180

## SEL PCT 012.ST25

tggtacata tacgtatata cgtatatgta tacatatata cgtttatgta tatatacgta 240

tatacgata tatgtatatg tatacatata tacatatatg tgtatatatg tatatacgta 300

tatgtgtata tatacaatat acatacatgc acatatatgt gtatatgcac ata 353

<210> 95

<211> 345

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(345)

<223> MAR of chromosome 1 genomic contig; 1435510..1435854

<400> 95

atcatatata ttatatatca tatatatgat atataaaaat tatatatcat atatatgata 60

tataaaaatt atatatatca tatataatat atataatata ttatatatat aaattatata 120

taatatatat aaattatata tatcatatat atgatatata atttatatat catatatatg 180

atatatataa tatattattt atatataata tattatatat tatataatat gtaatatata 240

ttatatatta catattatat tttttataaa taatatttta taatatatat aatattatat 300

aatatagaat atttatatatt atatattaca tattatataa tatat 345

<210> 96

<211> 521

<212> DNA

<213> Homo sapiens

## SEL PCT 012.ST25

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(521)

&lt;223&gt; MAR of chromosome 1 genomic contig; 39695..40215

&lt;400&gt; 96

tatatatata atagatatta tatatctatt atatatctat tatatatata atagatatta	60
tatatctatt atatatataa tagatattat atatctatta tatatataat agatattata	120
tatctattat atataatata tatctattat atattatata tctattatat ataatatata	180
tctattatat atattatata tctattatat atataataga tattatatat ctattatata	240
taatatatat ctattatata ttatatatct attatatata tgtatctatt atatatatta	300
tgtatctatt atatatataa tatatctatt atatatatat tatatataat atatattata	360
tatattatat atctattata tataatatat atctattata tatattatat atctattata	420
tatattatat atctattata tataatatat atctattata tatattatat atctattata	480
tataatatat attatatata tattatatat tgtatatcta t	521

&lt;210&gt; 97

&lt;211&gt; 484

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(484)

&lt;223&gt; MAR of chromosome 1 genomic contig; 1286007..1286490

## SEL PCT 012.ST25

&lt;400&gt; 97

atatcatata tatttatatat catatatatg atatataaaa atttatatc atatatatga 60

tatatatata ttatatatat catatataat atatataata tatttatatat ataaattata 120

tataatatta tatataaatt atatatcaca tatatgacat ataaattata tatcacatat 180

atgatatata atttatatat cacatatatg atatataatt tatatatcat atatatgata 240

tataatttat atacatatata tatgatatat aatttatata tcatatatat gatatatata 300

atatattatt tatatataat atatttatata ttatataata tgtaatatat atttatatt 360

atataatatg taatatatat tatatattac atatttatatt atttataaat aatattttat 420

aatatatata atatttatata atatagaata ttatatatta tatattacat atttatataat 480

atat 484

&lt;210&gt; 98

&lt;211&gt; 244

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(244)

&lt;223&gt; MAR of chromosome 1 genomic contig; 73556..73879

&lt;400&gt; 98

atttatatt atatttatata atatataata atatttatata atttatatt acatttatata 60

atatataata atatttatata ataatatata atttatataat atataataat atttatataat 120

atttatataat atttatataat atataaatat ataataatat atatttatatt atataatagt 180

atatattata ttatataata tatgttatta tatttatataa tataaactat tatataatat 240

SEL PCT 012.ST25  
244

aata

&lt;210&gt; 99

&lt;211&gt; 463

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(463)

&lt;223&gt; MAR of chromosome 1 genomic contig; 179038..179500

&lt;400&gt; 99

tacaatatat ttctattat atatatttg tattatatat aatatacaat atattttcta 60

ttatatataa tatatttgt attatatata ttacaatata tttgtatta tataatatat 120

aatacaatat aatatattgt attatataat atataatact atataatata ttgtattata 180

tattatatat aatactatat aatatatttt attatatatt atatataata ctatataata 240

tattttatta tatattatat ataatacaat atataatata ttgtattata atacaatgta 300

ttataatgta ttatatataa tatataatac aatatataat attatatata tttatatata 360

tatatatatt gtattatata tttgtatta tatatatatt gtattatata ttatatattt 420

atattataat tatgttttgc attatatatt tcatattata tat 463

&lt;210&gt; 100

&lt;211&gt; 390

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

## SEL PCT 012.ST25

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(390)

&lt;223&gt; MAR of chromosome 1 genomic contig; 55617..56006

&lt;400&gt; 100

```

tgtataatat atataacttta tatataatat atataacttta tatatatact atataactaat   60
atatataata tatactatat ataatatata ctaatatata taatatatac actatatata   120
atatatacta atatatatta tatatacttt atataatata tactaatata tataatatat   180
atactttata tataatatat actaatatat ataatgtata tactttatat ataatatata   240
ctaatatata atatatatac ttatatata atatatacta atatatatta tatatacttt   300
atatatataa tatatactta tatattatat atgcttatat ataatatata cactaatata   360
taatatatat actttatata ttatatatta                                     390

```

&lt;210&gt; 101

&lt;211&gt; 582

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(582)

&lt;223&gt; MAR of chromosome 2 genomic contig; 1157405..1157986

&lt;400&gt; 101

```

tgtatatgta tatatacaca tacgcacata tatgtatatg tatatataca catacgcaca   60

```

## SEL PCT 012.ST25

tatatgtata tgtatatata cacatacgca catatatgta tatgtatatg tatatgtata 120  
 tatacacata tacacatata tgtatatgta tatatacaca tatacacata tatgtatatg 180  
 tatatatata catatacaca tataatgtata tgtatatata cacatacaca tatatgtata 240  
 tgtatatgta tatatacaca tacacatata tgtatatgta tatgtatatata tacacatata 300  
 cacatatata catatatgta tacatatatg tgtatatata tacacatata tatacatata 360  
 tgtatacata tatgtgtata tatacacata tatatacata tatacatata catatatatg 420  
 tgtatgtata tatacacata tacatatata tgtatatgtg tatatatatt agacagatat 480  
 atatgtacat atacatatat atgtatatgt atatgtatat gtatatgtat atgtatatgt 540  
 atatgcatat ataatacaca tatacatata tgtatatgta ta 582

<210> 102

<211> 322

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(322)

<223> MAR of chromosome 2 genomic contig; 1858638..1858959

<400> 102

acacatatata tacacatatat atatacatat catatatata ccatatatat acataccata 60  
 tatataccat atatatacat accatatata cccatatat atacatacca tatatatata 120  
 ccatatatat acataccata tatataccat atatatacat accatatata tacatatat 180  
 atacatacca tatatatata ccatatatat acataccata tatatacacc atatatatat 240  
 atacatatata tatacatatat atacacata tatatacacc atatatacac accatatata 300

## SEL PCT 012.ST25

ccatatatat acaccatata ta

322

&lt;210&gt; 103

&lt;211&gt; 914

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(914)

&lt;223&gt; MAR of chromosome 2 genomic contig; 5712196..5713109

&lt;400&gt; 103

aaatatatat tctatatata gaaaatatat attctatata tatagaatat atatagaata 60  
 tatattctat atatattcta tatatataga atatatatat aaaacatata ttctatatat 120  
 aaaatatata ttctatatat ataaaatata tattctatat atatagaatg tatataaaat 180  
 atatattcta tatatataga atgtatataa aatatatatt ctatatatat agaattgata 240  
 taaaatatat attctatata tatagaatgt atataaaata tatattctat atatatagaa 300  
 tatatataac atatatatga aatatatata aaatatatat aaatacatat ttctatatat 360  
 aaatatatat aaatacatat ttctatatat aaatatatat caatacatat ttctatatat 420  
 aaatatatat aaatatatat tcatatatat aaaaatatat aaatatatat tcatatatat 480  
 aaaatatata tgaatatata ttctctatat ataaaatata tataatatat attatatata 540  
 taaaatatat ataatatata ttatatatat aaaatatata taatatatat tcatatatat 600  
 aaattatata taaatatata ttcatatata taatatatat aaatatttat ttcatatata 660  
 aaatatattt aaatatatat ttctatatag aatatatatt ctatatataa aatatatata 720  
 taaatatatt ttctatatag aaatatatat gaaatatata gaatatatat aaatatatat 780

## SEL PCT 012.ST25

tatatatact atatatacaa tatatattat atataaaata tatatacaat atatattcta 840

tatattaata tatagaatat atattaacat atatttcaat atattaatat atgaaatata 900

tataaatatt tcat 914

<210> 104

<211> 370

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(370)

<223> MAR of chromosome 2 genomic contig; 5713613..5713982

<400> 104

tatttcatat ataatatata tataaaatat atatttcata tacataatat atataatata 60

aataaaatat atatttcata tatataatat atataatatata tataaaacat atatttcata 120

tataatatat ataaactata tatttcatat ataatatata taaactatat atttcatata 180

cataatatat ataatatata tttcatttat attatatata taatatatat ttcatatata 240

taatatataa aatagatata aatatatata aatatatatatt tcatatatata tatatatataa 300

atatatatata atatatatatt tatatatatat atatatatatt catatatataa tataaaaaaa 360

tatatatttc 370

<210> 105

<211> 442

<212> DNA

<213> Homo sapiens

## SEL PCT 012.ST25

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(442)

&lt;223&gt; MAR of chromosome 2 genomic contig; 7481647..7482088

&lt;400&gt; 105

atataaatta tataatatgt tatataatat ataaatatat tatataacat gttatataat 60

atataacatg ttatataata tataacatgt tatataatat ataacatggt atataatata 120

taacatgfta tataatatat tatgtaatat gttatataat atataatata ttatataaca 180

tggtatataa tatataacat gttatataat atggtatata atatataaat atattatatt 240

atatgttata taatatataa atatattata ttatatgfta tataatatat aaatatatta 300

tattatatgt tatataatat ataaatatat tatattgtat gttatataat atataaatat 360

attatattgt atgttatata atatataaat atattatatt gtatgttata taatatataa 420

atatattata ttatatatgt ta 442

&lt;210&gt; 106

&lt;211&gt; 338

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(338)

&lt;223&gt; MAR of chromosome 2 genomic contig; 9594557..9594894

## SEL PCT 012.ST25

&lt;400&gt; 106

tatataaata tataccatat atataaatat atatattcca tatataaata tatatattcc 60

atatatataa atatatatat tccatatata aatatatata ttccatatat ataaatatat 120

atataaatat atatattcca tatatataaa tatatatata aatatatata ttccatatata 180

aatatatata tattccatat ataaaaatat atatatattc catatatataa aatatatata 240

tattccatat atataaatat atatatattc catatatata aatatatata tattccatat 300

atataaatat atatatattc catatatata aatatata 338

&lt;210&gt; 107

&lt;211&gt; 364

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(364)

&lt;223&gt; MAR of chromosome 2 genomic contig; 10519720..10520083

&lt;400&gt; 107

ttatatatat ttataaatat atatataagc tatatatatt tatatataat atattatata 60

tattagctat atatatttat ataataatat attatatatt agctatatat atttatatat 120

aataatatat ataagctata tatttatata tattatatat tagctatata ttttatata 180

taatataffa tatattagct atatatttat atataataaa taatatatat attagctata 240

tatatttata tataataata tatataagct atatatttat atataatata ttatatatta 300

gctatatata ttatatata ataatatatt atatattagc tatatatatt tatatataat 360

atat 364

## SEL PCT 012.ST25

&lt;210&gt; 108

&lt;211&gt; 342

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(342)

&lt;223&gt; MAR of chromosome 2 genomic contig; 11481943..11482284

&lt;400&gt; 108

tacatataat atataattat atataatata tattatatat tacatatata atatatatat 60

tacatatgta atatatatat tatatatgta atatatatta tatatgtaat atatatatta 120

tatatgtaat atatattata tatatgtaat atatatatta tatatgtaat atatatatta 180

tatgtaatat atatatgtaa tatatatata atatatatgt aatatatata taatatatat 240

gtaatatata tataatatat atgtaatata tatattatat atatgtaata tatatcatat 300

atatgtaata tatatcatat atatgtaata tatatcatat at 342

&lt;210&gt; 109

&lt;211&gt; 415

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(415)

## SEL PCT 012.ST25

<223> MAR of chromosome 2 genomic contig; 13499598..13500012

<400> 109

tatatatata tatatatata atataatata atatatatat aaatatatat aatataaatt 60  
 tatatatata tatttatata tacatatata aatatatatt tatatttata tataaatata 120  
 tataaatata tataaatata tatttatata tacatatata aatatatattg ttcatataaa 180  
 tatatatgta tatatacata tataaatata tatttatatat gtatatatat aatataatat 240  
 ataataataa tataatatat attatataaa tataatatat tatatataat atatataata 300  
 tataatatat aatatataat atataatata tatttatatat tatataatat ataaaatata 360  
 tatttatataa tatatataca taatatatat aaataaatat atataaagat ataaa 415

<210> 110

<211> 330

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(330)

<223> MAR of chromosome 2 genomic contig; 16370976..16371305

<400> 110

catttacata tgtatgtata agtatgtata ttacatactt atacatacat acttataaat 60  
 atataagtat aatacatata tacttataaa tatataagta taatacatat acacttatac 120  
 atatataagt ataatacata cataacttata catatataag tataatacat acatacttat 180  
 acatatataa gtataatata tacatactta tacatataag tataatacat acatacttat 240

## SEL PCT 012.ST25

acatatataa gtataataca tacataactta tacatatata agtataatac ataacttatta 300

catatgtata taagtatatt acataacttat 330

<210> 111

<211> 702

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(702)

<223> MAR of chromosome 2 genomic contig; 626641..627342

<400> 111

tatatataca catatacata tataatatat atacatatatac atatataatta tatatacata 60

tatattacat atatcatata tacatatata ttatatatac atatataatta tatatatcat 120

atatacatat atatattata tattatatat atcatatata catatatatt atatataatta 180

tatatatcat atatacatat atattatata tattatatat acatatatat tatatatatc 240

atataaacat atatattata tatatcatat atacatatat attatatata ttatatatat 300

catatatata tatatattat atatatcata tataatatat attatatata ttatatataa 360

tatatattat atatacatat atattatata tacatatata ttatatatac atatataatta 420

tatatacata tatattatat atacatatat attatatata tacatatata ttatatatac 480

atatataatta tatatacata tattatatat acatatatat tatatatata tatattatat 540

atatacatat atattatata tacatatatt atatatatatac atatataatta tatatacata 600

tattatatat atacatatat attatatata catatatatt atatacatat atattatata 660

tacatatata ttttatatat atataatata tttttatat at 702

## SEL PCT 012.ST25

&lt;210&gt; 112

&lt;211&gt; 679

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(679)

&lt;223&gt; MAR of chromosome 2 genomic contig; 3196047..3196725

&lt;400&gt; 112

atatattata tattcatata tcataaatat atatattata tattcatata ttatatatct 60

atatatttat atattcatat attatatatc tatatattta tatattcata tatttatatat 120

ctatttatat attcatatat tatatatcta tatattttat atattcgtat attatatatc 180

tatatattat atattcgtat attatatatc tatatattat gtattcatat atatctatat 240

attatatata ttcatatata ttataaatta tattcatata gtatatatct attataaatg 300

tatattcata tagtatatat ctatatatta taaatataca tatattatat atttatatat 360

tatatattca tatagatcta tatattatat atattcatat atgaatatat atatttatg 420

tatatatatt ataaatatat ttatatagta tagatattat atagtatatg catatttata 480

ttataaataa ttacatagt atatgtatat ttataaatta tatatattta catattacat 540

gtatatttat atattataaa tacatattta catattataa atatatttat atattatgaa 600

tataatttat atattattac atatttcat atatgcatag ttatatatta taaatatgca 660

tttatgtaaa tatatattt 679

&lt;210&gt; 113

&lt;211&gt; 728

## SEL PCT 012.ST25

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(728)

&lt;223&gt; MAR of chromosome 2 genomic contig; 3196778..3197505

&lt;400&gt; 113

tacataaata tatatttaca atagttaa atctgatatg taaatatgta ttataatat	60
ataaatatac atataatatg taaatatata aatatacata tactatgtaa atatatgtta	120
tatatacata tactatataa atatagaata tataaatata catatactat ataaatatgt	180
aatatataaa tatatactat ataaatatac atatactata taaatgtatt tataatatat	240
aaatatacat atactatata aattcatata tgaatatata atatataaat atatataata	300
tatgaatata tactcatata taaatatata tgaatatata ttataatat atagatataa	360
tatgaatata tatttataat atatagatat atatttatg aatatatatt tataatatat	420
agatatatac catatgaata tatattatac actatatgaa tatatattta taatatataa	480
atagatatat actatatgaa tatataatat atatactcta tgaatatata atatatatac	540
tatatgaata tattatatac tgtatgaata tataatatat agatgtatac tatatgaata	600
tataatatat agatatatat actatatgaa tatatataat atatagatat atactatatg	660
aatatatatg atatatagat atatactata tgaatatata atatatagat atatatttat	720
gatatatg	728

&lt;210&gt; 114

&lt;211&gt; 413

&lt;212&gt; DNA

## SEL PCT 012.ST25

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(413)

<223> MAR of chromosome 2 genomic contig; 2560638..2561050

<400> 114

atataaatat atatttatat attttatata aatatatata tttatatatt tttatataaa 60

tatatatatt tatatatatt tatataaata tatatatatta tatatatatta tataaatata 120

taaatatata tatttatata aatatataaa atatataaat atatttatat aaatatataa 180

aatatataaa tatatttata taaatatata aaatatataa atatatttat atataaatat 240

ataaaatata taaatatctt tatatataaa tatataaaat atataaatat ctttatatat 300

aaatatataa aatatataaa tatatttata tataaatata taaaatatat aaatatattt 360

atatacaaat atataaaata tataaatata tttatatata aatatataaa ata 413

<210> 115

<211> 361

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(361)

<223> MAR of chromosome 2 genomic contig; 4965309..4965669

## SEL PCT 012.ST25

&lt;400&gt; 115

tatacgtata tatacatata tatacgtata tatatacata tatatacgtata tatatacata 60

tgtatatatg tgtgtacatg tatatatata catatgtaca tatatatgta cacatatata 120

tatacatata tatgtacaca tatacatata tatgtacaca tatacatata catatatatg 180

tacacatata tatacatata tatgtacaca catatatata catatatatg tacacacata 240

tatacgtata tatgtacaca catatatatg tatatatatg tacacacata tatacgtata 300

tatatgtaca cacatatata tacgtatata tatgtacaca tatatatata cgtatatata 360

t 361

&lt;210&gt; 116

&lt;211&gt; 325

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(325)

&lt;223&gt; MAR of chromosome 2 genomic contig; 5258150..5258474

&lt;400&gt; 116

tacacacaca tatacatata tacatatata cgtgtatacgtata tatacgtata tacgtatata 60

tacatatatg tatacgtata cgtatatatg tatatatata tatatgtata cgtatacgtata 120

tatacgtata tatacatata tgtatacgtata tacgtatata cgtatatata catatatgta 180

tacgtatacgtata tatatacgtata tatatacata catatgtata cgtatacgtata tatatgtata 240

tatacgtata tgtatacgtata tacatatata cgtatatata cgtatatgta tatgtatata 300

cgtatatgta tatatgtata tatac 325

## SEL PCT 012.ST25

&lt;210&gt; 117

&lt;211&gt; 1508

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(1508)

&lt;223&gt; MAR of chromosome 2 genomic contig; 6057499..6059006

&lt;400&gt; 117

atataatata tataaattat ataatatata aaaattaata tataatatat ataaattata 60  
 taatatataa attaattata taatatatat aaattatata atatatataat taattatata 120  
 atatatataa attatataat acatatataat taattatata atatatataat tatataatat 180  
 atacaaatta tatactatat taattatata ttatataatt aattatataa tatatatataa 240  
 ttatatatta ttaaattaat tatataatat ataaattata taatatataa attaattata 300  
 taatatataa attatataat atataaatta attatataat atataaatta tataatatat 360  
 aaattaattg tataatatat aaattaatta tataatatat aatatataat taataaataa 420  
 ttatatatta attatataat taataaataa ataataaata tatataatta atatatataa 480  
 tacatcatat atatcacata tagattatat aatagttata tattatataa taaattatat 540  
 ataatatata ataaacatat ataacatatg ttatatatta cataatatag tataatatat 600  
 aacatatggt atatattaca taatatagta taatatataa catgttatat attacataat 660  
 atagtataat atataacata tgttatatat tacataatat agtataatat ataacatatg 720  
 ttatatatta cataatatag tataatatat aacatatggt atatattaca taatatagta 780  
 taatatataa catatgttat atattacata atatagtata atatatataa tatgttatat 840

## SEL PCT 012.ST25

attacataat atagtataat atataacata tgttatatat tacataatat agtataatat 900  
 ataacatatg ttatatatta cataatatag tataatatat aacatatgtt atatattaca 960  
 taatatagta taatatataa catatgttat atattacata atatagtata atatataaca 1020  
 tatgttatat attacataat atagtataat atataacata tgttatatat tacataatat 1080  
 agtataatat ataacatatg ttatatatta cataatatag tataatatat aacatatgtt 1140  
 atatattaca taatatagta taatatataa catgttatat attacataat atagtataat 1200  
 atataacata tgctatatat tacataatat agtataatat atatgttata tattacataa 1260  
 tatagtataa tatataacat atgttatata ttacatatta tagtataata tatatgttat 1320  
 atattatata atatagtata atatataatg tatgttatat attatataat atagtataat 1380  
 atataacatg ttatatatta tataatatag tataatatat atgttatata ttatataata 1440  
 tagtataata tataatatat gttatatatt atataatata gtataatata tatgttatat 1500  
 attatata 1508

<210> 118

<211> 415

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(415)

<223> MAR of chromosome 2 genomic contig; 7996866..7997280

<400> 118

caattatata atatacatat tatataattg tataaattat acaatcatat aattatatta 60  
 tatataatat acatataata taattatata taattatata atttataat ataattatat 120

## SEL PCT 012.ST25

ataattatat aattatatat aatatatatt ataattatat atataatata tatattatat 180  
 atattatata taatatataa ataatatata taatatatat ataattatat ataataatat 240  
 atgtaatat tataatatat atataatata ttatttataa ttatatatta tatatatatt 300  
 ataatatata taattataaa taatatatat tataatatat ataataatat atatataatt 360  
 atatataata atatatatta taattatata taataatata tataatttat ataatt 415

<210> 119

<211> 526

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(526)

<223> MAR of chromosome 2 genomic contig; 8300930..8301455

<400> 119

tatatcatat gatataattat acaatatatc atataaatatg atatattata tgatatattg 60  
 facaatatat catatgatat atgatataatt atacaatatata tcatataagg tatatattat 120  
 atcatatata atatataata taatatatga tataatatat gatatatgat atataatata 180  
 tgatatatga tatatgatat ataatatatg atatatgata tatgatatat aatatatgat 240  
 atatgatata tgatatataa tatatgatat atgatatatg atatgatata tgatatatga 300  
 tataatatat gatataatat atgatatatata ttatatgata tataatatat gatataattt 360  
 atatgatata taatatatga tatataatat ataatatatg atatgatata tattatatca 420  
 tatataatat ataataatat atatgatata tattatatat tttatacat tatatatata 480  
 aactatataa caatataaca tattatgtgt ataatatata ttacat 526

## SEL PCT 012.ST25

&lt;210&gt; 120

&lt;211&gt; 402

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(402)

&lt;223&gt; MAR of chromosome 2 genomic contig; 8576553..8576954

&lt;400&gt; 120

```

atgtatatta tatacaatat agtatatcat atatagtata tattatatag taatgtatta    60
tatataatgt ataatgtata aatatataat atatactaca tactatacta ttatatatac    120
tatatattat atatgataca tatactatat aatagtctat atattatact atataaatatg    180
ctatatatta tactatataa tatgctatat attatactat ataatatgct atatattata    240
ctatataata tgctatatat tatactatat aatatactat ataatatgct atatattata    300
ctatataata tactatatat tatactatat aatatactat ataacatact atatattata    360
tatgatacat atactatatt acatatataa tatatatata ta                        402

```

&lt;210&gt; 121

&lt;211&gt; 477

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

## SEL PCT 012.ST25

&lt;222&gt; (1)..(477)

&lt;223&gt; MAR of chromosome 2 genomic contig; 8785649..8786125

&lt;400&gt; 121

tatttatata tatatttata tatatatatta tatatatatta tatatatatt tatatatata 60

tttatatata tatttatata tttatatata tatattttta tatatttata tatatatatta 120

tatatttata tatatttata tttatatata tatttatata tatttatata tatttatata 180

tatatatatta tatatatatta tatatatata tttatatata tttatatata tttatatata 240

tatttatata tatatttata tatatatattca tatatatatta tatatatatt catatatatt 300

tatatatata ttcatatata tttatatata tatttatata tatatttata tatatttata 360

tatatttata tatatatatta tatatatatt tatatatata tatttatata tatatttata 420

tatatatatt tatatatata tttatatata tatatttata tatatatatta tatatat 477

&lt;210&gt; 122

&lt;211&gt; 773

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(773)

&lt;223&gt; MAR of chromosome 2 genomic contig; 10064737..10065509

&lt;400&gt; 122

atattatata tattacatat atatttatatt gtatataata tatatatatt attgtatata 60

atatatatat tatattgtat ataatatata tatttatattg tatataatat atatattata 120

## SEL PCT 012.ST25

ttgtatataa tatattatat tgtatatatt atattgtata tattatattg tataacaatat 180  
 atattatatt gtatacaata tatattatat tgtatataat atattatatt gtatataata 240  
 tattatattg tatatattat attgtatata atatattata ttgtatataa tatattatat 300  
 tgtatatatt atattgtata taatatatta tatgtatata ataatgtgta tactatatta 360  
 tataatatat attatataca atatataata tattgtatat catatatgat atattgtata 420  
 taatatataa tatatgatat attgtatata atatattata tatgatatat tgtatattat 480  
 atattatata tgatatattg tatattatat attatattat gtatatattgta tattatatat 540  
 tatatattgt atataaatg ttatatattg tatataatat gttatatatt atatattgta 600  
 tatatgttat atattatgta ttgtatataa tatgttatat attatattat gtatataatg 660  
 tattatatat tatatatatt atatattgta tataatgtat tatatattgt atattatata 720  
 ttatatattg tatataatat attatataca ttatatata tattatatat tgt 773

<210> 123

<211> 1554

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(1554)

<223> MAR of chromosome 2 genomic contig; 1039775..1041328

<400> 123

ataatatatt aaatgtatat ataatatatt aaatataaat atatttataa tatataaata 60  
 ttatataaaa tataaaatat atattaaata taaatatata taaaatatat attaaatatata 120  
 taaaatataa atatatatta aatatatatt aaatatataa aatataaata tatattaaat 180

## SEL PCT 012.ST25

atattttaa tatataaaat ataaatat ataaatata tttaaatat ataaatata	240
aatatatatt aaatatatt taaatatatt aatatataat acatatatta aatatatt	300
atatatataa aatatataaa atataaatat atattaaata tatataaaat atatatgta	360
aatatataaa agatatataa aatatataa tatattaaat atatatataaa tatatatata	420
ttaaatatat atattaaata taaatatata taaaatataa atatatgtat taaatatata	480
tattaaatat aaatatatgt attaaatata tattaaatat gaatatatgt attaaatata	540
tattaaatat aaatatatgt attatatata tagaatataa atatatgtat taaatatagt	600
atattaaata taaatatata taaaatatat attaaatatg aatatatata aaatatatat	660
attaaaaata tatataatat aaatatatat aaaatatata tattaaaaat atataata	720
taaatatata taaaatatat atattaaaaa tatatatataa atatatatat taaaatatata	780
tataaaatat atatatataa aatatatata aaatatatat attaaaaata tatattaaat	840
ataaatatat atattaaaaa tatatatata atataactat atattaaata tatattaaat	900
ataactatat attaaatata tattaaatat aactatatat taaatatata ttaaatataa	960
ctatatatta aatatatt aaatataact atatatataa tatatatata atataactat	1020
atattaaata tatattaaat ataactatat attaaatata tattaaatat aactatatat	1080
taaatatata tgaaatataa ctatatatta aatatatt aaatataact atagtatta	1140
aatataaata tatgtcttaa atatatatta aatataaata tatgtattaa atatatatta	1200
aatataaata tgtgtattaa atatatatta aatataaata tgtgtattaa atatatatta	1260
aatataaata tgtgtattaa atatatatta aatataaata tgtgtattaa atatatatat	1320
taaatatataa tatatgtatt aaatatatat taaatatataa tatatatata atatatatat	1380
taaatatataa tatatatata atataaatat atatatataa tatatatatt aaatatataa	1440
atatatataa tatatatatt aaatatataa ataaatatata aatatatt aaatatataa	1500
acatatatta aatatatgta ttatatatat atataaaata tatgtattaa atat	1554

&lt;210&gt; 124

&lt;211&gt; 650

## SEL PCT 012.ST25

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(650)

&lt;223&gt; MAR of chromosome 2 genomic contig; 3944813..3945462

&lt;400&gt; 124

catgatatat tatgtataat atatattata gattacatat aaattatata tataatatat 60  
aattatataa tatataatat tatataatat attatatata ttatacaatt atataatata 120  
tataatatac aattatataa tatataatat acaattatat aatatataat acaatataat 180  
atatatttaa tatattatat aatacatatt taatatatta tatattatat gttatatact 240  
aatatataa tatgtattta atatatacta ttatatatgt aatatattat ataatttatg 300  
taacatatta tatattatat atgcaatata ttacatgtta catatatatt acatataata 360  
tatgtaatat ataatataca ctatattatt atagtatata atatactata ttatgtaatt 420  
atataatata gtatattata cactatatta tattatcata taattatata ttatatacta 480  
tattacatat atattatgta atataatatg caatatgtta catatataat atatatgtat 540  
tatatagtat atatactata gtatatataa aatatatgct ataatatata ttttatatat 600  
tatataatac atataatgta tcatatatta tatataatat attttataat 650

&lt;210&gt; 125

&lt;211&gt; 441

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

## SEL PCT 012.ST25

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(441)

&lt;223&gt; MAR of chromosome 2 genomic contig; 5314265..5314705

&lt;400&gt; 125

tataaatata tatgaaatat atataaatta tatataattt atatatacat atataaatta 60

tatataaatt atatataaat tatatatata tatataaatt atatattata tataaaaattg 120

tatatattta tatataaatt gtatatataa ttatatata aattgtatat ataatttata 180

tatacaatgt atatattaat ttatatatac attgtatata taatttatat atacattgta 240

tatacaattt atatatacat tgtatatata atttatatat acattgtata tacaatttat 300

atataaatta tattatttat atatagtata tataaatata tatactatat ataaattata 360

tatttattta tatatttat ttttatata taaattatat attatttata tatacattat 420

atataaatta tatattatt a 441

&lt;210&gt; 126

&lt;211&gt; 1169

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(1169)

&lt;223&gt; MAR of chromosome 2 genomic contig; 5953971..5955139

&lt;400&gt; 126

## SEL PCT 012.ST25

atgtattcat attatatatt tatatatataa taatatatcat tcatattata ttttatata	60
ttaaataat atattcatat tatatatatta tatataaata tataatatat ttatgtataa	120
ataatatata tattcatatt atatatttct atataaataa tatatatatt catatttat	180
atttatatat aaatatataa tatatttata tataaatata taatatattt atatataata	240
tatatattca tatttatatat ttatatataa atatataata ttttatata taaataat	300
atatattcat attatatatt tatatatataa taatatatat tcatattata ttttatata	360
taaataat atattcatat tatatactta tatataaata atatatttc atatttatata	420
cttatatata aatatatat attcatatta tatatttata taaaaaat atatattcat	480
attatatatt tatatataat atatatttc atatttatata ttatatatt ctatatattc	540
atatttatata ttatatata aataatgtat attcatatta tatatttata tataaataat	600
gtatatcat attatatatt tatatatataa tatatatcca tatttatatat ttatatataa	660
atatatttc atatttatata ttatatata aatatatatt catatttat atttatataa	720
aatatatata ttcatattat atttatatat aaatatatat attcatatat atatttatat	780
ataatatata tattcatatt atatatttat atataatata tatattcata ttatatattt	840
atatataaat aatatatata ttcatattat atatttatat ataaataatg tatattcata	900
ttatatattt atatataaat aatgtatatt catatttat atttatatat aaatatatat	960
attcatatta tatatttgta tataaatata tattcatatt atatatttgt atatatttc	1020
atatatattt atatataaat atataatatt catatttat ataaatatat atattcatat	1080
tatatattta tatatatataa taatatatat tcatattatt tatatatata aatatatat	1140
attcatatta ttatatata taaataata	1169

&lt;210&gt; 127

&lt;211&gt; 653

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

## SEL PCT 012.ST25

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(653)

&lt;223&gt; MAR of chromosome 2 genomic contig; 6427669..6428321

&lt;400&gt; 127

tatatatgta tacatatatg tatatatgtg tatatatgta tacatatatg tatatatgtg 60  
tatatatgta tacatatatg tatatatgtg tatatatgta tacatatatg tatatatgtg 120  
tatatatgta tacatatatg tatatatgtg tatatatgta tacatatatg tatatatgtg 180  
tatatatgta tacatatatg tatatatgtg tatatatgta tacatatatg tatacatgtg 240  
tacatgtgta tacatatatg tatacatgtg tacatgtgta tacatatatg tatacatgtg 300  
tacatgtgta tacatatatg tatatatgtg tatacatata tgtatatatg tgtatatatg 360  
tatacatata tgtatataag tgtatatatg tgtatatgta tataagtgtg tatatgtgta 420  
tatgtatata agtgtatata tgtgtatatg tatataagtg tatatatgtg tatatatgta 480  
tacatatatg tatatatgtg tatatatgtg tatatgtata taagtgtata tatgtgtata 540  
tatgtataca tatatatgtg tatatatgta tacatatatg tatatatgtg tatatatgta 600  
tacatatatg taaatatgtg tatatatgtg tatatgtata taagtgtata tat 653

&lt;210&gt; 128

&lt;211&gt; 414

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(414)

## SEL PCT 012.ST25

<223> MAR of chromosome 2 genomic contig; 10890453..10890866

<400> 128

tatattttgt aaatatatat atagtaaata tatgtaaata tatatatttt gtaaatatat	60
atatattttg taaatatatg taaatatata tttttgtaa atatatgtaa atatatatat	120
ttgtaaata tatgtaaata tatatatttt gtaaatatat gtaaatatat atattttgta	180
aatatatgta aatatatata tttgtaaat ttatgtaaat atatatattt tgtaaataata	240
tgtaaataata tatatatttt gtaaatatat atacatatat attttgtaa tatataaaca	300
tatatatttt ataaatatat ttataaatat atatatgtta aatatattta taaatatatt	360
tataatatat atattgtaaa tatgtttata aatatatata ttgtatatat aaat	414

<210> 129

<211> 496

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(496)

<223> MAR of chromosome 2 genomic contig; 13952568..13953063

<400> 129

taatatatcat attatatatt atatatgtta tatataatat acatatattata tattatatat	60
tgatatatata atatacatat tatatattat atattgtata tataatataac atattatata	120
ttatatattg tatatataat atacatatta tatattatat attgtatata taatatatcat	180
attatatatt atatatgtta tatataatat acatatattata tattatatat tgatatatata	240

## SEL PCT 012.ST25

atatacatat tatatattat atattgtata tataatatac atattatata ttatatattg 300  
 tatatataat atacatatta tatattatat attgtatata taatatacat attatatatt 360  
 atatatgtta tatataatat acatattata tattatatat tgtatatata atatacatat 420  
 tatatattat atattgtata tataatatac atattatata ttatatattg tatatataat 480  
 atacatatta tatatt 496

<210> 130

<211> 317

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(317)

<223> MAR of chromosome 2 genomic contig; 16942865..16943181

<400> 130

tctcctagta gttatatata tatatatgtg tatatatata tctcctagta gatatatata 60  
 tatatatatc ctagtagata tatatatata tatatcctag tagatatata tatatatata 120  
 tcctagtagt tatatatata tatatatacct aacagttata tatatatata tcctagtagt 180  
 tatatatata tatatcctag tagttatata tatatatata tcctagtagt tatatatata 240  
 tatatcctag tagttatata tatatatatc ctagtagtta tatatatata ttatatatta 300  
 tataatatat atataat 317

<210> 131

<211> 464

<212> DNA

## SEL PCT 012.ST25

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(464)

&lt;223&gt; MAR of chromosome 2 genomic contig; 17217049..17217512

&lt;400&gt; 131

```

acatactata tatatacaca tactatatat actatatata gtatatagta tacatatact   60
atacatatac atatactata catatacata tacatatact aagtatacgt atatacagta   120
catagtatat gtatactata tagtatgtat atatagcata tagtatgcgt atactctata   180
tagcatatag tatgcatata cgctatatag catatagtat gcatatacta tatatagtat   240
agagtatgcg tatactatat atatagtata gagtatgcgt atactatata tatagtatag   300
agtatgcgta tactatatat atagtataga gtatgcgtat actatatata tagtatagag   360
tatgcgtata ctatatatat agtatagagt atgcgtatac tatatatata gtatagagta   420
tgcgtatact atatatatag tatagagtat gtatatatat agta                     464

```

&lt;210&gt; 132

&lt;211&gt; 430

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(430)

&lt;223&gt; MAR of chromosome 2 genomic contig; 19647266..19647695

## SEL PCT 012.ST25

<400> 132  
 tgtaaatata tgtaaatata ttttatatt atatattata taaaaatata atatataata 60  
 tataatatat aaactatata ttaatataat atatataaac tattatataa atacatatta 120  
 aatatattat attttaata tttatatatt aaatataata tatatttaatt atttatatat 180  
 taaatatata atatatttaa ttttatata atatatagca tttttatat ttatattata 240  
 tataacattt tatatttata ttttatatta tatatattta atttatattt atatttatatt 300  
 tatatttata ttatatataa cataattata tatattttca tattgtatat aataaagaaa 360  
 tgtatatttg ttatatataa tatatattat ataatttatt atatattata taatatatat 420  
 tatataatat 430

<210> 133

<211> 2131

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(2131)

<223> MAR of chromosome 2 genomic contig; 20481223..20483353

<400> 133  
 tatatataaa tatatttata tttatatat atttatataa atatattttt atataaatat 60  
 atatttaata taaatatctt tatatttaatt atatatttaa tataaatatc ttttatatta 120  
 atatatattt atatataaat atatatttat atttaataata tattaatatt taatatacgt 180  
 ttatatttaa tatatatttc tatataaata ttttatattt aacatatatt tatatataaa 240

## SEL PCT 012.ST25

tatatttata tttaatatat ttacatataa atatatttat atgtaatat tttacatata	300
aatatattta tatttaatat atatgcatat gtaaatatat ttatatftaa taatatttat	360
atataaatat atttatattt aataatattt atatataaat atatttatat ttaatatata	420
ttaaatatat atttatattt aatatatat ttatatttaaat atatatttat atttaatat	480
tatttatatat aaacatatat ttatatftaa tatatatttat atataaacat atatttatat	540
ttaatatata ttatatataa acatatattt atatttaata tatattftata tttaatatat	600
tatatataaa catatatfta tatttaatat atatttatat taaatatata ttatatataa	660
acatatattt atatttaata tatattftata ttaaatatat atttatattt aatatatata	720
tattaaatat atatttatat ttaatatata ttatatftaa atatatattt atattaaata	780
tatttatatt taatatatat ttatatftaaa tatatattftaa atatttaata tatattftata	840
tttaatatat acatatatat ttatatftaa tatatacata tatattftata tttaatatat	900
acatatatat ttatatftaa tatatacata tatattftata tttaatatat aaatttatat	960
tttatatata taaaaatatat tatttatatt taatatatat aaatatatat ttatatftaa	1020
tatatatatt tatattgaat atatacataa atatatattt atatttaata tataaacata	1080
tatttatatt tatatattftaa atatatattt atatttaata tataaatata tatttatatt	1140
taatatattt atatatacta atatatftat atttaatat tttatatata gatatatfta	1200
tatttaatat atttatgtgt attaatatat ttatatftaa tatattftata tattaatat	1260
tttatattt atatttatat attaatatat ttatatfta tatttatatt ttatatattt	1320
atatattaat atatttatat ttatatatat ttitatatat taataaattt atattftata	1380
tatttatata ttaataaatt tatattftat acagttatat aaatatattt atattftata	1440
cagttatat aatatattfta tattttatag ttatataaat atatttatat ttatacagt	1500
tatataaata tatttatatt ttatacagtt atataaatat atttatattt tatacagtta	1560
tataaatata ttatatattt atacagttat ataaatatat ttatatfta tacagttata	1620
taaatatatt tatattftat acagttatat aaatatattt atattftata cagttatat	1680
aatatatfta tattttatag agttatatata atatatftat attttatata gttatatata	1740

## SEL PCT 012.ST25

tatatttatg tttatacat ttatataaat atatttatat ttatacatt tgtatttaat 1800  
 atatatttat atataaatat attttatatt taatatattt atatataaat atatattgat 1860  
 atttaatatata tatttatata taaatatata ttgatattta atatgtttat atataaatat 1920  
 atatttatat ttaatatata tgtttatata tcaatatata ttatattta atatatttt 1980  
 acatataaat atatatttat attgatata tatttatatt tgatatatat ttatatata 2040  
 ttaatatatt tacatttgat atatatattta tatatattaa tatatttaca ttgatatat 2100  
 attttatata tattaatat ttacatttg a 2131

&lt;210&gt; 134

&lt;211&gt; 842

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(842)

&lt;223&gt; MAR of chromosome 2 genomic contig; 20483478..20484319

&lt;400&gt; 134

tatatattta tgtttaatat atatttatag ataaatatat atttacgttt aatatatatt 60  
 tatagataaa tatatattta cgtttaatat atatttatct ataaatatat ttacgtttaa 120  
 tatatattta tatattaata tatttatggt taatatatat ttatatatat taatatattt 180  
 atgtttaata tatatttata tattaatat ttatgttta atatatttat atatattaat 240  
 atatttatgt ttaatatata ttatatggt aatatattta ggtatatata tatttatatg 300  
 ttaatatata ttatatttaa tatatttat ttatatataa aagtatatat aatatataaa 360  
 tatttatataa attatttat agtattttta tatatattta tatataaatt ttatatattt 420

## SEL PCT 012.ST25

tatatatata aatatatatt tatatatata ttttatatat aatatatat ttatatata 480  
 attatatata taaatatata ttttatatt ttatatataa atatatatat ttatatata 540  
 attttatata ttttatatat gtaaatat atataaattt tatatatgt atatatatt 600  
 ataaatttta tatatatatt tatatatata atatatataa tatatatata ttttatatat 660  
 attatatata ttttatatt atattata ttttttta tatatttta tatgtatat 720  
 atatttatat ttatatatt ttttttta tatatttat atatatatt atatatgtat 780  
 attatatata ttatatatta tataatatat tatatatatt atattatata tttatatat 840  
 at 842

<210> 135

<211> 645

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(645)

<223> MAR of chromosome 2 genomic contig; 20897566..20898210

<400> 135

gtatattat attatatatt atataatata tattatatat taataaatta tatataatat 60  
 aatatatatg tatatttata ttatgttat aatatacata taattatata tgtatgtata 120  
 catgtatata tatacgtata tgtgtatatg tatacatata ggtatatgtg tacatgtata 180  
 catataggta tatgtatatg tatacatgta tacatataat ataattacat atgtatgtat 240  
 acatacatat gtaattatat tatatatgta tatgtatatt tatataatat ataatatgta 300  
 ttatatatta tacatgcata tttatatgta tattatatat acacatatata tataattata 360

## SEL PCT 012.ST25

tatgtatgta tatatacaca tatatatatta tattatatat gtatattata tacatatatt 420  
 tatattatat atgtatatat atttatcata ttatatgta atatgcatgt gtaataaata 480  
 atatacacat ttatatatgt atattatata catatatatta tattgtatat gtatatatat 540  
 ttatatatat ttgtatatca tatatttata tattgtatat ttatgtatat tatatatatta 600  
 tatattatat atgtattata taatatatat gtaaatatat attat 645

&lt;210&gt; 136

&lt;211&gt; 722

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(722)

&lt;223&gt; MAR of chromosome 2 genomic contig; 21664541..21665262

&lt;400&gt; 136

tataatatat attatatattct atataaatatg taaaatatat attatatattct atataatgta 60  
 ttatatatag aatataaatat attctatgta ttctataatc tatataaatac atattatata 120  
 ttatatagaa tattataaat aatatattct atattatata tagaatatat tctatatgtt 180  
 tatattctat atattatata tgaaatagta tataaaatat atataatata tataaaatat 240  
 gatataaat atataaaaa taatatataa tgtataaat ataaaaaat atataatgta 300  
 taatatataa aataatatat aatgtataat atataaaata atataaatg tatattatat 360  
 aaaataatat ataatgtata ttatatataa aataatatat aatgtatatatt atataaaaa 420  
 taatatataa tgtatataaa ataatatata atatattata tataaaataa tatataaat 480  
 attatatata aaataatata tattatatat aaaataatat ataatatatt atataaaaa 540

## SEL PCT 012.ST25

taatatatat tatatatataa ataatatata atatattata tataaaataa tatatattat 600  
 atataaaata atatattatta tatataaaat aatataatat atattatata taaaataata 660  
 tataatatat tatataaaaa tataaatata ttatataaaa atataaaata taaaatatta 720  
 ca 722

<210> 137

<211> 305

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(305)

<223> MAR of chromosome 2 genomic contig; 22834991..22835295

<400> 137

aatataaaat atatgatata taatacgtat tatatatgta taatacgtat tatatattaa 60  
 tatataatat ataatacata ttatatatgt atataatata tactaatata tataatgtat 120  
 acattatata ttacataat atataataca taatatagaa ttataattat atataataca 180  
 taatatataa ttatatatat tattatatat gtatttatat tatatataat atattatata 240  
 taatatatat tatataatta tataagtata taattatgtt atatacataa taatatataa 300  
 tatat 305

<210> 138

<211> 352

<212> DNA

<213> Homo sapiens

## SEL PCT 012.ST25

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(352)

&lt;223&gt; MAR of chromosome 2 genomic contig; 25277762..25278113

&lt;400&gt; 138

taatatatat aatatattat atattatata taatatattt tataatatat aaaatatatt 60

atatataata tataatatat ttataatat atataatata ttatatataa tatataatat 120

atattataat atatataata tattatatat attatatatt tatatttatt tatatatcca 180

taaatatata ttatatata atatatttta taatatatta tatataatat ataatatatt 240

ttataatata ttataatata taatatataa tatattttat aatatatata atatataata 300

tattatatat ttatatttat ttatatattc ataaatatat atatttatat ta 352

&lt;210&gt; 139

&lt;211&gt; 342

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(342)

&lt;223&gt; MAR of chromosome 2 genomic contig; 25378452..25378793

&lt;400&gt; 139

tatgtacata tatattttat atattatata taatatatat tatatgatat atataatata 60

## SEL PCT 012.ST25

ttatataata taatatataa aatatatata atatatatta tattatataa attatattat 120  
 atatatcata taatatattt tatatattat ataatatata ttatattata tatattttat 180  
 atattatatt atatatata tatatcatat aatatatatt atattatata ttttatatat 240  
 tatataaat atattatata tttttatata ttatataata tatattatat attttatata 300  
 ttatataata catatattat atataatata atatatatta ta 342

<210> 140

<211> 663

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(663)

<223> MAR of chromosome 2 genomic contig; 30209437..30210099

<400> 140

aatatatatt acatattgta tatatagtat atgtaatgta tataatatag tatattctat 60  
 attgtataat agtaatatat agtatatgat atactatata ttacttatca tatatacaat 120  
 atatatata tcgtatattg tatattatat attgtatata tgtaatatat gatatgtaca 180  
 tatgttatat atgtatataa tatactatat tatatatattg atattatata catatataac 240  
 actatttatac aatatataat atagcatatt atatacaata tagcatatac aatatataat 300  
 atagcatatt atataataa tagtatatta tatacaatat ataatatagc atattatata 360  
 taatataata tagtatatta tatacaatat ataatatagc atatacaata tagtatataa 420  
 tatataaat agcatatata atatagtata ttatatataa tatataaat agcatgtaca 480  
 atatagtatg ttatatataa tatataaat agcatatata atatagtata ttatatataa 540

## SEL PCT 012.ST25

tatataatat agcatataca atatattata ttatatacaa tatataatat agcatataca 600  
 atatagtata ttatatacaa tatataatac agcatataca atatagtata ttacatacag 660  
 tat 663

<210> 141

<211> 1200

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(1200)

<223> MAR of chromosome 2 genomic contig; 31725089..31726288

<400> 141

tgtacttata tattataatg tatatataaa gtatatactt tatatatact tatatattat 60  
 aatgtatatt attgtatata agtatataic ataatatata cttacatatg ctcacatata 120  
 ttataatgta tattgtatat attatataca tattatatat gtataatgta tatatacatt 180  
 atatatgtat aatgtatata tacattatat atgtataatg tatatataca ttatatatgt 240  
 ataatgtata taatatatac aatatatgta taatatataa tatatacaat atatgtataa 300  
 tatacaatat atgtataata tacaatatat gtatagtata taatatatat tatatatgta 360  
 tagtatatta tatattatat atgtatagta tataatatgt ataatgtata tattataata 420  
 tattatatat aatatctata acaatataat atattgtata tattatatat aatatatatt 480  
 tatataatat atattatata taatatatta tgtattttatt tatattatat ataatatataa 540  
 tatatataat ataaataata ttattatat attaatataa atatttatat taatatatat 600  
 ttattatata taaataatat ctatgatata aataatatat aatatacatg tatatgttat 660

## SEL PCT 012.ST25

aatatataca tataatatac atgtgtatat atactataca tgtatatata acatgtatat 720  
atatacatgt atatatatta tgtatacatg tatagtatat atacatgtat atatatat 780  
atatactata catgtatata tacatgtata tatatacata tatactatac atgtatatat 840  
acatgtatat atacacatat atactataca tgtatatata catgtatata tatacatgta 900  
tggtatatac attattataa tatacatata tagtatacat tatatacatt atataatatg 960  
cattattata atataatata cattattata atatacatta ttataatata atatacatta 1020  
ttataatata cattattata atatacatta taataatata cattattata atatacatta 1080  
taatattgaa gtatatatac tataatatat gtatatatta taatgtatat aatatacatt 1140  
attatatata agtatgtatt atatataagt atatattata atatatgtat atacatatat 1200

&lt;210&gt; 142

&lt;211&gt; 325

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(325)

&lt;223&gt; MAR of chromosome 2 genomic contig; 32147252..32147576

&lt;400&gt; 142

aaatacaaat atttatttat atataatata taatataata tatttattta tatataatat 60  
ataatttata attatataaa tatataatat atttatatat aatatataat ttatttatat 120  
attaattata tatataataa atatatataa tatataattt tatttatatat taattatata 180  
tataataaat atatatataa tataataata ttatatacat tatatataaa tataaatatt 240  
tatataatat ataataaat atatttattt atatatataa atataatata taattatata 300

SEL PCT 012.ST25  
325

aatatataat atatttatat ataac

<210> 143

<211> 507

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(507)

<223> MAR of chromosome 2 genomic contig; 32312662..32313168

<400> 143

attatttata taaatattat atttatatta ttatataaa tattatattt atattattta 60

tataaatatt atatttatat tatttatata aatatttat ttatattatt tatataaata 120

ttatatttat attatttata taaatattat atttatatta ttatataaa tattatattt 180

atattattta tataaatatt atatttatat tatttatata aatatttat ttatattatt 240

tatataaata ttatatttat attatttata taaatattta ttatattat ttatataaat 300

attatattta tattatttat ataaatattt atttatatta ttatataaa tattatttta 360

tatttatata aataatatat aaataaatat ttatatgta tataaatatt atttatatta 420

ttattttaa taaataatat aaattaatat aaatattaat attatttatt ttattataaa 480

taataataat attatattta tatttat 507

<210> 144

<211> 339

<212> DNA

<213> Homo sapiens

## SEL PCT 012.ST25

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(339)

&lt;223&gt; MAR of chromosome 2 genomic contig; 33651118..33651456

&lt;400&gt; 144

aaatataata tattatttat atataatata aatgatatat tatgtatata taaaatataa 60

ataatatatt atgtatatataaaaatataaaa tattatttat atataaaata taaataatat 120

ttatatataa aatataaata ttatattatt tatatatataa atataaataa tatattattt 180

atatataaaa tataaataat atattattta tatataaata atatataaaa tataaatata 240

tattatatat aaataaaaata tatatattat atatataaat ttatatataa tatataaaat 300

ataatatata tatttaatat ttattatata atatataat 339

&lt;210&gt; 145

&lt;211&gt; 461

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(461)

&lt;223&gt; MAR of chromosome 2 genomic contig; 45073053..45073513

&lt;400&gt; 145

tgtgtatata tatatacgtg tacatatata tatatacatg tgtatatata tacgtgtaca 60

## SEL PCT 012.ST25

tatacatata tacatgtgta tatataatgta catatacata tatacatgtg tatacatata 120  
 tatatacatg tacatatata tatatacatg tgtatacata catatatata tgtacatata 180  
 catatatata tgtgtatact tacatatata catgtacata tacatatata catgtgtata 240  
 tatacatata tacacgtaca tatatacatata tacatgtaca tatatacatg tatacatata 300  
 tacatgtaca tatgtacata tatatacatgta tacatatata catgtacata tgtacatata 360  
 tacatgtata catatatata tgtacatatg tacatatata catgtatata tatatacata 420  
 tgtacatacg cacagataga catatatata tatgtacata c 461

&lt;210&gt; 146

&lt;211&gt; 1162

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(1162)

&lt;223&gt; MAR of chromosome 2 genomic contig; 45487691..45488852

&lt;400&gt; 146

attatattat ctatataaat ctattatatt tattatatta tctatataat atctattata 60  
 tatattatat tatctatata aatctattat atatattata ttatctatat aaatctatta 120  
 tatatattat attatctata tatctattat atattatatt atattatatt atatataata 180  
 tctattatat atattatatt atattatatt atatataata tctattatat atattatatt 240  
 atctatataa tatctattat atattatata ttatattata tataatatct attatatata 300  
 ttatattata ttatatataa tatctattat atctattata tatattatat atatctatta 360  
 tatctattat atatattata tataatatct attatatcta ttatatatat tatatataat 420

## SEL PCT 012.ST25

atctattata tctattatat tatattatat ataatatcta ttatatctat tatatatatt 480  
 atatatatct attatatcta ttatatatat tatatataat atctattata tctattatat 540  
 atattatata taatatctat tatatctatt atatattata tatataatat ctattatata 600  
 tattatatat tatatatata atatctatta tatctattat atctattata tatatatcta 660  
 ttatatctat tatatatatt atatacataa tatctattat atctattata tatattatat 720  
 atataatata tattatatct attatatata tactatctat tatatctatt atatatatta 780  
 tatatgtact atctattata tctattatat ctattatata tatactatct attatatcta 840  
 ttatatatat tatatatata ctatctatta tatctattat atatattata tatatactat 900  
 ctattatata tctattatat atattatttt atattatata tagtatctat tacatatatt 960  
 atattatatt atatataata tctattatat atattatatt atattataaa taatatatat 1020  
 aatatctgtt atatataata gatattatat ataatatata atatatataa tagatattat 1080  
 atatattata ttatataata tataatatat aatataatta atataaaata tatataatat 1140  
 ataattaata taatatgtaa ta 1162

&lt;210&gt; 147

&lt;211&gt; 562

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(562)

&lt;223&gt; MAR of chromosome 2 genomic contig; 45516233..45516794

&lt;400&gt; 147

acatattata tatattatat ataatatata ttatatatac atattatata tattatatat 60

## SEL PCT 012.ST25

aatatatatt atatatacat attatatata ttatatatac atatatatat tgtatataat 120  
atatacatat tatatatatt atatatacat attatatatt atatataata tatacatatt 180  
atatattata tataaatatt atatattata tataaatatt atatatataa atattatata 240  
ttatatataa atattatata tcttatatat aaatataata tataatatat ataattatta 300  
tatattatat ataaatatta tatattattat ataattattat atataatata taaatatata 360  
tattatataa atattgtata tattatataa atattatata tattatatat aaatattata 420  
tatattatat aaatatatat aaatatataa aatatataaa tatgtaaaat ttattttat 480  
aaatatataa tataaatata taaatatataa tataaattat atataatata taatatatta 540  
tacataatat atactatata ta 562

&lt;210&gt; 148

&lt;211&gt; 801

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(801)

&lt;223&gt; MAR of chromosome 2 genomic contig; 45727251..45728051

&lt;400&gt; 148

atatatatat ataatatata catatataga atatatatat tattatattc tatatataga 60  
atatatatat agaatatata tatatagaat atatatatag aatatatata tagaatatat 120  
atatagaata tatatataga atatatatat atagaatata tatatagaat atatatatat 180  
agaatatata tatatagaat atatatatat agaatatata tatatagaat atatatatag 240  
aatatatata tagaatatat atatatagaa tatatatata gaatatatat atatagaata 300

## SEL PCT 012.ST25

tatatataga atatatatat atagaatata tatatagaat atatatatat agaatatata 360  
 tatagaatat atatatatag aatatatata tagaatatat atatatagaa tatatatata 420  
 gaatatatat atatagaata tatatataga atatatatat atagaatata tatatagaat 480  
 atatatatat agaatatata tatagaatat atatatatag aatatatata tagaatatat 540  
 atatatagaa tatatatata gaatatatat atatagaata tatatataga atatatatat 600  
 atagaatata tatatagaat atatatatat agtatatata gaatatatat atatagtata 660  
 tatagaatat atatatatag aatatatata tagaatatat atatatagaa tatatatata 720  
 gaatatatat atatagaata tatatataga atatatatat atatagaata tatatataga 780  
 atatatatat atatatagaa t 801

&lt;210&gt; 149

&lt;211&gt; 346

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(346)

&lt;223&gt; MAR of chromosome 2 genomic contig; 50937238..50937583

&lt;400&gt; 149

taaaattata tatattatat ataatatata atatattata tataatatat attataatat 60  
 atataatat tattatataa aatatattct atagaatata tattctatta tataatatat 120  
 attctattat aatatatatt atatataata tatattctat tataatatat attatatata 180  
 atatattcta ttatgatata tattatatat aataacatat attatatata atatatattc 240  
 tattatataa aatatatatt atataaaata tatattctat tatataaaat atatattata 300

## SEL PCT 012.ST25

taaaatatat attatattat ataaaatata tattatacta tatata 346

<210> 150

<211> 462

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(462)

<223> MAR of chromosome 2 genomic contig; 55672627..55673088

<400> 150

taaatatata ttatatatta tattatatat aatatattta ttttatata tactataatt 60

tatatataat atatattata tatataatat atttataata tatatcatat aaataatata 120

tatttataat atatatcata taaataatat atatttataa tagatatcat ataaataata 180

tatatttata atagatatca tataaataat atatatttat aatagatatc atataaataa 240

tatatattta taatagatat catataaata atatatattt ataatatata tcatataaat 300

aatatatatt tataatatat atcatataaa taatatatat ttataatata tatcatataa 360

ataatatata tttataatag atatcatata aataatatat atttataata gatatcatat 420

aaataatata tatttataat agatatcata taaataatat at 462

<210> 151

<211> 401

<212> DNA

<213> Homo sapiens

## SEL PCT 012.ST25

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(401)

&lt;223&gt; MAR of chromosome 2 genomic contig; 56081352..56081752

&lt;400&gt; 151

tatacatgta tgtattcgta tatgtatggt atatatgtat atgtgttata tacatatata	60
tatatacatg tatatgtggt atatacatat acatatatac atgtatatgt gttatatata	120
tatacatata tacatgtata tgtgttatat acatatacat atatacatgt atatgtgtta	180
tatacatgtg tatgtgtata tgtatatata catatatgtg tatgtgcatg tgtatatata	240
catatatgta tatgtgtata tgtatatata catatatgta tatgtgtatg tgtatacgta	300
tatatacata tatgtgtatg tgtatgtgta tacgtatata tatacatata tgtgtatgtg	360
tatacgtaca tatacatata tgtgtatgtg tatacgtaca t	401

&lt;210&gt; 152

&lt;211&gt; 765

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(765)

&lt;223&gt; MAR of chromosome 2 genomic contig; 56404208..56404972

&lt;400&gt; 152

tatattatat aaagaatata tattatataa tatgtaaaga atatatatta tataatatgt	60
---	----

## SEL PCT 012.ST25

aaagaatata tattatatat tatgtaaaga atatatatta tatataatat ataaagaata 120  
 tatattatat aatatataaa gaatatatat tatatattat ataaagaata tatattatat 180  
 ataatatata aagaatatat aatatataat atataaagaa tatatattat atataatata 240  
 taaagaatat atattatata taatatataa agaatatata ttatatatta tataaagaat 300  
 acatatatat aatatataaa gaatatatat tatatataat atataaagaa tatatattat 360  
 atataatata taaagaatat atattatata taatatataa agaatatata ttatatataa 420  
 tatataaaga atatatatta tatataatat ataaagaata tatattatat atattatata 480  
 aagaatatta tatattatat aaagaatata tattatatat aatatataaa gaataaacat 540  
 atatactata tataaagaat atacattata tatactatat ataaagaata tacattatat 600  
 atactatata taaagaatat atataatata taaagaatat acattatata taatatataa 660  
 agaatatatt atatattata taaagaatac attataatat aaagaatata ttatatataa 720  
 tataaagaat acattataat atataaagaa tatatataat atata 765

<210> 153

<211> 443

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(443)

<223> MAR of chromosome 2 genomic contig; 61953416..61953858

<400> 153

ttatatatt atagataaaa ttatattata ttacatgtaa tatataatat gtaaaatata 60  
 ttatattaca tatataatat ataatatgta aaatatatta tattacatat ataatatata 120

## SEL PCT 012.ST25

ataigtataaa tatatttat tacatatata atataaaata ttacatatata tatattttac 180  
 ataaatatat attatctatt acatatttat tatatgtaat aatatgtaca tatgtataaa 240  
 tatgtatata ttatacata tgtatatatt atatatatat atatatgtat atattatata 300  
 tacatatata tgtatatatt atattatata tacatatata tgtatatatt atattatata 360  
 tacatatata tgtatatatt atattatata tacatatata tgtatatata ttataaatat 420  
 gtataataaa gatttatatg taa 443

&lt;210&gt; 154

&lt;211&gt; 372

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(372)

&lt;223&gt; MAR of chromosome 2 genomic contig; 62076211..62076582

&lt;400&gt; 154

tatataaat tatatatgta attatatatc agtatatata attatatata attatcaata 60  
 tatataatta tatataatta tcaatatata taattatcaa tagatatata taattatata 120  
 tataattata tataattata tatcagtata tataattata taattatata tatgtatata 180  
 taattatag tataaattat ctataagtat atataactat aatatatatc aattatatat 240  
 acttatgtat aattatatat actgatatat aattatatac aattatatat atcaattata 300  
 tataaattat tataattata tatacatata tataaattata tatataaatt atatgtaatt 360  
 atataattac ac 372

&lt;210&gt; 155

## SEL PCT 012.ST25

&lt;211&gt; 484

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(484)

&lt;223&gt; MAR of chromosome 2 genomic contig; 62158581..62159064

&lt;400&gt; 155

attatatata atataaaaat tatacatatt atttattat atattatata cataatatat 60

atatttcata tataatatat attatatata atataaaata tatattatgt ataattatat 120

ataaaatata ttatataatt atatataaca taaaatatat ataatatata attatatata 180

atataaaata tatatatata ataaaatata tattatatgt aattatatat aatatataat 240

atatatataa tataaaatat atattatata taattataat ataaaatata tattatatag 300

tatatattat ataaaatata tattatatat aattatatat tatataaaat atatattata 360

tataattata taatatataa tatatatgt atataattat atataatata aaatatatat 420

aatatatgaa ataagatata tactatatat aatatatata attacatat aagatatata 480

tcat 484

&lt;210&gt; 156

&lt;211&gt; 644

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

## SEL PCT 012.ST25

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(644)

&lt;223&gt; MAR of chromosome 2 genomic contig; 68145036..68145679

&lt;400&gt; 156

tatatatg ctaatatg taatatat tatatatg ctaatatata tatgctaata 60

tataatat attatatata aatatataat atatattat ataaatatat aatatattat 120

atataaatat ataatatata tatatatat atatactata ttatatatta tgtataacat 180

ataatacata ttgttatat ataatatata tatttatgt tatatattat atattatata 240

taataataca atatatttta tatattatat gttatatatt atatattata tataatataa 300

cataatatat aatatatatt atattatata ttacatatat tagcaatatt atatataaaa 360

tatatataat atatataaaa tatatatata aatataaaat atatatcaaa atataaacta 420

tataatatat aaaaatatat tatatatat atataaaaat ataaactata taatatataa 480

aaatatatta tatataatat ataaaaatat attatatatt atatataaaa atatattata 540

tataatatat aaaaatatat ataaaaatata aaaaatatat ataaaaatata aaaaatatat 600

aaaataatat aaaaatatata atatataata atataatata taat 644

&lt;210&gt; 157

&lt;211&gt; 530

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(530)

&lt;223&gt; MAR of chromosome 2 genomic contig; 71257289..71257818

## SEL PCT 012.ST25

&lt;400&gt; 157

atatctatta tatttatata ctttatataa attatatcta ttatatitat atactttata 60  
 taaattatat ctatttatatt tatatacttt atataaatta tatctattat atttatatac 120  
 tttatataaa ttatatctat tatatttata tactttatat aaattatatac tatttatatt 180  
 atatacttta tataaatata taatttatatt tatatacttt atataaatat aattataaat 240  
 atatttatat actttatata aatataatta taaatatatt tatatacttt atataaatat 300  
 aattataaat atatttatat actttatata aatataatta taaatatatt tatatacttt 360  
 ataattatat gttatattta taatttatatt tatataattc ataattatat acattatggt 420  
 tatagttata taatttataa ttatatatat tatatttata tttatataat ttataattat 480  
 ataaattata taaattatat aaattatctt taatttatat tatataatct 530

&lt;210&gt; 158

&lt;211&gt; 337

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(337)

&lt;223&gt; MAR of chromosome 2 genomic contig; 73413615..73413951

&lt;400&gt; 158

acttatatta tatataacta tattattgta tattaatata aattaatgat atataatata 60  
 ttaattatat attattatat gtgatataaaa atacttatat ttatactgta tatatgtata 120  
 tacacacata tatgtatata tgtatatata cacatatgta tatatgtata tgtatatatg 180

## SEL PCT 012.ST25

tatactgtat atatgtatat acacacatat atgtatatat gtatatgtat atatgtatac 240

tgtatatatg tatatacata tatacatata tgatatatat cacatatatg tgatatataa 300

atatatttat ataaatataa tattaatatt tatatta 337

<210> 159

<211> 1340

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(1340)

<223> MAR of chromosome 2 genomic contig; 77011049..77012388

<400> 159

atgtatttta tatagtatat attatgtatt atattgatat aattatataa caattattta 60

tatataaaat aacaaataaa tatataaaat aataaatata tatttattat taaataataa 120

atatatattt attattaaat aataaatata taaagtaata aatatatatt tatatattaa 180

ataattcata tatatttata tattaaataa ttcatatata tttaaataat taatacatat 240

ttaaataatt aatatatatt tatataatat atatttatat attaaataat taatatatat 300

ttatagatta aattaatata tatttatata ttaaattaaa tttaatatat tatatattta 360

tataatttaa atttaataat ttatataatt taatttaatt taatataatt aaaatatatt 420

aaacattata taatatataa tatatttaatt atataatata tatttaatat ataatatatt 480

taatataata tatatttaatt atataatata tatttaatat ataatatatt taatatataa 540

tatatttaatt atataatata tatttaatat ataatatatt taatatataa tatatattta 600

atatataata tatttaatat ataatatata tttaatatat aatatattta atatataata 660

## SEL PCT 012.ST25

tatatttaaat gtataatata tttaatatat aatatatatt taatgtataa tatatttaaat 720  
 atataatata tatttgatgt ataatatatt taatatatat ttgatgtata atatatttaa 780  
 tatataatat atatttgatg tataatatat ttaatatata atatatattt gatgtataat 840  
 atatttaata tataatatat atttgatgta taatatattt aatatataat atatatttga 900  
 tgtataatat atttaatat taatatatat ttgatgtata atatatttaa tatataatat 960  
 atatttgatg tataatatat ttaatatata atatatattt gatgtataat atatttaata 1020  
 tataatatat atttgatata taatatattt aatatataat atatatttga tatatattta 1080  
 atatataata tatatttgaat atataatata tttaatatat aatatatatt tgatatataa 1140  
 tatatttaaat atataatata tatttgatat ataatatatt taatatataa tatatatttg 1200  
 atatataata tatttaatat ataatatata ttgatatat aatatattta atatataata 1260  
 tatatttgat atataatata ttttcttatt aattatttat atataatata taaatatata 1320  
 ttaattaatt atatattaaa 1340

<210> 160

<211> 937

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(937)

<223> MAR of chromosome 2 genomic contig; 78226855..78227791

<400> 160

tgtgtatata tacatatatg tgtatctatg tgtatatata catatgtgta tatatacata 60

tatgtgtata tatacatatg tgtatatatg tgtatatatg tgtatatata catatatgtg 120

## SEL PCT 012.ST25

tatatatgtg tatatatgtg tatatataca tatatgtgta tatatgtgta tatatacata 180  
 tgtgtatata tgtgtatata tacatatatg tgtatatatg tgtatatata catatatgtg 240  
 tatatatgtg tatatataca tatatgtgta tatatgtgta tatatgtgta tatatacata 300  
 tatgtgtata tatgtgtata tatacatata tgtgtatata tgtgtatata tacatatatg 360  
 tgtatatatg tgtatatgtg tgtatatata catatatgtg tatatacaca catatatgtg 420  
 tatatatgtg tatatataca tatatgtata tatacatata tgtgtatata tgtgtatata 480  
 tacatatatg tgtatatatg tgtatatata catatatgtg tatacataca tatatgtgta 540  
 tatatgtgta tatatacata tatgtgtata catacatata tgtgtatata tgtgtataca 600  
 tacatatatg tgtatacata catatatgtg tgtatatgtg tatacataca tatatgtgtg 660  
 tatatatgtg tatacatatg tgtgtatag tgtatatata catatatgtg tgtatatatg 720  
 tgtatatata catatatgtg tgtatatatg tgtatatata catatatgtg tgtatatatg 780  
 tgtatatata catatatgtg tgtatatatg tgtatatata catatatgtg tgtatatatg 840  
 tgtatatata catatatgtg tgtatatatg tgtatatata catatatgtg tgtatatatg 900  
 tgtatatata catatatgtg tgtatatatg tgtatat 937

&lt;210&gt; 161

&lt;211&gt; 1350

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(1350)

&lt;223&gt; MAR of chromosome 2 genomic contig; 79287748..79289097

&lt;400&gt; 161

## SEL PCT 012.ST25

tatatatatt atatatatag taactgttct attatatata tattatatat atttctgttc 60  
 tattatatat tatatatatt atattatata ttatatgtaa tatattatat atattataag 120  
 taatatatta tatatatatt atgtaatat ttatatatat tatatgtaat atattatata 180  
 tattatatgc aatatgttat atatattata tgcaatatgt tatatatatt atatgcaata 240  
 tattatatat attatatgca atatattata tataatatat gtaatatatt atattatata 300  
 ttatatgtaa tatcttatat attatatgta atatattata tatattatat gtaatatctt 360  
 atatatatatta tatgtaatat attatatatt atatgtaata tattatctta tatatatatt 420  
 atgtaatat ttatattata tattatatgt aatatatat attatgtaata tattacatat 480  
 tatatgtaat atatattata tgtaatatat tacatattat atgtaatat tattatatgt 540  
 aatatattac atattatatg taatatatta catattatat gtaatatatt atatgtatta 600  
 tatgtaatat attatatgta ttatatgtaa tatattatat gtattatatg tattatatgt 660  
 aatatattat atgtattata tgtaatatat tatattatt atgtaattat attatatgta 720  
 atatattata ttatatatta tatattatt atgtaatat ttatattata tattatatat 780  
 attatatgta atatattata ttatatatta tatattatt atgtaatat ttatattata 840  
 tattatatat attatatgta atatattata ttatatatta tatattatt atgtaatat 900  
 ttatattata tattatatat attatatgta atatattata ttatatatta tatattatt 960  
 atgtaatat ttatattata tattatatat attatatgta atatattata ttatatatta 1020  
 tatattatt atgtaatat ttatattata tattatatat attatatgta atatattata 1080  
 ttatatatta tatattatt atgtaatat ttatattata tattatatat attatatgta 1140  
 atatattata ttatatatta tatattatt atgtaatat tttatatta tatattatt 1200  
 attatatatt atatgtaata tattatatta ttattatat attatatatt atatgtaata 1260  
 tattatatta ttattatat atattatatt attattata tataatatat tatattatat 1320  
 atattatatt atatatttt ctgttctaatt 1350

&lt;210&gt; 162

&lt;211&gt; 332

## SEL PCT 012.ST25

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_difference

&lt;222&gt; (1)..(332)

&lt;223&gt; MAR of chromosome 2 genomic contig; 81142998..81143329

&lt;400&gt; 162

ctatgtatat aactatatat aactattata taacttaata agatatataa ctattatata 60

acttaataag ttatatataa ctattatata taacttaata agttatatat aactattata 120

taacttaata agttatatat aactattata taacttatta agttatatat aactatatat 180

aacttaataa gttatatata actattatat aacttaataa gttatatata actattatat 240

aacttaataa gttatatata actattatat aacttaataa gttatatata actatatata 300

acttatatac aacttattaa gctatatata ta 332

&lt;210&gt; 163

&lt;211&gt; 327

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(327)

&lt;223&gt; MAR of chromosome 2 genomic contig; 84019536..84019862

## SEL PCT 012.ST25

<400> 163  
 actgacagta tacatactgt atatatatac agtatgtata catatacagt atgtatacta 60  
 tatacagtat gtatactgta tatatatata cagtatgtat actgtatata tatacagtat 120  
 gtatacgtat gtatactgta tatatgtatt atagtgtata tatgtattat agtgtatata 180  
 tgtattatat atattatagt gtatgtatta tatgtgtata tacatataat atattataca 240  
 tatacatatg cacaatatgt atatgtatta tatgtattca tatacatata tgtatatgta 300  
 taatatatgt atacatataa tacacat 327

<210> 164

<211> 407

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(407)

<223> MAR of chromosome 2 genomic contig; 1448030..1448436

<400> 164  
 tatataatat atattacata tatattatat ctatattatt tatattacat atgtaataata 60  
 tattatattt atattattta tataatatat tatatatatt atattattta tatgtaatat 120  
 atttatattg ttatatata ttatatttat attatttata tataatacat attatattta 180  
 tattatttat atataatata tataataaat atataatata tataaaaata tatatattta 240  
 atatatctat aatatatatt atatatatta tatataatat atataattgt acatatattt 300  
 attatatata ttatatatat aatatatatt ataaatataa tatataaata tatttataaa 360  
 tatatataaa tattatattt atacattata ttatatataca tattata 407

## SEL PCT 012.ST25

<210> 165  
 <211> 1959  
 <212> DNA  
 <213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(1959)

<223> MAR of chromosome 2 genomic contig; 2117630..2119588

<400> 165

```
tatacatgtt atagtgata tagtatacta atatataatg tatgtatgtg tatacatata   60
cacatataat atacacatat ataatatata tagtatataa taatgtataa tatataatat  120
ataatataaa atgtatagta tactacatat ttatatatag tatatagtat gcatagtaca  180
tatatactat atatgtagta tactatagtg tatatatagt acaccatata tagtataaat  240
atactatata gtatatgtac tatatatata ctatatagta tatacagtat acatatatag  300
tataacctata ctatatagta tatatagtg gcgtatacta tatagtatat atagtgtagc  360
tatactatat agtatatata gtgtgcgtat actatatagt atatatagtg tgcgtatact  420
atatagtata tatagtgtagc gtatactata tagtatatat agtatacata tatagtgtagc  480
gtatactata tagtatatat agtatacata tatagtgtagc gtatactata tatagtatac  540
atatatagta tatctagagt atatgtagta tgtatagtat atatagtcta catactgtat  600
atacagtata tatatactct atagtatact atacagtata gtatactata tagtatacaa  660
tatatgtata ctatagaaac acactatata tagtatacta tatatactat atactatata  720
ctatatatag tatactatat atactacata ctatatatag tgtatgtata gtatatataa  780
actatatata gtgtatatag tatatatatt atatataata tatattatat tatattatac  840
```

## SEL PCT 012.ST25

tatatattat atgtatatta tagtatatta tactattata tattatatat tatattatat 900  
 attatataat ataataat tatatattat aaaatatata ttttatatt atatatattt 960  
 aaatatattta taatatatat ttataatat atatattata attattttat atataatata 1020  
 aaatataata aatatattat aatatatat tttaaaatat aatatattata tattataaaa 1080  
 atataaatat ataatatatt atatattata tatagtataa tatataatat gttatatagt 1140  
 atcttatact attatactat atatattata tagtgtatat atagtatact atatatagtg 1200  
 tatatagtggt atactatagt gtatatagtg tatactatag tgtatatagt gtatactata 1260  
 tacactgtat atagtagtggt atactatata cactgtatat agtagtggtat actatataca 1320  
 ctgtatatag tagtgtatac tatatacact gtatatagta gtgtatacta tatacactgt 1380  
 atatagtagt gtatactata tacactgtat atagtagtggt atactatata cactgtatat 1440  
 agtagtggtat actatataca ctgtatatag tagtgtatac tatatacact gtatatagta 1500  
 gtgtatacta tatacactgt atatatagta tattatatat actatatatg tatatatagt 1560  
 atacatatat attatatata cagtatatat agtatatata ctatgtagta tatatagtat 1620  
 atatactata tagtatgtat agtatactat atagtatata tagtatatta tatagtatat 1680  
 atactatata gtatatatag tatattgtat atatagtata tatactatat agtatatata 1740  
 gtatatgtta tatatagtat attgtatata tagtatacat agtatgtata tatagtatat 1800  
 atagtataca tatatagtat gtacacagta tatatagtct atatgtatac tacatatagt 1860  
 atacatgtat actatactac atatagtata catgtatact atactacata tagtatacat 1920  
 gtatagtata ctacatatat tatacatgta tagaatact 1959

&lt;210&gt; 166

&lt;211&gt; 520

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

## SEL PCT 012.ST25

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(520)

&lt;223&gt; MAR of chromosome 2 genomic contig; 2119984..2120503

&lt;400&gt; 166

tatgtatgca tcgtatacat atatagtata tatatgtatg catcgtatac atatatacag 60  
 tatatatagt atgcatcgta tacatacagt atactatata tacagtatat acagtatact 120  
 atatatacag tatatacagt atactatata tacagtatat acagtatact gtatatacag 180  
 tatatacagt atatatagta tactatatat acagtatata tactatgtat tctatatata 240  
 gtatagtgt catagtatac atatagtata cactatacta tatatagtat actatatata 300  
 ctctatatag tatatatagt atactatata tagtatatat gtatactata tatagtgtat 360  
 atatatacta tatatagtgt atatatatac tatatatagt atatatatac actatatatt 420  
 gtatagtata gtgtatatat agtatagtat atgtatatat acacatgtat acatgtatat 480  
 atgtatacta atatatacta atatatgtat aaatatatat 520

&lt;210&gt; 167

&lt;211&gt; 954

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(954)

&lt;223&gt; MAR of chromosome 2 genomic contig; 2578285..2579238

&lt;400&gt; 167

## SEL PCT 012.ST25

tattatatat aactttataa tatataatat atattatata taactttata atatataa 60

tatattatat ataactttat aatatataat atatattata tataacttta taatatataa 120

tatatattat atataacttt ataatatata atatatatta tatataactt tataatatat 180

aatatatatt atatataact ttataatata taatatatat tatatataac ttataatat 240

ataatatata ttatatataa ctttataata tataatatat attatatata actttataat 300

atataatata tattatatat aactttataa tatataatat atattatata taactttata 360

atatataata tatattatat actatatata atatataact ttataatata taatatatat 420

tatatactat atataacttt ataatatata atatatatta tatattatat ataactttat 480

aatatataat atatattata tataacttta taatatataa tgtatattat atattatata 540

ttatatatta tatataactt tataatatat aatgtatatt atatattata tataacttta 600

taatatataa tatataatat aatatataac ttataatat atatatcata tattatatat 660

aactttataa tatatatcat atattatata taactataat atatatatca tatattatat 720

ataactataa tatatatatc atatattata tataacttta taatatatat atcatatatt 780

atatataact ttataatata tatcatatat tatatataac ttataatat atatcatata 840

ttatatataa ctttataata tatattatat ataactttat aatatatatc atatattata 900

tataacttta taatatatat catatattat atataacttt ataatatata tcat 954

&lt;210&gt; 168

&lt;211&gt; 452

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(452)

&lt;223&gt; MAR of chromosome 2 genomic contig; 3836217..3836668

## SEL PCT 012.ST25

&lt;400&gt; 168

tttatatata aatatataatc ttatatatat ttatatataa tacatatata tcttatatat 60

ataaaatata tatacatatt tatatatataa atacatatgt attatatata tttatatata 120

atacatatgt attatatata attatataat acatatgtat tatatacaat tatataatac 180

atatttataa atatatatat ttatatattat atatatttat atataaataa atatatattt 240

atagatttat ttatatataat atatatttat ataaatatat atttatatat atttatataa 300

atatatattt atatatattt ctatatatat atataaatat atgtataaat atatatattt 360

atacatatat tcatataaat atatatattt atacatgtat ttatatgaat atatatttat 420

acatgaatt atatgaatat atatttatac at 452

&lt;210&gt; 169

&lt;211&gt; 417

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(417)

&lt;223&gt; MAR of chromosome 2 genomic contig; 3837666..3838082

&lt;400&gt; 169

gatatatata ttatatataa tatatatata aagagatata ttatatatt tatttatata 60

aatatatttc ttatatataa gatatatgta aatatattta ttatatataa tatatttata 120

tatgtaaata tatatttata tatttatata ttatatatt tatttatata aatatatata 180

ttatatatt tatttatata tataaaaata tataaatata aatatatata aatatatata 240

## SEL PCT 012.ST25

attataaata tagaaataaa tataaatata aatatataaa tatatataaa tataaatata 300  
 tataaatata aatatatata aatataaata tataaatgta taaatatata aatataaata 360  
 tatataaata tgtataaata tataaatata taaatatata aaaatatata taaatac 417

<210> 170

<211> 1197

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(1197)

<223> MAR of chromosome 2 genomic contig; 6294846..6296042

<400> 170

tatatactaa tatgtatata taaatatata aatatatata cacgtgtata tataaatata 60  
 tatgtatata taaatatata tacatatatg tatataaaaa tatatacgta tatacgata 120  
 tacgtatata tagatatata cgtatatatg tatatacgta tatatagata tatacgata 180  
 tacgtatata tagatatata cgtatatatg tatatacgta tacatgtgta tatacgata 240  
 tacacatata cgtatacatg tgtatatatg tatatgtata cattatatat acgtatatat 300  
 acatatatgt atacatgtat atataaatat atacatatat gtatatatta tacatatatg 360  
 tatatataat atatatatta tatataaat atatatatta tataatatat atattatata 420  
 taatatatat attatatata atatatatta tattatatat aatatatata tatataaat 480  
 attatatatg tacatatgta cataatgtat atatgtatat atataatata tatgcacatg 540  
 tatatataat atatgtatat tatatatata tatgtatata tgtacatatt atatatgtat 600  
 atatgtacct attatatata catatgtata tatgtacctt ttatatatac atatgtatat 660

## SEL PCT 012.ST25

atgtacatat tatatatata tatgtatata tgtacatat atatatatat atgtatatat 720  
 gtgcatgcat atataatata taatatatta tagattataa tattatatac atatcatata 780  
 ttatatactt atatatatat gtatatatta tatacatatt atatattata tacatataat 840  
 atgtatatat aatatatata tatattatat attatatata atacattatg ttatatatta 900  
 tgttatataa tatatattat ataatatata tatattatat ataatatata catatataat 960  
 aaataatata taattatata tataatatat gcatataaat atgtaatata ttttatatta 1020  
 tatatgatca tatataatat gacatattat atagtattat atatatgata tattatatat 1080  
 gattatatat attatatata aatatatgat tatatataat catatatata aatatatgat 1140  
 tatatgatta tatataaata tatatatatg attatatgat tatatataat tgattat 1197

<210> 171

<211> 362

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(362)

<223> MAR of chromosome 2 genomic contig; 6506971..6507332

<400> 171

tatatatagt gtatactata tatacgctat atgcacacat aaactatata tacagtatat 60  
 aatatgcgta tactatatac acagtatata ctacatgtat actatatata gtatataaga 120  
 tatatactat gtatataata tatatactag gtatatatat ccatatatat actatatact 180  
 atagtatata catatatatg tacgtatata tgtatatgta catatatatg tagtatgtat 240  
 atatatatat atatacacac tatagtatat acatatatat actatatata ccctatatag 300

## SEL PCT 012.ST25

agtatattat atacagtata ctatatatac tatatatacc ctatatagag catgtctatg 360

ct 362

<210> 172

<211> 2578

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(2578)

<223> MAR of chromosome 2 genomic contig; 6507395..6509972

<400> 172

ggtatactat atatactata gagtatactt tatagtatat atacctatat tatatatata 60

tacatacact gtatagtata tatggtatat atactatata tggcatatat agtttatata 120

tatactatat atggtatata tagtttatat atatactata tatggtatat atagtttata 180

tataccatat atggtatata tagtttatat agtacatata gtatatatac acactgtata 240

gtatatatta ttagtatat atactatata tactgtatat atagtataaa tactatatat 300

agtatacact atatactata cactatatat actatatact atatactata tatagtatac 360

tatatagtat atagtatact ctatatgtac tatagagtat actatatata ctatacataa 420

aatattttta tatatagtac agcgtatact atatactata tatagtatac tctatatgta 480

ctatagagtg tagtatatac tatacagtat actctatata tactatacag tacactatat 540

atactatata tagtatattt tatatatagt acagtatata cagtatatat attatactat 600

atgtagtaca tatatagttt agtatatata gtatatatac tatactatat gtactacata 660

tataatagta tatatagtat atatactata ctatatgtag tacatatata gtttagtata 720

## SEL PCT 012.ST25

tatactagta tatagatata tagttatata gatataat agtatatata gtatatatag 780  
 catatatagt atatatgcta tatatactat atagcatata ctataacta tatatacagt 840  
 atatatagca tatatagcat atataatata tatacttttg atatacatat tatatacagt 900  
 atatatagta tatatactgt ataaatatac tatatatacc gtatatgcac actatatgct 960  
 atatatacta tataactat atacagtata tatagtacac tatactatat aaagtatata 1020  
 tagtatacag tacactatac tatatacatt atatatagta tatattatac atagtatata 1080  
 gtatataaat agtatatata gtatatacag tatatatata gcatacttta tatagtatac 1140  
 acagtatata gatactatat atgctatata tagtatctat atactgtata ttatatatac 1200  
 taatatagta tatatgtata tatatactgt atatataata tatacatata tagtatatat 1260  
 actatacata cacactatac atagtatat atactatata tactatatac tatatatacct 1320  
 atatatacta tatagtatat tatatatacct atatatacta tatagtatat tatatatacct 1380  
 atatatacta tatagtatat tatatactat atataccata tatactatat atactgtata 1440  
 gtatactata tatactatat agtatactgt atatactata tagtatactg tatatactat 1500  
 atagtatact gtatatacta tatagtatac tgtatatact atatagtata ctgtatatac 1560  
 tatatagtat actgtatata ctatatatac tatatagtat actgtatata ctatatagta 1620  
 tactatatat actatatacc atatatacta tgtatatact atatatagta tatactatgt 1680  
 atatgctata tatagtatat atagtatata tgctatatat agtatatata gtatatatgc 1740  
 tatatataca gtctatatat agtatatata ctatatagac tatatatata gcatatatac 1800  
 tatatatact atatataata tatatgggtat atacatagta tctatatgta gtatctatat 1860  
 atagtaccta tatatactat atataggtac tatatatagt atatatactt tatatagata 1920  
 ctatatatag tatatactat ttatatagta tatatagtat atgtagcata tatagtatat 1980  
 atagtatata tagtatatag tatgtatagt atatatagat tatattgtat atacagtata 2040  
 tatactgtat atactatata aatagtacat acagtatata cagtatatat gtactatata 2100  
 tagtatatac agtatatata gtatatatgt accatatata gtatatacag tatatacagt 2160  
 atatatgcac tatatggtat atacagtata tacagtatat atgtactata taaatagaat 2220

## SEL PCT 012.ST25

atactctata tacagtatat atgtactata taaatatata cactatgtac agtatatatg 2280  
 tactatataa atagtatata cactatatac agtatatatg tactatatag tgtatacagt 2340  
 atatacagta tatagggtact atatatggta tatacagtat atatgcacta tatggtatat 2400  
 acagtatata tgcactatat atggtatata cagtatatat gtactatata tggatatatac 2460  
 agtatatatg tactatatat ggtatatata gtatatatgt actatatatg gtatatacag 2520  
 ttatacagt atatatgcac tatatatggt atatacagta tacatgtact atatatgg 2578

<210> 173

<211> 598

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(598)

<223> MAR of chromosome 2 genomic contig; 7770400..7770997

<400> 173

gtgtattgta tatacatata cgtatctacg tatatacata tatgtattgt atatacatat 60  
 atgtattgta tatacatata tgtatatacg tatatacata tatgtattgt atatacatat 120  
 atgtatatac gtatatatat atatgtatat acgtatatag atatacatat atatgtattg 180  
 tatatacata tatgtatata catatataca tatatatga tatacatata tatgtattgt 240  
 atatacatat acaatatatg tatatatata tatacatata caatatatgt atatacatat 300  
 atatgtattg tatatacata tatatgtatt gtatatatat atattgatat acatatatgt 360  
 atatatatat atatgcatat atgtatatat acatatatgc atatatgtat atatacatat 420  
 atacatatgt acatatatac atatatacat atatgtatat atacatatat acatatgtac 480

## SEL PCT 012.ST25

atatatacat atatacatat gtacatatat acatatatac atatgtacat atatacatat 540

atagatatat atacacatat atagatatat ttatatgtat atatacatat atacatat 598

<210> 174

<211> 1048

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(1048)

<223> MAR of chromosome 2 genomic contig; 8332422..8333469

<400> 174

cattatatat aatatataat atattattat atataatata tataacatta tatatagtat 60

atatgacata tataacatat attatatata acatatataa aatataacat attatatata 120

acatatataa aatataacat atattatata taacatatat aaaatataac atatattata 180

tataacatgt ataaaatata acatatatta tatataacat gtataaaata taacatatat 240

tatataacat gtataaacta taacatatat tatatatataa atatattata tgttatatat 300

tataaataaa atatattata tgttatatat tataacatat tatataaata atatataata 360

tataacatat attatatataa taatatataa catatattat ataaataata tataacataa 420

catatattat ataacatata acatataaca tatattatat ataacatata acatataaca 480

tatattatat ataacatata acatataaca tatattatat ataacatata acatatatta 540

tattatatat aacatataac atatattata ttatatataa catataacat atattatatt 600

atatataaca tataacatat attatattat atataacata taacatatat tatattatat 660

ataatatata acatatatat tatatatata atataacata taacatatat tatatatata 720

## SEL PCT 012.ST25

ataatatata acatatatta tatataatat aatatataac atatattata tataatataa 780  
 tatataacat atattatata taatataata tataacatat attatatata atataatata 840  
 taacatatat tatatataat ataatatata acatatatta tatataatat aatatataac 900  
 atatattata tataatataa tatataacat atataatata taacatatag catatataat 960  
 atataacata taacatatat tatatataac atataacata tattatatat aacatataac 1020  
 atatataata tgaacatta tatataac 1048

&lt;210&gt; 175

&lt;211&gt; 375

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(375)

&lt;223&gt; MAR of chromosome 2 genomic contig; 8909678..8910052

&lt;400&gt; 175

tatatacaca tatatacgta tgaatatata tacacatata cgtatgaata tatataccca 60  
 tatacgtatg aatatacaca tatatatacg tacgtatata tatacacata tatacgtacg 120  
 tatatatata cacatatata cgtacgtata tatatacaca tatatacgta cgaatatata 180  
 tacacatata tacgtacgaa tatatatata catatatacg tacgaatatata tatacacata 240  
 tatacgtacg aatatatata cacatatata cgtacgaata tatatacaca tatatacgta 300  
 cgaatatata tacacatata tacgtacgaa tatatatata catatatacg tacgaatatata 360  
 tatacacata tatac 375

&lt;210&gt; 176

## SEL PCT 012.ST25

&lt;211&gt; 563

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(563)

&lt;223&gt; MAR of chromosome 2 genomic contig; 10572503..10573065

&lt;400&gt; 176

atttataata tatatgtata aatatatgta tatatttata tttaaataa tgtatatata 60

tttatattta aatatacgta tatatattta tatttaaata tacgtgtata tatttatatt 120

taaataatcg tgtatatatt tatattttaa tatacgtgta tatatttata tttaaataa 180

cgtgtatata ttatatattta aatatacgtg tatatattta tatttaaata tacgtgtata 240

tatttatatt taaataatcg tgtatatatt tatattttaa tatacgtgta tatttatatt 300

taaataatcg tgtatatatt tatttaaata tatgtatgta ttataaata tatattttaa 360

gtatatattt ataaatgtat acatgtatat ataaatatat atattttaa tatatattta 420

tatatatatt tatatattta tataagtata tatatattta aatatatgta tatatttata 480

tatttatata agtatatata tttaaataa tgtatatatt tataatatat attttaaata 540

tatatttata tatttattat ata 563

&lt;210&gt; 177

&lt;211&gt; 595

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

## SEL PCT 012.ST25

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(595)

&lt;223&gt; MAR of chromosome 2 genomic contig; 11609694..11610288

&lt;400&gt; 177

tataaatact atatatagta tatataatat tatatatact atatataaat atagttagta 60  
taaataatat ataatataga tatataatat aatataatat gttataaata taaatatatt 120  
tatataattt aatttataat atataatata taatatataa ttttaattta taatatataa 180  
tatataattt aattttataa tatataatat ataatatgta aattatatat aatttaatat 240  
atctaaatta tataatttaa atataaatat aatataaata tatctaacat aatatacata 300  
acataaatat atatagtata tatagtacat ataaatatat atagtacata tagtatatat 360  
aaatatatag tatatataaa tatagtatat ataaatatat agtatatata tagtatatat 420  
aaatatatag tatatataaa tatatatagt atatataaat aatatatagt atataaataa 480  
tatatattat taaatataat aataatttat tatatatact atatattatt atgtattata 540  
ttatatatat tattttatat ttaatatata ttattttata tatttatatt aatat 595

&lt;210&gt; 178

&lt;211&gt; 662

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(662)

&lt;223&gt; MAR of chromosome 2 genomic contig; 12699804..12700465

## SEL PCT 012.ST25

&lt;400&gt; 178

gtatatatat atatatatat atgggtgtata tatatatata tatatatggt gtatatatat 60  
 atatatatat atgggtgtata tatatatata tgggtgtatat atatatatgc tgtatatata 120  
 tatgggtatat atatatggta tatatatatt tgctatatat atagcagatc tgctatatat 180  
 atatatattgc tatatatata gcagatctgc tatatatatt tgctatatat atgctatata 240  
 tatgctacat atagctata tatatgctat atatatgcta tatatatgct atatatatgc 300  
 tatatatatg ctacatatat gctatatata tgctacatat atgctatata tatgctatat 360  
 atatatgcta tatatatgct atatatatat gctatatata tgctatatat atagctata 420  
 tatatgctat atatatatgc tatatatatg ctatatatat gctatatata tagcatatat 480  
 atatagctat atatatgcta tatatatagc ttatatatat gctatatatg ctatatatat 540  
 gctatatata tagctatata tatgctatat atagctatat atagctaca tatatgctat 600  
 atatatgccca tatgtatgct atatatatgc tatatatata tgctatatat atgctatata 660  
 ta 662

&lt;210&gt; 179

&lt;211&gt; 649

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(649)

&lt;223&gt; MAR of chromosome 2 genomic contig; 12821904..12822552

&lt;400&gt; 179

## SEL PCT 012.ST25

tatgtaatat tatatatata aattatatat tatacatatg taatattata tatatataaa 60  
 ttatatatta tacatatgta atattatata tatataaatt atatattata catatgtaat 120  
 attatatata tataaattat atattatata tatgtaatat tatatatata taaattatat 180  
 attatacata tgtattatat atataaatta tatattatac atatataata tatatataaa 240  
 ttatatatta tacatgtata atatataata attatatatt atacatatat aatatatata 300  
 aattatatat tatacatata taatatatat aaattatata ttatacatat ataatatata 360  
 taaattatat attatacata tataatatat ataaattata tattatacat atataatata 420  
 tataaattat atattatata tatataatat atataaatta tatattatac atatataata 480  
 tatataaatt atatattata catatataat atatataaat tatatattat acatatataa 540  
 tatatataaa ttatatatta tacatatata atatataata attatatatt atacatatat 600  
 aatatatata aattatatat tatacatata taatatatat aaattatat 649

&lt;210&gt; 180

&lt;211&gt; 3191

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(3191)

&lt;223&gt; MAR of chromosome 2 genomic contig; 15356889..15360079

&lt;400&gt; 180

tacaattata tataactata aatataatat aatatatatt atctatatta catattaata 60

tataatatat attacctatt aatatataat ataatatata taatatatat tacctattaa 120

tatataatat aatatatata atatataatta cctattaata tataataaaa tatatataat 180

## SEL PCT 012.ST25

atatatata tattatata taatatata tatataacat atataacata tactatata 240  
 tatataacat atataattgt atagtatta tatatattat atatacttat acataatata 300  
 taaataatta aatatatgtt ataaatataa caaatatata acatatataa catatataac 360  
 atatatataa ttacataaaa tatataatac ataatatata ttatgcaaca tattatataa 420  
 tatataacat ataagtata ttatattata tcatatataa tacataatat ataatatatg 480  
 atataatata atatattata tatgatataa tataatatat tatatatgtt ataataaat 540  
 atatattata tataggatat attataacat attacatatg atataataaa tttatctta 600  
 tatataggat atattataat atacacata tagcatatat taaaatatat tacatatagt 660  
 atattatata tactatatgt atatatacat atagtatatt atagtatatt atacagtata 720  
 tattatata actatatata gtagtatata gtatatatta tatatactat atatagtagt 780  
 atacagtata tattatacag tatatattat atacactata ttatatatta tgtataatat 840  
 atactatata tagtatatta ttagtatat attaaacata atagatatat agtatatact 900  
 atagataata gatattatat agtatatagt atatattata tataatatat ataatatata 960  
 ttatatacat atagtatata tgatatatta tatataatat atataatata taatatatgt 1020  
 aatataatac atattatata taatatatgt aatataatat aatatataat atagtataa 1080  
 taataatata tattatataa tataacatat ataaatataa taatatatat tatatgat 1140  
 aacatacata aatataataa catatataat atatattata tattatattg tatatatgat 1200  
 atactatata ttacacatta tacattattt ataatatata attaatatat aacatatatt 1260  
 agataacata taattatatt tgaacatat ataagatat attacatata taacatatat 1320  
 aattatatat atatttatct aattatatat gaaattatat atgacatata aaattatata 1380  
 ttatatatgt tatatgtatt atatattata tatgttatat atgttatata taacatatat 1440  
 aacatatata acacacacat ataacatata taacatatat tacatatata acatatataa 1500  
 cacatatata attatctaac atagataata tatataatat ataataaac atatatatta 1560  
 tatattatac actctattat attatatata ttatacataa tatataatat atagtatata 1620  
 atataatata ttgtatatat gatataatat atattgtaca tagtataata tacatatata 1680

## SEL PCT 012.ST25

gtatattatg tataacataa tatatagtat attatgtata acataatata tagtatatta 1740  
 tgtataacat aatatatagt atattatgta taacataata tatagtatat tatgtataac 1800  
 ataatatata gtatattatg tataacataa tatatagtat attatgtata acataatata 1860  
 tagtatatta tgtataacat aatatatagt atattatgta taacataata tatagtatat 1920  
 tatgtataac ataatatata gtatattatg tataacataa tatatagtat attatgtata 1980  
 tataatatac atattatata gtatattatg tatatataat atacatatta tatagtatat 2040  
 tatgtatata taatatacat attatatagt atattatgta tatataatat acatattata 2100  
 tagtatatta tgtatatata atatacatat tatatagtat attatgtata tataatatac 2160  
 atattatata gtatattatg tatatataat atacatatta tatagtatat tatgtatata 2220  
 taatatacat attatatagt atattatgta tatataatat acatattata tagtatatta 2280  
 tgtatatata atatacatat tatatagtat attatgtata tataatatac atattatata 2340  
 gtatattatg tatatataat atacatgta ttagtatatat tatgtatata taatatacat 2400  
 gttatgtagt atattatgta tatataatat acatgttatg tagtatatta tgtatatata 2460  
 atatatataa ggtgtatata tattatgtat atataatata taaggatatat atattatgta 2520  
 tatataatat atataagggtg ttatatataat gtatatataa tatataagggt atgtatatta 2580  
 tgtatatata atagtatat tatatataat atataattat tatatacatat atgtatctat 2640  
 ataatatata ttatgtatat attaggtatc tatataatat atattatgta tatatattat 2700  
 gtatctatat aatatatata ttatgtatat atattatgta tctatatata atatatatta 2760  
 tatgtatatt atgtatctat ataatatata taatgtatat agatatatta tatattatgt 2820  
 atatatatta tgtatctatt ttatatataa tgtatataga tatacaatat atattatgta 2880  
 tatattatgt atctatatata tatatattat ttatatagat atatatatta tgtatatata 2940  
 cataatatat tacatattat gtatatatac ataatatata atattatg tatatatata 3000  
 taatatataa tatattatat attacatata ttatatataa tatattatat tatgtatata 3060  
 tattatgtat atataatgta tatataatat ataaagtgta tatatattgt gtatatataa 3120  
 tgtatatata ttacatatat tatgtgtata tatattatac ataatatata tactacatta 3180

SEL PCT 012.ST25  
3191

tacataatat g

<210> 181

<211> 314

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(314)

<223> MAR of chromosome 2 genomic contig; 728676..728989

<400> 181

tgtgtatata tgtatatata atatatatta tataatatgc atagtataa aatatgtata 60

ttatatatgt atatfffata tatatgtata tattatatgt atatfffata tatgtatatt 120

ttatatatat gtatatatta tatatgtata ttttatatat atgtatatat tatatgtata 180

ttttatatat atgtatatat tatatatgta tttttatat atagtatat attatatatg 240

tatatfffat atatatgtat atfffatata tatgtatatc atatatatgt atatattata 300

tatatgtata tcctt 314

<210> 182

<211> 423

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

## SEL PCT 012.ST25

&lt;222&gt; (1)..(423)

&lt;223&gt; MAR of chromosome 2 genomic contig; 737493..737915

&lt;400&gt; 182

ataatatata gtgtctttta tattatctaa tatgtaatat aatgtatfff atattatgta 60

ttttatatta tataatatat aatataatgt atfftatatt atatgttata taatatatag 120

tgcattatat attatgttat attatatata ttttatttat ataaattata tattatatgt 180

tattttatat atattatata acatataata taacaatgca ttatatatta taaaatatat 240

aatacattac atattatata taatatataa tacattacat atattatata atatatata 300

cattatcata tattacaaat attacattag tataatagta attataatat aatatattat 360

atattacata tattatatta atgtaatagt aattataata taatatatat tatattttat 420

att 423

&lt;210&gt; 183

&lt;211&gt; 724

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(724)

&lt;223&gt; MAR of chromosome 2 genomic contig; 1069556..1070279

&lt;400&gt; 183

tattataata tattatatac attatattgt atatatacta tatatgggat atatatgata 60

cataatataa aatgtatatt gtaatatata ttatatatat acatagtgta cattatataa 120

## SEL PCT 012.ST25

tataatataa tgtatattat aatatacatt ataataaat agtgtactat gtatatagta 180  
 tatataatgt atattataat gtattatata gtataatata atataatata cattatatag 240  
 tattgcatta tataatgciat ataatatata atatattatg tatatatata ttatatatac 300  
 tatattatat agtacatata atgtatatta tatagtatat ataataaat acattatata 360  
 tacaatatat aatgtatatt atatagtatg tataatgtaa tacattatac atagtacata 420  
 aagtatatta taatatatta taatatataa tatacattat atattataat gtatataata 480  
 tattgtatat atactatata taatgtatat acaattatat ataattgtat atatacatgt 540  
 atatgtatat gtatatatac atgtatatgt atgtgtatat atacatatat gtatatgtat 600  
 gtgtatatat gtatatgtat atatgtatat gtatacgtat atatgtatat acaatgtata 660  
 tataatgtat ataaaaatat ataatatata caatatgtat ataatgtata taattatata 720  
 atat 724

&lt;210&gt; 184

&lt;211&gt; 383

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(383)

&lt;223&gt; MAR of chromosome 2 genomic contig; 2719918..2720300

&lt;400&gt; 184

atatttatat ttatatatt atttatatat aaatatatat ttatatatta tatattattt 60

atatataaat atatatttat attttatata ttatttatat ataaatatat atttatattt 120

tatatattat ttatatataa atatatatatt atattttata tattatttat atataaatat 180

## SEL PCT 012.ST25

atatttatat ttatatatt atttatatat aaatatatat ttatatatta tatattatt 240  
 atatataaat atatatttat atttatata ttatatatt atatattata tatattata 300  
 ttaattgtg tataatatat attattaat ataataaata tattatttt tatatattat 360  
 ataaaaatat ataatatata aaa 383

&lt;210&gt; 185

&lt;211&gt; 309

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(309)

&lt;223&gt; MAR of chromosome 2 genomic contig; 4994249..4994557

&lt;400&gt; 185

tataatatat aattgttata acattataac aattatatat tatatataat acaattatat 60  
 aatatatatt atataattgt aatatataat ataattatat aatatatatt atataatata 120  
 atatataata tatcatatat gttatatatt ttattatata atatatatta tatataatat 180  
 tatatataat atatattata tataatatta tatataatat atattatata taatatatt 240  
 atatatatta tatataatat atattatata ttaaattatta tatatataat atatataaca 300  
 ttattgtta 309

&lt;210&gt; 186

&lt;211&gt; 740

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

## SEL PCT 012.ST25

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(740)

&lt;223&gt; MAR of chromosome 2 genomic contig; 5034916..5035655

&lt;400&gt; 186

```

ttatatata aaatattata tataatatta tatataatat ttctatata aaatgtgtat    60
ataattatat ataattatat aaaatataat atagaatatc taataatgta taatatataa   120
catataaaaa taatattatt taatatataa tttttatat ataattttt tatatataat   180
ataatatata ttttatatat aattattaat tatataatta atatataata tatattttat   240
acataattat taattatata taattaatat ataatatatc ttatacataa ttatcaatta   300
tatataatta atatataata tatattttat acataattat taattatata taattaatat   360
ataatatatc ttatacataa tatatataaa tatattatat ataatatata ttatatataa   420
tattatatat aatatatatt atatatataa aatttatata taatattata tataatatta   480
tatattttat atacaatatg atatataata taatttatat attatatata ttatatata    540
attattatat aaattatata aatataaatt atatatttat atataattat tatataaatc    600
attatataat tattataatt ataatatata atataatata atattatata taatatatag    660
tattctatat aaataatata acatatattt tatatagaat attatatata atataatata    720
tattttatat agaattattat

```

740

&lt;210&gt; 187

&lt;211&gt; 847

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

## SEL PCT 012.ST25

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(847)

&lt;223&gt; MAR of chromosome 2 genomic contig; 6074678..6075524

&lt;400&gt; 187

```

aatatagaca taaatatata tgcataaata tatatatgca taaatatata taaaaatata   60
tataaatata tacataaata tatataaata tatacaaaaa tatatataaa tatataaaaa   120
aatatataaa tatatataca catatatataa tatatataca tacatatata aacatatata   180
cataaatata tatgtataaa tatatataca cataaatata tgtatgaata tatatacata   240
aatatatatg tataaatata tatacataaa tatataaaga tatatacata aatatatata   300
aatatatata cataaatata tataaatata tataaataga tatataaata tatatataaa   360
tatataaata tatatataaa tatataaata tataaaaaata gatataataa tatatatata   420
aatatataaa tatatatata aatatatata aatatataaa tatatatata aatatatata   480
aatatataaa tatatataaa tatataaata tatatataaa tatatataaa tatataaata   540
tatataaata tatataaata tataaatata tatataaata tatataaata tataaatata   600
tatataaata tataaatata taaatatata tataaatata taaatatata taaatatata   660
taaatatata aatatatata aatatatata aatatataaa tatatataaa tatatataaa   720
tatatataaa tatataaata tatataaata tatataaata tatataaata tatataaata   780
tataaatata tatataaata taaatatata taaatatata aatatatata taaatatata   840
taaatat                                     847

```

&lt;210&gt; 188

&lt;211&gt; 784

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

## SEL PCT 012.ST25

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(784)

&lt;223&gt; MAR of chromosome 2 genomic contig; 6108986..6109769

&lt;400&gt; 188

```

atttatttat atattttaata tataaaatat atattttaata tataaaatgt atatatatat 60
atatattata tataatacaa tatatattat atataatata tattatatat aatattatat 120
attatattat aatataatat atatttatata taatataata tatattatat attattatat 180
ataatataat atatattata tattattata tataatataa tatatattat atattattat 240
atataatata atatatatata tatattatta tatataatat aatatatatt atatatatat 300
tttatatata taatatataa tatatatatt atatatatat ttatatata taatatataa 360
tatatatatt atatatatat ttatatata taatatataa tatatatatt atatatatat 420
tttatatgta taatatataa tatatatatt atatatatat tatatatata taatatgtaa 480
tatatatatt atatatatat tatatatata atatatatata tacataaaat atatattata 540
tataatatat ataatatata ttatatataa aatatatttt atgtataata tatattatat 600
ataatatata atgtatattt atatataaaa tatatatatta tatacaatgt atatttatat 660
ataaaatata tatttatata caatgtatat ttatatataat atgtgtttaa tatatgaaat 720
atatatttat atataatatata tatttaattat ataaaaatata tattaaatat atatttatat 780
ttaa
784

```

&lt;210&gt; 189

&lt;211&gt; 381

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

## SEL PCT 012.ST25

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(381)

&lt;223&gt; MAR of chromosome 2 genomic contig; 10389032..10389412

&lt;400&gt; 189

tatacacata tagagtatat agagtatata tagagtatat ctatagagta tatatgtata 60

tagagtatat aatacagcct accatatata tagtatacat atatataac tctatatact 120

atatatatag tgtgtatata tatagtatag accctacat atatatatat aggagtatat 180

atatatacac actcctacta tatatagtat gtatatagag agtatataga gtatatatac 240

agtatatata cacagtatat atatgccata tatagtatct atatacttat atatagtatg 300

tatctatata cttatatata gtatgtatct atatactata tatagtatgt atctatatac 360

tatatagagt atatatgtat a 381

&lt;210&gt; 190

&lt;211&gt; 507

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_difference

&lt;222&gt; (1)..(507)

&lt;223&gt; MAR of chromosome 2 genomic contig; 11097807..11098313

&lt;400&gt; 190

## SEL PCT 012.ST25

aattatatat aatttattat atataatfff atatttataa tatttttata tacatatfff 60  
atatatcttt ataattatat attacatata taatattata taatatatat aatatatata 120  
atatatatta tatattatat aatatatatt atatatatta tatataatat atataatata 180  
tataatatat ataatatata taatatataa tatatattat ataatatata ttatatataa 240  
tatatattat atataatata tattatatat aatatataat atatatataa tatataacat 300  
ataataatat attatacata atttatatat aatttttata taattatata ttttatata 360  
ttttatata attatatata ttatatatt ttatatataat tatatatatt tatatatfff 420  
tatataatta tatatatataa ttttatataa atatatataa ttttatataa ttttatataa 480  
ttataaaata tataattata tataatt 507

<210> 191

<211> 329

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(329)

<223> MAR of chromosome 2 genomic contig; 11234628..11234956

<400> 191

ttatagttaa atatataaat ataaaatata cagttttata cagtatatat aaaatatata 60  
atatataata cataatacat tagttatata tactatatat actatatata ctacacgtat 120  
agtatatata tgaaactata tatatactat acgtgtagta tatatatgaa actatatata 180  
tactatacgt gtagtatata tatgaaacta tatactatac gtatagtata tatatgaaac 240  
tatatatact atatataact aactataatt gtatatagtt aaaaatataa atataaaata 300

## SEL PCT 012.ST25

tacagttaaa tatattaata tataatagt 329

<210> 192

<211> 584

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(584)

<223> MAR of chromosome 2 genomic contig; 797844..798427

<400> 192

tattatttta tgtataaat agataaaaaat atataactaat atatatgtac ttatatatac 60  
 atcaatatat aatgtattat ttatactaa cgtatattat atatactagt atataatcta 120  
 tattatttta tatgtataa atatataata aaatatataa atattttatg catatatataa 180  
 tatataatat atactaacat gctaatttat atatacttat atataattta tatagtatat 240  
 aatatataaa tgtatataat acataattta tatatttata tattaatagt ttatatatta 300  
 gtatatatac taattttata tactaataaa taaattatat aatatataaa ttatatatta 360  
 tagtacataa tatatattat atagttaa ataatatgtaa ctataatata taactatata 420  
 tgatatacag ttatatataa tataaatttt acatacagta tataaattat atactatata 480  
 ttatatata tatggatat aaattatata ctatacat ttatatacat ggtatataaa 540  
 ttgtatacta tataatgtgt attagtatat atactaatat atac 584

<210> 193

<211> 363

<212> DNA

## SEL PCT 012.ST25

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(363)

<223> MAR of chromosome 2 genomic contig; 1093824..1094186

<400> 193

```
tatacacaca catatatata cacatatata tacacatata tatatacaca tatatatata    60
catatatata cacgtatatata tgtatacaca tatatatgta tatatatata catatatata    120
cacatatata cgtgtatatata cgtatatacg tacatatata cgtgtatatata cgtatatgcg    180
tacatatata cgtgtatatata cgtatatgcg tacatatata cgtgtatatata cgtatatgcg    240
tacatatata cgtgtatatata cgtatatgcg tacatatata cgtgtatatata cgtatatgcg    300
tacatatata cgtgtatatata cgtatatgcg tacatatata cgtgtatatata cgtatatgcg    360
tac                                                                    363
```

<210> 194

<211> 545

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(545)

<223> MAR of chromosome 2 genomic contig; 3456187..3456731

## SEL PCT 012.ST25

&lt;400&gt; 194

tattataata tataattata tattataata tatattatat tatatattta tatattataa 60

tatatattat attatatatt tatatattat aatatatatt atattataat atatatattat 120

attataatat attatattat aatatatatt atattattat atattataat ataatatata 180

ttataatata tattatatta taatttatat attatatata ttataatata tattatatta 240

tatatatatt tatattataa tatatattat tatatattat atattataat ttatattata 300

ttacaatata tattataaat atatatatta tattataaat atatatattt atattacaat 360

atatattata aatatatatt ttatattaca atatatatta taaatatata tattatatta 420

caatatatat tataaatata tattatatta caatatatat tatattataa tatatattta 480

tatatgatat attatattta atatatatta taacataata tataatata aatatattaa 540

tataa

545

&lt;210&gt; 195

&lt;211&gt; 356

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(356)

&lt;223&gt; MAR of chromosome 2 genomic contig; 5001567..5001922

&lt;400&gt; 195

tataaaatat atgttatata tataatatat attatataat atataatata tataatatat 60

aaaatatata aaatatataa tatataatat aatatataat atatatata tataaaatat 120

atataatata aaatatatat aatatataat atatataata tataatacat ataatatata 180

## SEL PCT 012.ST25

atatataata tataatatat ataatatata atatataata tataatatat ataatatata 240  
 atatataata tatataatat ataatatata atatataata tataaatata taaatatata 300  
 tacacacata cacacacata tatgcatata tatacatata catgtgtaca tagata 356

<210> 196

<211> 321

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(321)

<223> MAR of chromosome 2 genomic contig; 5457330..5457650

<400> 196

tatacaatat attataaatt atatataatt tatatataat atatattata tataaattat 60  
 atataattta tataatatat aaattatata taatataaat tatatataat ttatataata 120  
 tataaattat atattatata aattaaatat aatttatatt atatataaat tatatttaatt 180  
 ttatataata tataaattat atttaattta tatataatat aaattatatt tttatatatt 240  
 atgtataatt tatatattta tacatatata cattataata tattgtatag tatatataat 300  
 atatagtata tataaagcat a 321

<210> 197

<211> 361

<212> DNA

<213> Homo sapiens

## SEL PCT 012.ST25

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(361)

&lt;223&gt; MAR of chromosome 2 genomic contig; 8124469..8124829

&lt;400&gt; 197

```

tatatataat atatattata tatattatat aaattatata taatatgtaa tataaatttt   60
gtaatataaa ttatatatat aaattatata taatatatat taatatatat aatataaatt   120
aatatatata atatataatt atatataatt tatatgatat atataaatat atattatata   180
taaattatat atatcataaa ttatatatca tataaattat atataatata cattatgtac   240
ataatatatg atatataata tataatatat attatatata attatatata tataattata   300
taatatatat aaattataat atataatata tataaattat aatatataat atatataaat   360
t                                     361

```

&lt;210&gt; 198

&lt;211&gt; 418

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(418)

&lt;223&gt; MAR of chromosome 2 genomic contig; 11151485..11151902

&lt;400&gt; 198

```

atgtaactat atatatagta tatatagtat atatatacta tatagtgtgt atatatagta   60

```

## SEL PCT 012.ST25

tatatatact atagtagtg tatatatagt atatatatag tgtatatatc gtatatcac 120  
 tatatactat atagtgata tatagatat tagtatata tagtatatat agtatagtat 180  
 atagtagata tatagtgat atatactga tatatagtg acatagtata ctatatagta 240  
 tacatatagt acactgata gtatatatag tatagtatat atagtataca tagtatacta 300  
 tatatagtat agtatacata gtatactata tagtatatag agtatatata cagtatacta 360  
 tatagtatat agagtatata tacagtatac tatatcgtgt gtatagagta tatataca 418

<210> 199

<211> 394

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(394)

<223> MAR of chromosome 2 genomic contig; 13591477..13591870

<400> 199

ttatatatat ttatatata ttatatatat ttatatata ttatatatat attatatata 60  
 tattatatat aattatatat aatatatatt atatatatta tatataatta tatataatat 120  
 atattatata tattatatat ataatatata tataatatat atattttata tatgtattat 180  
 atatatatta tatatatatt atatatatta tatatatatt atatatatta tattttatat 240  
 atataatata acatatataa tatataatta tatattatat atatatatta ttatatataa 300  
 tatatatatt atataatata atatatatt atatatatta tatattttat atattttatat 360  
 aaaaattatt ttatatatt ttatatataa atat 394

<210> 200

## SEL PCT 012.ST25

&lt;211&gt; 1194

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(1194)

&lt;223&gt; MAR of chromosome 2 genomic contig; 14996824..14998017

&lt;400&gt; 200

```

taatatttat atatacatat aaaatttata tataatatat aatatttata tatacatata    60
aaatttatat atatatataa ttttatata tacatatataa atttatatat aatatataat    120
atttatatat acatatataaa tttatatata atatatataa tttatatata catataaaat    180
ttatatataa taaatattta tatatacata taaaatttat atataattta tatataacat    240
ataatattta tatataaaat ttatatataa catatattta tatataattt atatatataa    300
tataatattt atatatataa tatatttatt tatacaattt atatatataa tataataactt    360
atatatacat acataattta tatgatatat atttatataa taatttatat gatatatataat    420
atatctaata tatattatat atatttatata tatttatatat aatttatata atatatatta    480
tatatataat ttatatataa tatatattat atatatataat tatataatat atatattata    540
tatataattt atataatata tatttatatat ataatttata taatatatat tatatatata    600
atttatataa tatatattat atataattta tatataacat attttatata catatatataat    660
ttatatataa tatatattta catatacata tataattttt atataatata aaatatttct    720
atatacatat ataattttta tataatataa aatatttcta tatacatata taatttttat    780
ataatatata ttctatatata catgtctaata ttttatataa tatatatttc tatatacata    840
tataattttt atataatata taatattttt atatacataa tttttatata atatatattt    900

```

## SEL PCT 012.ST25

acatatacat atataatfff tatataatat atatttatat atacatatat aatttttaca 960  
 taatatatat tatatatata tatataatff atatacaaca tataatatat acatatataa 1020  
 ttatatataca acatataata ttatgtata catatataat gtatacacia tatataatat 1080  
 ttatatatac atatataatt tatatgtaat atatacatat ataatttata tgtaatatat 1140  
 atacatgtat aatttatatg tagtatatat acatgtataa ttatatgta gtat 1194

<210> 201

<211> 487

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(487)

<223> MAR of chromosome 2 genomic contig; 14998429..14998915

<400> 201

tagtatacat ttacacatac atgtataatt atatgtaata tataatattt acatatataa 60  
 ttatagataa tatatatffa catatacata tataattata tataatatat aatgtttaca 120  
 tatacatata taattatata taatatatat ttaaatacac atatacaatt atatataata 180  
 tatatttaca tatgcatata taattataga taatatatat ttacatatac atatataatt 240  
 atatataata tataatgttt acatatacat atataattat atataatata tatttaaata 300  
 tacatataca attatatata atatatatff acatatgcat atataattat agataatata 360  
 tatttacata tacatatata attatatata atatataata ttacatata catatataat 420  
 gtatatataa tatataatat ttacatatac atatataatt tatatataat atatattata 480  
 tatatta 487

## SEL PCT 012.ST25

&lt;210&gt; 202

&lt;211&gt; 421

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(421)

&lt;223&gt; MAR of chromosome 2 genomic contig; 16562490..16562910

&lt;400&gt; 202

tatatgaata tatatatgaa tatatacgta tatatgaata tatacatgta tgtatatatg 60

aatatatgta tatatatgaa tatatatgta tatatgaata tatgtatata tatgaatata 120

tatgtatata tgtatatata tgaatatata tgtatatatg tatatatatg aatatatatg 180

tatatatgta tatatgtata tatatgaata tatatgtata tatgaatata tatgaatata 240

tatgtatata tatgaatata tatgaatata tgtgtatata tatgaatata tatgtatata 300

tatgaatata tgtatatata tatgaatata tatgtatata tgtatatatg aatatatatg 360

tgtatatgaa tatatatatg aatatatatg tgtatatgaa tatatatgaa tatatatgtg 420

t 421

&lt;210&gt; 203

&lt;211&gt; 479

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

## SEL PCT 012.ST25

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(479)

&lt;223&gt; MAR of chromosome 2 genomic contig; 21592301..21592779

&lt;400&gt; 203

tatatgtata cgtatataat atattatata ttatatacgt gtacgtatat atgtaataata 60  
 taatgtatat gtacacgtat ataatatata atatattata tacgtatacg tatacattat 120  
 atattacata tatacgtata tacgtatata aaatatatgt atatattata tatacgtata 180  
 taatatatat tatataatat ataatatata cgtatacata taatatatta tatatacata 240  
 ttatatatta tatatttaaa ttatatatta tatcatatat aatatatatg atataatata 300  
 taatatacat atattacata atatattatta tatacatata catatataat atataatata 360  
 ttatatacat atacatatat aatatataat atattatata catatacata tataatatat 420  
 aatatattat atacatatat atatataata tataatatat tatatatata tattatata 479

&lt;210&gt; 204

&lt;211&gt; 870

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(870)

&lt;223&gt; MAR of chromosome 2 genomic contig; 22557584..22558453

&lt;400&gt; 204

tataatatat aatatacata atatgtatat ttatacaca atataaataa tatacataac 60

## SEL PCT 012.ST25

atatatgtat attttatata tgtatatttt atatatattt tatatatattt atatatatgt 120  
 atattttata tataatatat atattgtata taataatata taatatatta tattatatat 180  
 aatatatata atatatatat aaatatatat tatatataat atgtataata tataatattt 240  
 tatatataat atgtataata tatattttat atataataat atgtacaata tatattttat 300  
 atataataat atgtacaata tatattttat atataataat atgtacaata tatattttat 360  
 gtataaatatg tataatatat attttatgta taatatatat ttatgtata atatatattt 420  
 tacgtatatt ttatatataa tatataatat ttatatata atatataaca ttatatatat 480  
 aatatataat attatatata ttatatattt tatatataat atatataaat atatatattt 540  
 tatatataat atattttata tataatatat ataaatatat atattatata taatatattt 600  
 tatatataat atattttata tataatatat aatatatttt atatattata tataatatat 660  
 tatatattat atataatata ttatatataa tatataatat ataatatatt atatataata 720  
 tataatatat aatatattat atataatata taatatataa tatataatat attatatata 780  
 atatataata tgtaatatat aatatattt atataatata taatataata tataatattt 840  
 tatatataat atataatata taatatataa 870

<210> 205

<211> 1086

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(1086)

<223> MAR of chromosome 2 genomic contig; 30591960..30593045

<400> 205

## SEL PCT 012.ST25

gtatatataa tatatattat attatgttat atattatgta gactatgtat taaatatatg 60  
 tatatattat atataaatat ataatatata ttataattt ataattataa atatatttat 120  
 aatatatttt tctaaatatt tatatattat atattatatt taatgatata taataaatat 180  
 atttctaata tttttatat ttataaatat ttatatata ttatatattt tatatatact 240  
 atatattata tatttatatat ttatatata ctatatatta tatagtatat attttatata 300  
 tactatatat tatatattat atattttata tatactatat attatatatt atatatttta 360  
 tatatactat atactatttta ttatatattt tatatatact atatactatt tatttatatat 420  
 ttatatata ctatatacta ttatttatat attttatata tactatatat tatatattat 480  
 atattttata tataatatat atttattata tttttatat attatatata ttatatatta 540  
 tatatttata tatttatataa tatatattat atatagaata tataatatat attatatata 600  
 atataatata atatatatta tataaaatat atataatata taaaatatat aatatatgat 660  
 atatataata tatattctat atttatacat atatatttaa tatttatatta atatataatt 720  
 atatattatc atatgtaata atagatatata tatgtaatat ataaattata attatatatt 780  
 aatatttatat attatttaaat atgtatatatt acacatatat taattattaa atatatatat 840  
 ttaatatatt aaatattatg tattaaatat atataatata ttataaata ttttatatat 900  
 aatatataca tatattaaca tatatgtata tatgtatata ttatatataa cattatatat 960  
 attatgttac atatactata ttttatatgt tacatatact atatattata tgttacatat 1020  
 aatatatata acatatatta taatatgtaa catattatat ataacatatata atatatagta 1080  
 tatata 1086

&lt;210&gt; 206

&lt;211&gt; 406

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

## SEL PCT 012.ST25

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(406)

&lt;223&gt; MAR of chromosome 2 genomic contig; 36233909..36234314

&lt;400&gt; 206

attataaata tatattatag atattagata ttatagatat aatatatata atatatatta 60

tagatattat agatatagat ataatagata ttatagatat tatagatata atatatatta 120

tagatattat agatataata tatattatag atattataga tataatatat attatagata 180

ttatagatat aatatatatt atagatataa tatatattat agatattata gatatagata 240

ttatagatat tatatatatt atagatataa tatatattat agatattata gatatagata 300

ttatagatat aatatatatt atagatatta tagatataat atatattata gatattatag 360

atataatata tattatagat ataagatata ttatagatat tacaga 406

&lt;210&gt; 207

&lt;211&gt; 797

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(797)

&lt;223&gt; MAR of chromosome 2 genomic contig; 36271745..36272541

&lt;400&gt; 207

atataaacat atacgtatat acacatatat acaaatacat atatacatat attatatata 60

tgtatatata ttatattata catatattat atatatatta tattatacat atatacatat 120

## SEL PCT 012.ST25

acacataaac atattacata catatacaaa ttatacacat atacatatat acatatatgt 180  
 atatacatat attatatata aatatatgta tataaaatgt acattatata tacatatata 240  
 ttatgtataa ataatatata aaataaacat aatatatatt tatagatatg atatatataa 300  
 tatatatgta tacatatata catatatgta tatataatgt acattatata tacataaaca 360  
 tcatatataa atgttatata tataatataa atatatataa tatataatat atactttata 420  
 tactatatat aatatatata atagtatata acatatacta tatatactat atataatata 480  
 tactatatat actgttatata atatatataa taatatatac tatatatact aaatataata 540  
 tacataatat aatatatact atatatataa tataatatat aatatagtat atatactata 600  
 tataataatt acatattata tattatacat tatatatatt ataattatta tatataatta 660  
 tatattacat actttgtata taatgtaaat atacattaga atatatatg tatatatatg 720  
 tacatatata atgtatatat gtatacatta tataaactat atataaacat tatattatat 780  
 aaacattata tataaac 797

<210> 208

<211> 423

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(423)

<223> MAR of chromosome 2 genomic contig; 36498521..36498943

<400> 208

tattatatta tatatttaatt attatatatt taatatatta tatatttaatt attatatatt 60

taatatatta tatatttaatt attatatatt taatatatta tatatttaatt attatatata 120

## SEL PCT 012.ST25

taatatatta tatatttaaat attatatata taatattata tatataatat tatatattta 180  
 atattatata tataatatta tatatataat atattatata tttagtatta tgtatttaaat 240  
 atattatata tttagtatta tgtatttaaat atattattta tttagtatta tatatttaaat 300  
 atattattta tttagtatta tatatttaaat atattatata tttaatatat tatatattta 360  
 ttatatattg tatatttaaat atattatata ttattatat attatatata attatatatt 420  
 taa 423

&lt;210&gt; 209

&lt;211&gt; 304

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(304)

&lt;223&gt; MAR of chromosome 2 genomic contig; 37179891..37180194

&lt;400&gt; 209

gtgtatatat atcatatata ttatatcata tatatgtgta tatatatcat atattatattc 60

atatatatgt gtatatatat catatatata tcatatatgt gtatatatca tatatattat 120

atatcatata tgtgtatatata tatcatatat tatatatcat atatatgtgt atatatcata 180

tatattatat atatctcata tgtgtatatata tatcatatat aatatatatg tgtatatatc 240

atatatcata tataacatat atatgtgtat atatcatata tataacatat atcatatatg 300

tgta 304

&lt;210&gt; 210

&lt;211&gt; 693

## SEL PCT 012.ST25

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(693)

&lt;223&gt; MAR of chromosome 2 genomic contig; 38440448..38441140

&lt;400&gt; 210

tatatattct ttatatatt atatataata tatattcttt tatatattat atatagtata 60  
tattctttta tatattatat atagtatata ttcttttata tattatatat agtatatatt 120  
cttttatata ttatatatag tatatatatt ttatatatt atatatagta tatattcttt 180  
tatatattat atatagtata tattctttta tatattatat atataatata tattctttta 240  
tatatcatat ataatatata ttcttttata tattatatat aatatatatt cttttatata 300  
ttatatatca tgtatatata atatacaaaa tatatataga ttftatatat agattattac 360  
ataatagaat atattatata ttatatataa tatatacata atatataata ttatatatga 420  
tataatatat atcatatata tcatataata tatattatat atcatatatt atatataata 480  
atatatagat tatatataat tatatatata atatatataa ttatatatat tatctatata 540  
tagataatat atataattat atataatata ttatatagat tatatataat tatattatat 600  
acaaaatota tatataatat atattatatt atatataata tacataacta tataaaaaat 660  
ataatatata atatatataa tatataatat ata 693

&lt;210&gt; 211

&lt;211&gt; 471

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

## SEL PCT 012.ST25

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(471)

&lt;223&gt; MAR of chromosome 2 genomic contig; 38887582..38888052

&lt;400&gt; 211

aacatatata ctatatatat tatatactat attatatatt atatatataa acatatatac 60

tatatataat atataaacat attatattat acatgatata gataaacata tatattatat 120

ataatataga taaaatatgt tatatataat ataagtata gacatatatt atatatacat 180

atattctaca tatattatat atatatctca cacatattat attatatata catatatctt 240

acatatatta tatatacata tattctacat atattatata tacatatatt ctacatatac 300

atatatacat atattatata tacatatatt atagatatat aatatataaa catatataat 360

attattatat ataatatata taataatatt atataatata taataatatt atatcttata 420

tataaataat atatatattt tatatatata atattatata tatataatat a 471

&lt;210&gt; 212

&lt;211&gt; 1221

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(1221)

&lt;223&gt; MAR of chromosome 2 genomic contig; 43885944..43887164

## SEL PCT 012.ST25

&lt;400&gt; 212

catataaaca tatatttat gtaacatata aacatattat atgtaacata taatatataa 60  
 tatataaaca tatattttat atatttatg ttacatataa tatataatat ataaacatat 120  
 attatatatt atagtgaaca tataatatat aatatataaa catatatttt atatataata 180  
 tataaacata ttttatatat aatatataaa catattttat atataatata taaacatata 240  
 ttttatatat aatatataaa catattttat atataatata taaacatata ttttatataa 300  
 tatataaaca tataatatat ataatatata aaagtatata atataaatat atataatata 360  
 aacatatata atataaatat atataaaata taaacatatg taatatataa acatatatta 420  
 tatataatat ataaacatat attatacgta caatatataa acatatattg tacgtacaat 480  
 atataaacat atattatagc tacaatatat aaacatatat tatacgtaca atatataaac 540  
 atatattata cgtacaatat ataaacatat attatacgta caatatataa acatatatta 600  
 tacgtacaat atataaacat atattatagc tacaatatat aaacatatat tatacgtaca 660  
 atatataaac atatattata cgtacaatat ataaacatat attatacgta caataaacat 720  
 atattatagc tacaatatat aaacatatat tatacgtaca atatataaac atatattata 780  
 cgtacaatat ataaacatat attgtacgta caatatataa acatatatta tatgtataat 840  
 atataaacat ataatatata atatatttta tatatatggt tattatatat gtttatatat 900  
 tatataaac atatattatt atattatata tgtttatata ttatatatta tataatatat 960  
 atgtttatat attatatatt atataatata tatgtttata tattatatat tatataatat 1020  
 atatgtttat atattatata ttatatataa tatatgttta tatattatat attatataat 1080  
 atatatgttt atatattata tattatataa tatatatggt tatatattat atattatata 1140  
 atatatatgt ttatatatta tatattatat aatatatatg ttatatatt atataaataa 1200  
 taaacttaca tattttatta a 1221

&lt;210&gt; 213

&lt;211&gt; 543

&lt;212&gt; DNA

## SEL PCT 012.ST25

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(543)

&lt;223&gt; MAR of chromosome 2 genomic contig; 45818200..45818742

&lt;400&gt; 213

tatgtatatata tacatatata ttatacatg tatatatgta tatatacata tatatttata	60
catgtatatata tatacatata tttttataca tgtatatata tacatatata ttatacatg	120
tatatatata catatatatt tatacatgta tgtatatata catatatatt tatacatgta	180
tgtatatata catatatatt tatacatgta tgtatatata catatatatt tatacatgta	240
tgtatatata catatatatt tatacatgta tgtatatata catatatatt tatacatgta	300
tgtatatata catatatatt tatacatgta tgtatatata catatatatt tatacatgta	360
tgtatatata catatatatt tatacatgta tgtatatata catatatatt tatacatgta	420
tgtatatata catgtatatatt tatacatgta tgtatatata catgtatatatt tatacatgta	480
tgtatatata catgtatatatt tatacatgta tgtatatata catgtatatatt tatacatgta	540
tac	543

&lt;210&gt; 214

&lt;211&gt; 463

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

## SEL PCT 012.ST25

&lt;222&gt; (1)..(463)

&lt;223&gt; MAR of chromosome 2 genomic contig; 47055478..47055940

&lt;400&gt; 214

atacatacat atatacatat atacacatat atatatataa tacacacata ttacatata 60  
 tacacacata tatacatata tacatatata cacatatata catgcataca catatatata 120  
 tatatacaca catatacaca catatatata tatatacaca tatatacaca tatacacata 180  
 tatacacaca tatacatata tacacatata tacatatata catatatata cacatatata 240  
 catatatata tatacacata tatacacata tacatatata cacatatata cacatatata 300  
 catatatata catatatata tatatacaca tatatacaca catatacaca tatatacata 360  
 tatacatatg tatacacata tatacatatg tatacacata tatacacata tacatatata 420  
 catacacata tatacgtata tatgtgtata tatacacata tac 463

&lt;210&gt; 215

&lt;211&gt; 2482

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(2482)

&lt;223&gt; MAR of chromosome 2 genomic contig; 47492696..47495177

&lt;400&gt; 215

aatatatata aaatatatta tattctatgt aatatataga atatataaaa tatattctat 60  
 atattatata gaatatatat ttataatat atattattta tatattttta tatatttata 120

## SEL PCT 012.ST25

ttatttatat atttatatat aatttatata atttatatat ataatttatata tataatttat 180  
 ataaattata tatataattt atatataatt tatatataat ttatataaat tatatatata 240  
 atttatatat aatttatatg atttttatat ataatttatata tataatttat ataattttta 300  
 tatataattt atatataatt tatataattt ttatatataa ttatataat atatatatat 360  
 aatttatata taatttatat aatttatata tataatttat atataattta tataatttat 420  
 atatataatt tatatataat ttatataatt tatatatata atttatatat aatttatata 480  
 atttatatat ataatttatata cataatttat ataatttatata tatataattt atataattta 540  
 tatatataat ttatatatat aatttatata atttatatat atgatttatata taatttatat 600  
 atataattta tataatttat atatataaat tatatatata atttttatat aatttatata 660  
 ttataattt atatatttat ataatttatata tatttatataat ttatatattt atataattta 720  
 tatatttatata atttatatat ttatataatt tatatataat tattcatata ttatataat 780  
 ttacatatata ttatttatat attcatatat aatttatata ttatatata atttatatat 840  
 aattatttac atatttatat atttatatat aatttatata tatttatata taatttatata 900  
 ataaaatata taatatataa tatataatat tataatagat aaaatatata ctatatatta 960  
 tatattttac atttatattta atatttatatg tataattttta tatcatatat aatatatatg 1020  
 atatatatataa ttatatatca tatataatat ataggtatata tataattttta tatcatatat 1080  
 aatatatatg gtatatataa ttatatatca tatataatat atgatatata atttttatatc 1140  
 atataatatata tatttatatat aatttttatat ctacatatata tatatttatat atacaatttt 1200  
 atatctatct ataatatata ttatatataac aatttttatat ctatatataa tatatttatat 1260  
 atacttttat atttatatata aaatgtatat tatatatatact ttatatattat atataaaatg 1320  
 tatatttatat ataattttat ttatatata aaatgtatat tatatataat ttatttttat 1380  
 atataaaatg tatatttatat ataattttat ttatatata aaatgtatat tatatataat 1440  
 ttatttttat atataaaatg tatatttatat ataattttat ttatatataa aaatgtatat 1500  
 tatatataat ttatatattat atataaatatg tatatttatat ataattttat atttatatata 1560  
 ataggtatat tatatataat ttatatattat atataaatatg tatatttatat ataattttat 1620

## SEL PCT 012.ST25

attatatata atagtatat tatatataat ttgtattat atataatag tatattatat 1680  
 ataattttat attatatata atagtatat tatatataat ttatattat atataatag 1740  
 tatattatat ataattttat attatatata atagtatat tatatataat ttatattat 1800  
 atataatag tatattatat ataattttat attatatata aaatgtatat tatatataat 1860  
 ttatattat atataatag tatattatat ataattttat attatatata atagtatat 1920  
 tatatataat ttatattat atataaaatg tatattatat ataattttat attatatata 1980  
 aaatgtatat tatatatatt atataaaaa tgtatattat atatattata tataaaatgt 2040  
 atattatata tattatatata aaaatgtata ttatatatat tatatataaa atgtatatta 2100  
 tgtatattat atataatgta tattatgtat attatatata atgtatatta tatataatat 2160  
 atattatata taatgtatat tatataatat atattatata ttataatata taatatatcat 2220  
 tatatattac atattatata taatatatta tatattatat attacatatt atataata 2280  
 tattatatat tatattaaat atatatttta tatattatat attatatatt atataaaaata 2340  
 tatatattat atattatata aaatatatat atattatatt atatattata ttaaatatat 2400  
 attttatata taatatatat aatatataat atataaaaata tatattatat attatatata 2460  
 aattatatat attatatata aa 2482

<210> 216

<211> 539

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(539)

<223> MAR of chromosome 2 genomic contig; 47561069..47561607

## SEL PCT 012.ST25

&lt;400&gt; 216

aacagtaata taccactaat atataataat atataacagt aatatatcat taatatataa 60

tatatcatta gtatataata ttaatatata ttaatatata atatatcata tacaatatta 120

atatatatta atatataata atatattatt aatgtataat agtaatataa tatattatca 180

atatatatta ctaatatata ataatatatc gttaatatat aatagatcat taatatataa 240

tgttaatata ttatgaatag ataatatatc agtatataat attaatatat taatatatta 300

tatattattt aataatatat aatatattaa taaataatta tatattaata tagcaatata 360

ttaatatatg actgtattat attattaata tataacaata tattatatat tatataataa 420

tttattatat aatatataat aatatattat atattatata acatattaat aatacataat 480

aacattaata atatataata atgttaatat attattatat tatatattaa tatataata 539

&lt;210&gt; 217

&lt;211&gt; 336

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(336)

&lt;223&gt; MAR of chromosome 2 genomic contig; 52853648..52853983

&lt;400&gt; 217

tatatacata aaatatatat attttatata tatacataat atatatatgt atattttatg 60

tatatatcta taatatatat aatataataa aatatacata tatattttat atatataa 120

tatacatata aaatatacat acataaaaata tacatgtata ttttatgtat atataatata 180

tatataaaat atacatgtat attttatata tataatatatc atgtataatt aatatacatg 240

## SEL PCT 012.ST25

tatgttatat atattacatg tatattatat ataatafaca tataaatttt aaatttagtg 300

tatattacat gtatattata tataatatat gtatat 336

<210> 218

<211> 406

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(406)

<223> MAR of chromosome 2 genomic contig; 54866263..54866668

<400> 218

tacgtatata aaaatgtata ttacatata taaaataaat attttatata cgtatataaa 60

atatatattt attttatata cgtatataaa atatttattt tatatatgta tataaaatat 120

ttattttata tacatgtata ttaaatatat atttatatat gtatataaaa atatataffa 180

tatacatgta tataaaatat atatttatata tgtatataaa aatatatatg tatataaaat 240

atatataffa tatagatata taaaatatat atttatataga tatataaaat atatataffa 300

tatagatata taaaatatat atatttatata gatataataa atatatatat tatatagata 360

tataaaatat atatattata tagatatata aaatatatat attata 406

<210> 219

<211> 1452

<212> DNA

<213> Homo sapiens

## SEL PCT 012.ST25

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(1452)

&lt;223&gt; MAR of chromosome 2 genomic contig; 55113305..55114756

&lt;400&gt; 219

ataatatata atatatttg tatattatat tattatatat tatatattat taaatatata 60  
 tattatatta tatattatat aatatatatt atatataata atatagaata tataattata 120  
 tattatatta tatattatat aatatatatt atatataata atatagaata tataattata 180  
 tattatacta tatattatat aatatatatt atatataata atatagaata tataattata 240  
 tattatataa tatgtgaata atgtaata taattatatt attacatat tatataatat 300  
 ataattatat tatataatat ataattatat tatttgata ttatatataa catatacatt 360  
 atattatata taatataatt atatataatt aattataaat taattatata taattatata 420  
 atataatata taatatacat aatatataat atataatata taatatacat aatatatat 480  
 atattatata taatataata tatataatat aatatatat aatgtataat ataattatat 540  
 attatatata atatataatg ttatatatt atattatatt atataattaa ttatatgtaa 600  
 ttaatataat ataattatta tatataattt ttatatatat ataatatata attatataat 660  
 ataatatatat tatattatat tatataatat atatattatta tataatataa tataattata 720  
 ttatatatt atataatat atataattat atattatatt atataataaa tataattata 780  
 taatataata tgattatata atatattatg tatattatat attatatatt gtattatgta 840  
 tattatatat tatatattat gtatattata tattatgtat attatatatt atgtatatta 900  
 tatattatat attatattat gtataatata ttatgtatgt tatatatatat ataaattata 960  
 ttatatatta tgtatattat atataaatta tattatatat tatgtatatt atatataata 1020  
 taaagtatat attatgtata ttatatataa tataaagtat atattatgta tattatatat 1080  
 aatatataagt atatattatg tatattatat ataatatataa gtatatatta tgtatattat 1140

## SEL PCT 012.ST25

atataatata aagtatatat tatgtatatt atatataata taaagtatat attatgtata 1200  
 ttatatataa tataaagtat atattatgta tattatatat aatataaagt atatattatg 1260  
 tatattatat ataataataa gtatatatta tgtatattat atataatata aagtatatat 1320  
 tatgtatatt atatataata taaagtatat attatgtata ttatatataa tataaagtat 1380  
 atattatgta tattatatat aatataaagt atatattata tgtataaat tatatattgt 1440  
 tatatattat at 1452

<210> 220

<211> 502

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(502)

<223> MAR of chromosome 2 genomic contig; 56350637..56351138

<400> 220

atatattata gaaatataaa tatatagata tatctatata ttatagaaat ataaatatat 60  
 agatatatct atatattata gaaatataaa tatatagata tatctatata ttatagaaat 120  
 ataaatatat agatatacct atatattata gaaatataaa tatatagata tacctatata 180  
 ttatagaaat ataaatatat agatatacct atatattata gaaatataaa tatatagata 240  
 tatctatata ttatagaaat ataaatatat agatatatct atatattata gaaatataaa 300  
 tatatagata tatctatata ttatagaaat ataaatatat agatatatct atatattata 360  
 gaaatataaa tatatagata tatctatata ttatagaaat ataaatatat agatatatc 420  
 aacatatatg ttacatatta tatattatat atctatatat ctatataaca ttatatatct 480

SEL PCT 012.ST25  
502

atatatctat ataacatata ta

<210> 221

<211> 794

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(794)

<223> MAR of chromosome 2 genomic contig; 57051633..57052426

<400> 221

aactatatat actatattat atagttatac tatatatact atattatata gttatataac	60
tattatataa ctgtattata tagttatata actattatat aactgtatta tatagttata	120
taactattat ataactgtat tatatagtta tataactata ttatataact gtgttatata	180
gttatatatt atataactat attatataac tgtattatat agttatatat tatataacta	240
tattatataa ctgtattata tagttatata ttatataact atattatata actgtattat	300
atagttatat attatataac tgtattatat agttataaaa ctatattata taactgtatt	360
atatagttat aaaactacta tataactgta ttatataatt ataaaattat actatataac	420
tgtattatat agttataaaa ctatactata taactgtatt atagttat aaaactatac	480
tatataactg tattatatag ttataaagct atactatata actgtattat atagttatat	540
aactatacta tataactgta ttatatagtt ataaaactat actatataac tgtattatat	600
agttataaaa ttatattata taactgtatt atagttat ataaactatat tatataactg	660
tattatatag ttatataact atattatata agtgtattat atagttatat aactatatta	720
tataactgta ttatacagtt atataactat attatataac tgtattatat acttatataa	780

SEL PCT 012.ST25  
794

ctatattata taac

&lt;210&gt; 222

&lt;211&gt; 300

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(300)

&lt;223&gt; MAR of chromosome 2 genomic contig; 57069272..57069571

&lt;400&gt; 222

acacatacat atatgtatat atgcacacac atatatatgt atatatacac atacatatat 60

gtatatatac atatatgtat atacgcacat acatatatgt atatatacac gtacatatat 120

gtctctatat atacacatac acatatgtat atacatatat gtgtatatat acacaatcat 180

atatgtatat acatatatac acatatacac aaacatatat gtatatacat atatgtatat 240

acatatatac acatatacac aaacatatat gtatatacat atatgtatat acatacacia 300

&lt;210&gt; 223

&lt;211&gt; 370

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(370)

## SEL PCT 012.ST25

<223> MAR of chromosome 2 genomic contig; 57235143..57235512

<400> 223

```
tattttata tataactata tatatttat atataaatta tatatatgat catatatata    60
atcatatata taatcatata tgattatata tgatcatata tatatttata tatataatta   120
tatatactta tatataatta tatatatatt tatatatata attatgtata cttatatata   180
tttatatata taattatata tacaatttat atatataatt atatataatt tatatataat   240
tatatatata aattatatat aagtatatat aattatatat atgtttatat ataattatat   300
atataaatga tatgtataat atataactat atataattat atataaatat atatatagat   360
tttatatata                                     370
```

<210> 224

<211> 306

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(306)

<223> MAR of chromosome 2 genomic contig; 57693125..57693430

<400> 224

```
tacgtatata cacgtataaa tataaatata tacatgtata tacgtatata catgtataaa    60
tataaatata tatatgtata tacgtatata catgtataaa tatatatatg tatatacgta   120
tatacatgta taaatatata tatatgtata tacgtatata catgtataaa tatatataca   180
tgtatatacg tatgttgtgt atacatacaa atctgtacat atatacatat atgttgtgtg   240
```

## SEL PCT 012.ST25

tatatataca tctatacatg tgtatgcgta tatatgtata tgtatatata gtatatataa 300

tacatg 306

<210> 225

<211> 500

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(500)

<223> MAR of chromosome 2 genomic contig; 59810331..59810830

<400> 225

tttattatat gtaatatata ttgtattatt atatatatta tatataatat atattgtatt 60

attatatata ttatatataa tatatattgt attattatat atattatata taatatatat 120

tgtattatta tatatattat atataatata tattgtatat tatatatatt atatattata 180

ttattatata ttatatatat tatattatta tatattatat attatatata ttatattata 240

tattatatat tatattatat atattatatt atatattata tattatatta tatatattat 300

attatatatt atatattata ttatatatat tatattatat atattatata ttatatatta 360

tatatattat atattatata ttatatatat tatatattat atataatata tattatatta 420

ttatataata ttatatatta tatatattat atattatata taatatatat tatattatta 480

tataatatta tatattatat 500

<210> 226

<211> 565

<212> DNA

## SEL PCT 012.ST25

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(565)

<223> MAR of chromosome 2 genomic contig; 59974589..59975153

<400> 226

```
atatatgtat aatatgtata tatgtatata ttatgtatat gttatatatg taatatatgt   60
atgtatatat tatatatcat atataatata taatgtgtat atatgtatat atgtatgtat   120
acatgtatat actatgtata tattgtatat attatatatg tatatataca tatacatata   180
taatatatac atatattata tacaatatat acatgtatat tatatacgat atatacatat   240
atattatata caatatatac atagtatata aatgtataca tacatacata tatacatatt   300
atatatgtat atatgtatac ataaatgtat atataatata tatacatata taaatgtata   360
catacgtaca tatacgtata tgtatatgca tatatgtata tatgtgcata catatatatg   420
tatatacata tatgtacata tgtacatata cgtatatatg tacatatgta catatacgta   480
tatatgtaca tatgtacata tacgtatata tgtacatatg tacatatacg tatatatgta   540
catatgtaca tatatacata tatat                                     565
```

<210> 227

<211> 427

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

## SEL PCT 012.ST25

&lt;222&gt; (1)..(427)

&lt;223&gt; MAR of chromosome 2 genomic contig; 60605573..60605999

&lt;400&gt; 227

tatataatgt atataatgga tatagatata gatatagata tatatffffat ataatatata 60

ttatatatta tatataatat atgttatata tattatatat ttatatatat atatatatta 120

tataaattat atatatataa tatataatat atatattata tatatffffat ataatatata 180

ttatatatta tctattatat atffffatata atatatattt tatataatat ataatatata 240

atatatatatt tacataatat ataatatata atacgtatta tatataatat ataatacgta 300

tttatataa tatataatac gtattatata taatacgta tatatattat ataatatata 360

atacgtatta tataatatac gtaattatat ttattataa tacgtattat atattatata 420

atatata 427

&lt;210&gt; 228

&lt;211&gt; 1199

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(1199)

&lt;223&gt; MAR of chromosome 2 genomic contig; 61229949..61231147

&lt;400&gt; 228

gtatacatat ataaagtgtat tatataatgt atatacatat atacatatat aaagtatata 60

tataatatat acatatataa agtatatata taatatatac atatatataag tatatatatat 120

## SEL PCT 012.ST25

atatacatat ataaagtata tataatata acatatata agtatatata tcatatatac 180  
 atatataaag tatatatata atatatacat atatacatat ataaagtata tataacatat 240  
 atacatatat aaagtatata taacatatat acatatataa agtatatata taatatatac 300  
 atatatacat atataaagta tatataacat atatacatat atacagtata tataacatat 360  
 atacatatat acagtatata taacatatat acatatatac agtatatata acatatatac 420  
 atatatacag tatatataac atatatacat atatacatga agtatatata acatatatac 480  
 atatatacat gaagtatata taacatatat acatatatac atgaagtata tataacatat 540  
 atacatatat acatgaagta tatataacat atatacatat atacatatat aaagtatata 600  
 taacatatac atatatacat atataaagta taacatatac atatatacat atataaagta 660  
 tatataatat ataacatata catatataaa gtatatataa tatataacat atacatatat 720  
 aaagtatata taatatataa catatacata tataaagtat atataatata tacatatata 780  
 catatataaa gtatatataa tatatatata catatataaa gtatatataa tatatatata 840  
 tatatacata tataaagtat atataatata tatacatata taaagtatat ataatatata 900  
 tacatatata catatataaa gtatatataa tatatatata tatatacata tataaagtat 960  
 atataatata tatacatata tacatatata aagtatatat aatatatata catatatata 1020  
 tatataaagt atatataata tatatacata tatacatata taaagtatat ataatatata 1080  
 tacatatata catatataaa gtatatataa tatgtatata tatatacata tataaagtat 1140  
 atataatatg tatacatata tacatatata aagtatatat ataatatgta tacatatat 1199

<210> 229

<211> 454

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

## SEL PCT 012.ST25

&lt;222&gt; (1)..(454)

&lt;223&gt; MAR of chromosome 2 genomic contig; 62181058..62181511

&lt;400&gt; 229

tatatatcat atattatata tgatatatat tatgtatata atacatatta tatataataa 60  
 atatttatta tatatgatat atattatgta tataatacat attatatata ataaatatat 120  
 attatattat atataataaaa tatatattat attatatata atatatattt atatataaat 180  
 atattatata taaatatata ttatatataa aatatttata tattatatat aaatatatat 240  
 tatatataaa tatttatata ttatatataa atatttatat attatatata aatatttata 300  
 tattatatat aaaatatatt atatatatta tatatattat atattatata taatatattt 360  
 aatatataat atataaacat atattatata taatatataa acatatataa atatatttat 420  
 atataataga taaaaatata tataatatat ataa 454

&lt;210&gt; 230

&lt;211&gt; 658

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(658)

&lt;223&gt; MAR of chromosome 2 genomic contig; 62190919..62191576

&lt;400&gt; 230

tatatacaca actatatata taactatata tataacaacta tatatacaac tatatatata 60  
 actatatata taactatata taactatata tataactata tataactata tatataacta 120

## SEL PCT 012.ST25

tatatataac tatataaact tatatatata actatatata actatatata actatatata 180  
 taactatata tataaactata tatataacta tatatataac tatatatatact atatataata 240  
 ctatatatat ataactatat atataactat atatataata ctatatataa ctatatatat 300  
 ataactatat atataaactat atatataata ctatatatat aactatatat atataactat 360  
 atatataact atatatatat aactatatat aactatatat atataactat atatataact 420  
 atatatatat aactatatat ataactatat atatataact atatataata ctatatatat 480  
 ataactatat atataaactat atatataact atatataata ctatatatat ataactatat 540  
 atataactat atatataata ctatatatat aactatatat ataactatat atataactat 600  
 atatataata ctatatatat aactatatat atataactat atatataact atatatat 658

&lt;210&gt; 231

&lt;211&gt; 1486

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(1486)

&lt;223&gt; MAR of chromosome 2 genomic contig; 62384127..62385612

&lt;400&gt; 231

attatatcta atctattata tattatatct aatacatatt atatctaac tattgtatat 60  
 tatatctaata atataatata ttatatataa tatattatat attatatatt atatacaata 120  
 tattatatat tatataatat ataatatatt atatataata tattatatct aatatattac 180  
 atattatatc taatctatta tagatataat atgtaatata ttatatatta tatctaataag 240  
 atattagata taatatataa tatattatta atataatata ttagatataa tatataatat 300

## SEL PCT 012.ST25

aataatatat aatatatatt attggaata tataatatat aattaatat atatattata	360
tataattatt atgaataata tatcatatat aatatctagt atattatata ttaataacat	420
ataaatatta tattaataat aaataacata ttaatattat attaataata tataatatac	480
taatattata ttaataatat ataataact aatattatat taataatata taatatacta	540
atattatatt aataatatat aatatactaa tattatatta ataatatata atatacta	600
attatattaa taatatataa tataactaata tattaagaat atataatata ctaatatatt	660
aagaatatat aatatactaa tattatatta ataatatata ttatattaa taatatatta	720
attattatta attaattatt aataattata taatattgat tatattaata ttatcaattt	780
aataatattg attatatatt atatattata tattatatat tatatattat atattatata	840
ttatatatta ataatatata ttagatataa tataatatat taataatata taagatataa	900
tataatatat taataatata tattagatat aatataatat attaataata tatattagat	960
ataatatat atattaataa tatatattag atataatata atatattaat aatatatatt	1020
agatgtaata taatatatta ataatatata ttagatgtaa tataatatat taataatata	1080
tattagatgt aatataatat attaataata tatattagat gtaatataat atattaataa	1140
tatatattag atgtaatata atatattaat aatatatatt agatgtaata taatatatta	1200
ataatatata ttagatgtaa tataatatat taatatatat tagatgtaat ataatatatt	1260
aataatatat attagatata atataatata ttaataatat attagatata atataatata	1320
ttaataatat ataagatata atataatata ttaataatat ataagatata atataatata	1380
ttaataatat ataagatata atataatata ttaataatat atattagata tataatatat	1440
taataatata tattagatat ctaatatcta ttagatatct aataga	1486

&lt;210&gt; 232

&lt;211&gt; 333

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

## SEL PCT 012.ST25

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(333)

&lt;223&gt; MAR of chromosome 2 genomic contig; 62538649..62538981

&lt;400&gt; 232

ttatatatat tatatatata tttttatat atatattata tatatatatt atatatatat 60

tatatatata ttttatatat atattatata tatattttat atatatatta tatatatatt 120

ttatatatat tatatatata ttttatatat attatatata tttttatat atatattata 180

tatatatatt atatatatat tatatatata ttttatatat atattatata tatattttat 240

atatatatta tatatatatt ttatatatat attatatata tttttatat atatattata 300

tatatatatt atatatatat tatatatata ttt 333

&lt;210&gt; 233

&lt;211&gt; 480

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(480)

&lt;223&gt; MAR of chromosome 2 genomic contig; 63240325..63240804

&lt;400&gt; 233

tatatatata atatatatatt tttaaataata aaatatatat atattttaat attaatatat 60

atatatttta atatataata tatatatatt atattttata tataaaatat atatattata 120

## SEL PCT 012.ST25

tattttatat ataaaatata tatattatat attttatata ttaaaatata tattttatat 180  
 attttaatta ttaaaatata tatattatat attttaata taaaatatat atattatata 240  
 tttaatatata taaaatatat atattatata tttaatatata taaaatatat atattttata 300  
 ttatatata taaatatata tattatatat tttaatatat aaaatatata tattatatat 360  
 tttaatatat aaaatatata tattatatat tttaatatat aaaatatata tattatatat 420  
 tttaatatat ataaaatata tatattatat attttatata tattaaatat atattttata 480

<210> 234

<211> 302

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(302)

<223> MAR of chromosome 2 genomic contig; 63935480..63935781

<400> 234

atatatataa atatatatat aattatatat agatatatat aattatatat agatatatat 60  
 attctatatt ctatatatat ataatatata atatataaat tatatataga atatatatta 120  
 tatataatat attatatata ttatatataa tatatatatt atatatatta tatataatat 180  
 atatattata tatattatat ataatttata tatattatat atagaatatata tattatatat 240  
 agaatataga atatatatata tatatataga atacagaata tatatagaat atagaatatata 300  
 ta 302

<210> 235

<211> 407

## SEL PCT 012.ST25

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(407)

<223> MAR of chromosome 2 genomic contig; 63935888..63936294

<400> 235

tataatatat taatataata tatagacagt atataatata atatacagac agtatataat 60  
atacagacag tatataatat ataattattat atataatatt atatataata ttatataata 120  
tattatatta tatatattat ataattattat atattatata taatatatgt aatattatat 180  
attatattat acataatata ttatatataa tatattatat ataattattat atatattata 240  
tataatatat ataataataa tattataata tataatatat aatagtacag tatatattat 300  
atatataatt ctatatataa tatatagaat tctatctatt tataatatat atagaattct 360  
atatataata tataatatatc agaattctat atatattata tatagaa 407

<210> 236

<211> 302

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(302)

<223> MAR of chromosome 2 genomic contig; 66958350..66958651

## SEL PCT 012.ST25

<400> 236  
 tattatatat attgtatata tatgtatatatt atatatattg tatatataat gtatattata 60  
 tatatttatat atatatgtat attatatata ttgtatatat atgtatatta tatatattgt 120  
 atatatgtat atgtatatat gtatgtgtat atatatacac atatacacat atatgtgtat 180  
 gtatatatat gtgtgtatat acgtatatat acatatatac aatttttgta tatatacata 240  
 tatacacata tatatgtgta tgtgtatata tatacacata tatgtgtgtg tatatacaca 300  
 ta 302

<210> 237  
 <211> 651  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_binding  
 <222> (1)..(651)  
 <223> MAR of chromosome 2 genomic contig; 68307125..68307775

<400> 237  
 gatatttatat attgtatata ttatatatgt atataatata ctattatata ttatatatgt 60  
 atataatttt attaatatat atatttatatt atattatata ttatattata ttatattata 120  
 tatataatat taatattata tattattata tattatatta tattaatatt atatatatat 180  
 aatatatata atatatataa tagtattata tataatatat ataatagtat tatatattat 240  
 atatatataa tactattata tatatttatat ataatagtat tatatatatt atatatataa 300  
 tactattata tataatatat actatttatat aatatatata atactattat atatattata 360

## SEL PCT 012.ST25

tataatacta ttatatataa tatatatataat actattatat ataatatata taatactatt 420  
 atatataata tatataatac tattatatat aatatatata atactattat atataatata 480  
 tataatacta ttatatataa tatatatatt atatataatt atattaatat ataatagtat 540  
 catatataat aatagtatat ataatatata atatatatat tatatatatt ataatagtat 600  
 atataacata taatatagta tatatatatt atattatata taaaatattt a 651

<210> 238

<211> 367

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_binding

<222> (1)..(367)

<223> MAR of chromosome 2 genomic contig; 68308243..68308609

<400> 238

atatatatat atgagtcac catacacata tatatatata atgtttatat atataatgta 60  
 tatatatataat gtttatatat aatgtatata tataatgttt atatatatataa tgtatatata 120  
 taatgtttat atatataatg tatatatata atgtttatat atataatgtg tatatatataat 180  
 gtttatatat ataatgtgta tatataatgt ttatatataa tgggtatata taatgtttat 240  
 atatataatg tgtatatata atgtttatat atataatgtg tatatatataat gtttatatat 300  
 ataatgtgta tatatatataat gtttatatat ataatgtgta tatatatataat gtttatatat 360  
 ataatgt 367

<210> 239

<211> 499

## SEL PCT 012.ST25

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(499)

&lt;223&gt; MAR of chromosome 2 genomic contig; 410241..410739

&lt;400&gt; 239

ataatatgta tatatattat attatatatt atattacata ttatatatta tattacatat 60

tatatattta tatattacat attatatatt atattttata ttatatatta tatcatatat 120

atgttatgca ttatataata cataatatat tatatatgat ataatatata ttatatatta 180

ttatatataa tataattaat atattatgta ttatataata tatattatgt tataatatat 240

aatatatatt atataattat ataatatatt atgtattata taatatatat tatgttataa 300

tatattatat tatatatatt atatatatat tatatatata atgtatatta tatataatac 360

ataatatatt atatattata tattatttta tataatatat tatataatgt gatattattat 420

ataatatatt atataacata gtatattata taatatatta tataatgtaa tatattatat 480

attatataat atattgtat 499

&lt;210&gt; 240

&lt;211&gt; 402

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

## SEL PCT 012.ST25

&lt;222&gt; (1)..(402)

&lt;223&gt; MAR of chromosome 2 genomic contig; 31531..31932

&lt;400&gt; 240

cacattatat atataaacat tatatatata cacattatat atataaacat tatatatata 60  
 cacattatat atataaacat tatatatata cattatatat ataaacatta tatatacaca 120  
 ttatatatat aaacattata tatacaaatt atatataata acattatata tacaattat 180  
 atatataaac attatatata tacattatat atataaacat tatatatata cattatatat 240  
 ataaacatta tatatatata ttatatatat aaacattata tatatacatt atatataata 300  
 acattatata tacaattat atatataaac gttatatata tacattatat atataaacat 360  
 tatatgtata cattatatat ataaacatta tatatatatg tg 402

&lt;210&gt; 241

&lt;211&gt; 421

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_binding

&lt;222&gt; (1)..(421)

&lt;223&gt; MAR of chromosome 2 genomic contig; 32415..32835

&lt;400&gt; 241

ataaatattt tatatataat atataatata tatactatat tatatgttat atatactatt 60  
 ataatatata taatatatat attatatatt atatatacta ttattatata tgatactatt 120  
 atatattaat ataattatat ataatatata tattatataa tatactatta tatattatat 180

## SEL PCT 012.ST25

ataatagtat attatataat atatatatta tatataatag tattatatat actattatat 240

attatatata ttatatatat ataaaatata atataatata tataatatat aatattaata 300

ttatatatat aatataatat aatatataat ataataatata atatatatta ataaaattat 360

attaatatat aatatataat agtatattat atacatatat aatatatata atataata 420

t 421